

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets



(11)

EP 1 365 034 A2

(12)

EUROPEAN PATENT APPLICATION

(43) Date of publication:
26.11.2003 Bulletin 2003/48

(51) Int Cl.⁷: **C12Q 1/68**, B01L 3/00,
A61K 48/00, G06F 17/00

(21) Application number: **03010447.5**

(22) Date of filing: **09.05.2003**

(84) Designated Contracting States:
**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR
HU IE IT LI LU MC NL PT RO SE SI SK TR**
Designated Extension States:
AL LT LV MK

(30) Priority: **21.05.2002 EP 02010291**
13.02.2003 EP 03003112

(71) Applicant: **Bayer Aktiengesellschaft**
51368 Leverkusen (DE)

(72) Inventors:
• **Wirtz, Ralph, Dr.**
50674 Köln (DE)
• **Munnes, Marc, Dr.**
40699 Erkrath (DE)
• **Kallabis, Harald, Dr.**
50161 Köln (DE)

(54) **Methods and compositions for the prediction, diagnosis, prognosis, prevention and treatment of malignant neoplasia**

(57) The invention provides compositions, methods and uses, for the prediction., diagnosis, prognosis, prevention and treatment of malignant neoplasia and breast cancer in particular. Genes that are differentially

expressed in breast tissue of breast cancer patients versus those of normal people are disclosed.

EP 1 365 034 A2

Description**TECHNICAL FIELD OF THE INVENTION**

5 [0001] The invention relates to methods and compositions for the prediction, diagnosis, prognosis, prevention and treatment of neoplastic disease. Neoplastic disease is often caused by chromosomal rearrangements which lead to over- or underexpression of the rearranged genes. The invention discloses genes which are overexpressed in neoplastic tissue and are useful as diagnostic markers and targets for treatment. Methods are disclosed for predicting, diagnosing and prognosing as well as preventing and treating neoplastic disease.

BACKGROUND OF THE INVENTION

10 [0002] Chromosomal aberrations (amplifications, deletions, inversions, insertions, translocations and/or viral integrations) are of importance for the development of cancer and neoplastic lesions, as they account for deregulations of the respective regions. Amplifications of genomic regions have been described, in which genes of importance for growth characteristics, differentiation, invasiveness or resistance to therapeutic intervention are located. One of those regions with chromosomal aberrations is the region carrying the HER-2/neu gene which is amplified in breast cancer patients. In approximately 25% of breast cancer patients the HER-2/neu gene is overexpressed due to gene amplification. HER-2/neu overexpression correlates with a poor prognosis (relapse, overall survival, sensitivity to therapeutics). The importance of HER-2/neu for the prognosis of the disease progression has been described [Gusterson et al., 1992, (1)]. Gene specific antibodies raised against HER-2/neu (Herceptin™) have been generated to treat the respective cancer patients. However, only about 50% of the patients benefit from the antibody treatment with Herceptin™, which is most often combined with chemotherapeutic regimen. The discrepancy of HER-2/neu positive tumors (overexpressing HER-2/neu to similar extent) with regard to responsiveness to therapeutic intervention suggest, that there might be additional factors or genes being involved in growth and apoptotic characteristics of the respective tumor tissues. There seems to be no monocausal relationship between overexpression of the growth factor receptor HER-2/neu and therapy outcome. In line with this the measurement of commonly used tumor markers such as estrogen receptor, progesterone receptor, p53 and Ki-67 do provide only very limited information on clinical outcome of specific therapeutic decisions. Therefore there is a great need for a more detailed diagnostic and prognostic classification of tumors to enable improved therapy decisions and prediction of survival of the patients. The present invention addresses the need for additional markers by providing genes, which expression is deregulated in tumors and correlates with clinical outcome. One focus is the deregulation of genes present in specific chromosomal regions and their interaction in disease development and drug responsiveness.

30 [0003] HER-2/neu and other markers for neoplastic disease are commonly assayed with diagnostic methods such as immunohistochemistry (IHC) (e.g. HercepTest™ from DAKO Inc.) and Fluorescence-In-Situ-Hybridization (FISH) (e.g. quantitative measurement of the HER-2/neu and Topoisomerase II alpha with a fluorescence-*in-situ*-Hybridization kit from VYSIS). Additionally HER-2/neu can be assayed by detecting HER-2/neu fragments in serum with an ELISA test (BAYER Corp.) or with a quantitative PCR kit which compares the amount of HER-2/neu gene with the amount of a non-amplified control gene in order to detect HER-2/neu gene amplifications (ROCHE). These methods, however, exhibit multiple disadvantages with regard to sensitivity, specificity, technical and personnel efforts, costs, time consumption, inter-lab reproducibility. These methods are also restricted with regard to measurement of multiple parameters within one patient sample ("multiplexing"). Usually only about 3 to 4 parameters (e.g. genes or gene products) can be detected per tissue slide. Therefore, there is a need to develop a fast and simple test to measure simultaneously multiple parameters in one sample. The present invention addresses the need for a fast and simple high-resolution method, that is able to detect multiple diagnostic and prognostic markers simultaneously.

SUMMARY OF THE INVENTION

50 [0004] The present invention is based on discovery that chromosomal alterations in cancer tissues can lead to changes in the expression of genes that are encoded by the altered chromosomal regions. Exemplary 43 human genes have been identified that are co-amplified in neoplastic lesions from breast cancer tissue resulting in altered expression of several of these genes (Tables 1 to 4). These 43 genes are differentially expressed in breast cancer states, relative to their expression in normal, or non-breast cancer states. The present invention relates to derivatives, fragments, analogues and homologues of these genes and uses or methods of using of the same.

55 [0005] The present invention further relates to novel preventive, predictive, diagnostic, prognostic and therapeutic compositions and uses for malignant neoplasia and breast cancer in particular. Especially membrane bound marker gene products containing extracellular domains can be a particularly useful target for treatment methods as well as diagnostic and clinical monitoring methods.

[0006] It is a discovery of the present invention that several of these genes are characterized in that their gene products functionally interact in signaling cascades or by directly or indirectly influencing each other. This interaction is important for the normal physiology of certain non-neoplastic tissues (e.g. brain or neurogenic tissue). The deregulation of these genes in neoplastic lesions where they are normally exhibit of different level of activity or are not active, however, results in pathophysiology and affects the characteristics of the disease-associated tissue.

[0007] The present invention further relates to methods for detecting these deregulations in malignant neoplasia on DNA and mRNA level.

[0008] The present invention further relates to a method for the detection of chromosomal alterations characterized in that the relative abundance of individual mRNAs, encoded by genes, located in altered chromosomal regions is detected.

[0009] The present invention further relates to a method for the detection of the flanking breakpoints of named chromosomal alterations by measurement of DNA copy number by quantitative PCR or DNA-Arrays and DNA sequencing.

[0010] A method for the prediction, diagnosis or prognosis of malignant neoplasia by the detection of DNA sequences flanking named genomic breakpoint or are located within such.

[0011] The present invention further relates to a method for the detection of chromosomal alterations characterized in that the copy number of one or more genomic nucleic acid sequences located within an altered chromosomal region (s) is detected by quantitative PCR techniques (e.g. TaqMan™, Lightcycler™ and iCycler™).

[0012] The present invention further relates to a method for the prediction, diagnosis or prognosis of malignant neoplasia by the detection of at least 2 markers whereby the markers are genes and fragments thereof or genomic nucleic acid sequences that are located on one chromosomal region which is altered in malignant neoplasia and breast cancer in particular.

[0013] The present invention also discloses a method for the prediction, diagnosis or prognosis of malignant neoplasia by the detection of at least 2 markers whereby the markers are located on one or more chromosomal region(s) which is/are altered in malignant neoplasia; and the markers interact as (i) receptor and ligand or (ii) members of the same signal transduction pathway or (iii) members of synergistic signal transduction pathways or (iv) members of antagonistic signal transduction pathways or (v) transcription factor and transcription factor binding site.

[0014] Also disclosed is a method for the prediction, diagnosis or prognosis of malignant neoplasia by the detection of at least one marker whereby the marker is a VNTR, SNP, RFLP or STS which is located on one chromosomal region which is altered in malignant neoplasia due to amplification and the marker is detected in (a) a cancerous and (b) a non cancerous tissue or biological sample from the same individual. A preferred embodiment is the detection of at least one VNTR marker of Table 6 or at least on SNP marker of Table 4 or combinations thereof.. Even more preferred can the detection, quantification and sizing of such polymorphic markers be achieved by methods of (a) for the comparative measurement of amount and size by PCR amplification and subsequent capillary electrophoresis, (b) for sequence determination and allelic discrimination by gel electrophoresis (e.g. SSCP, DGGE), real time kinetic PCR, direct DNA sequencing, pyro-sequencing, mass-specific allelic discrimination or resequencing by DNA array technologies, (c) for the dertermination of specific restriction patterns and subsequent electrophoretic separation and (d) for allelic discrimination by allele specific PCR (e.g. ASO). An even more favorable detection of a hetrozygous VNTR, SNP, RFLP or STS is done in a multiplex fashion, utilizing a variety of labeled primers (e.g. fluorescent, radioactive, bioactive) and a suitable capillary electrophoresis (CE) detection system.

[0015] In another embodiment the expression of these genes can be detected with DNA-arrays as described in WO9727317 and US6379895.

[0016] In a further embodiment the expression of these genes can be detected with bead based direct flourescent readout techniques such as described in WO9714028 and WO9952708.

[0017] In one embodiment, the invention pertains to a method of determining the phenotype of a cell or tissue, comprising detecting the differential expression, relative to a normal or untreated cell, of at least one polynucleotide comprising SEQ ID NO: 2 to 6, 8, 9, 11 to 16, 18, 19 or 21 to 26 or 53 to 75, wherein the polynucleotide is differentially expressed by at least about 1.5 fold, at least about 2 fold or at least about 3 fold.

[0018] In a further aspect the invention pertains to a method of determining the phenotype of a cell or tissue, comprising detecting the differential expression, relative to a normal or untreated cell, of at least one polynucleotide which hybridizes under stringent conditions to one of the polynucleotides of SEQ ID NO: 2 to 6, 8, 9, 11 to 16, 18, 19 or 21 to 26 or 53 to 75 and encodes a polypeptide exhibiting the same biological function as given in Table 2 or 3 for the respective polynucleotide, wherein the polynucleotide is differentially expressed by at least at least about 1.5 fold , at least about 2 fold or at least about 3 fold.

[0019] In another embodiment of the invention a polynucleotide comprising a polynucleotide selected from SEQ ID NO: 2 to 6, 8, 9, 11 to 16, 18, 19 or 21 to 26 and 53 to 75 or encoding one of the polypeptides with SEQ ID NO: 28 to 32, 34, 35, 37 to 42, 44, 45 or 47 to 52 or 76 to 98 can be used to identify cells or tissue in individuals which exhibit a phenotype predisposed to breast cancer or a diseased phenotype, thereby (a) predicting whether an individual is at risk for the development, or (b) diagnosing whether an individual is having, or (c) prognosing the progression or the

outcome of the treatment malignant neoplasia and breast cancer in particular.

[0020] In yet another embodiment the invention provides a method for identifying genomic regions which are altered on the chromosomal level and encode genes that are linked by function and are differentially expressed in malignant neoplasia and breast cancer in particular.

[0021] In yet another embodiment the invention provides the genomic regions 17q12, 3p21 and 12q13 for use in prediction, diagnosis and prognosis as well as prevention and treatment of malignant neoplasia and breast cancer. In particular not only the intragenic regions, but also intergenic regions, pseudogenes or non-transcribed genes of said chromosomal regions can be used for diagnostic, predictive, prognostic and preventive and therapeutic compositions and methods.

[0022] In yet another embodiment the invention provides methods of screening for agents which regulate the activity of a polypeptide comprising a polypeptide selected from SEQ ID NO: 27 to 52 and 76 to 98 or encoded by a polynucleotide comprising a polynucleotide selected from SEQ ID NO: 1 to 26 and 53 to 75. A test compound is contacted with a polypeptide comprising a polypeptide selected from SEQ ID NO: 27 to 52 and 76 to 98 or encoded by a polynucleotide comprising a polynucleotide selected from SEQ ID NO: 1 to 26 and 53 to 75. Binding of the test compound to the polypeptide is detected. A test compound which binds to the polypeptide is thereby identified as a potential therapeutic agent for the treatment of malignant neoplasia and more particularly breast cancer.

[0023] In even another embodiment the invention provides another method of screening for agents which regulate the activity of a polypeptide comprising a polypeptide selected from SEQ ID NO: 27 to 52 and 76 to 98 or encoded by a polynucleotide comprising a polynucleotide selected from SEQ ID NO: 1 to 26 and 53 to 75. A test compound is contacted with a polypeptide comprising a polypeptide selected from SEQ ID NO: 27 to 52 and 76 to 98 or encoded by a polynucleotide comprising a polynucleotide selected from SEQ ID NO: 1 to 26 and 53 to 75. A biological activity mediated by the polypeptide is detected. A test compound which decreases the biological activity is thereby identified as a potential therapeutic agent for decreasing the activity of the polypeptide encoded by a polypeptide comprising a polypeptide selected from SEQ ID NO: 27 to 52 and 76 to 98 or encoded by a polynucleotide comprising a polynucleotide selected from SEQ ID NO: 1 to 26 and 53 to 75 in malignant neoplasia and breast cancer in particular. A test compound which increases the biological activity is thereby identified as a potential therapeutic agent for increasing the activity of the polypeptide encoded by a polypeptide selected from one of the polypeptides with SEQ ID NO: 27 to 52 and 76 to 98 or encoded by a polynucleotide comprising a polynucleotide selected from SEQ ID NO: 1 to 26 and 53 to 75 in malignant neoplasia and breast cancer in particular.

[0024] In another embodiment the invention provides a method of screening for agents which regulate the activity of a polynucleotide comprising a polynucleotide selected from SEQ ID NO: 1 to 26 and 53 to 75. A test compound is contacted with a polynucleotide comprising a polynucleotide selected from SEQ ID NO: 1 to 26 and 53 to 75. Binding of the test compound to the polynucleotide comprising a polynucleotide selected from SEQ ID NO: 1 to 26 and 53 to 75 is detected. A test compound which binds to the polynucleotide is thereby identified as a potential therapeutic agent for regulating the activity of a polynucleotide comprising a polynucleotide selected from SEQ ID NO: 1 to 26 and 53 to 75 in malignant neoplasia and breast cancer in particular.

[0025] The invention thus provides polypeptides selected from one of the polypeptides with SEQ ID NO: 27 to 52 and 76 to 98 or encoded by a polynucleotide comprising a polynucleotide selected from SEQ ID NO: 1 to 26 and 53 to 75 which can be used to identify compounds which may act, for example, as regulators or modulators such as agonists and antagonists, partial agonists, inverse agonists, activators, co-activators and inhibitors of the polypeptide comprising a polypeptide selected from SEQ ID NO: 27 to 52 and 76 to 98 or encoded by a polynucleotide comprising a polynucleotide selected from SEQ ID NO: 1 to 26 and 53 to 75. Accordingly, the invention provides reagents and methods for regulating a polypeptide comprising a polypeptide selected from SEQ ID NO: 27 to 52 and 76 to 98 or encoded by a polynucleotide comprising a polynucleotide selected from SEQ ID NO: 1 to 26 and 53 to 75 in malignant neoplasia and more particularly breast cancer. The regulation can be an up-or down regulation. Reagents that modulate the expression, stability or amount of a polynucleotide comprising a polynucleotide selected from SEQ ID NO: 1 to 26 and 53 to 75 or the activity of the polypeptide comprising a polypeptide selected from SEQ ID NO: 27 to 52 and 76 to 98 or encoded by a polynucleotide comprising a polynucleotide selected from SEQ ID NO: 1 to 26 and 53 to 75 can be a protein, a peptide, a peptidomimetic, a nucleic acid, a nucleic acid analogue (e.g. peptide nucleic acid, locked nucleic acid) or a small molecule. Methods that modulate the expression, stability or amount of a polynucleotide comprising a polynucleotide selected from SEQ ID NO: 1 to 26 and 53 to 75 or the activity of the polypeptide comprising a polypeptide selected from SEQ ID NO: 27 to 52 and 76 to 98 or encoded by a polynucleotide comprising a polynucleotide selected from SEQ ID NO: 1 to 26 and 53 to 75 can be gene replacement therapies, antisense, ribozyme and triplex nucleic acid approaches.

[0026] In one embodiment of the invention provides antibodies which specifically bind to a full-length or partial polypeptide comprising a polypeptide selected from SEQ ID NO: 27 to 52 and 76 to 98 or encoded by a polynucleotide comprising a polynucleotide selected from SEQ ID NO: 1 to 26 and 53 to 75 or a polynucleotide comprising a polynucleotide selected from SEQ ID NO: 1 to 26 and 53 to 75 for use in prediction, prevention, diagnosis, prognosis and

treatment of malignant neoplasia and breast cancer in particular.

[0027] Yet another embodiment of the invention is the use of a reagent which specifically binds to a polynucleotide comprising a polynucleotide selected from SEQ ID NO: 1 to 26 and 53 to 75 or a polypeptide comprising a polypeptide selected from SEQ ID NO: 27 to 52 and 76 to 98 or encoded by a polynucleotide comprising a polynucleotide selected from SEQ ID NO: 1 to 26 and 53 to 75 in the preparation of a medicament for the treatment of malignant neoplasia and breast cancer in particular.

[0028] Still another embodiment is the use of a reagent that modulates the activity or stability of a polypeptide comprising a polypeptide selected from SEQ ID NO: 27 to 52 and 76 to 98 or encoded by a polynucleotide comprising a polynucleotide selected from SEQ ID NO: 1 to 26 and 53 to 75 or the expression, amount or stability of a polynucleotide comprising a polynucleotide selected from SEQ ID NO: 1 to 26 and 53 to 75 in the preparation of a medicament for the treatment of malignant neoplasia and breast cancer in particular.

[0029] Still another embodiment of the invention is a pharmaceutical composition which includes a reagent which specifically binds to a polynucleotide comprising a polynucleotide selected from SEQ ID NO: 1 to 26 and 53 to 75 or a polypeptide comprising a polypeptide selected from SEQ ID NO: 27 to 52 and 76 to 98 or encoded by a polynucleotide comprising a polynucleotide selected from SEQ ID NO: 1 to 26 and 53 to 75, and a pharmaceutically acceptable carrier.

[0030] Yet another embodiment of the invention is a pharmaceutical composition including a polynucleotide comprising a polynucleotide selected from SEQ ID NO: 1 to 26 and 53 to 75 or encoding a polypeptide comprising a polypeptide selected from SEQ ID NO: 27 to 52 and 76 to 98.

[0031] In one embodiment, a reagent which alters the level of expression in a cell of a polynucleotide comprising a polynucleotide selected from SEQ ID NO: 1 to 26 and 53 to 75 or encoding a polypeptide comprising a polypeptide selected from SEQ ID NO: 27 to 52 and 76 to 98, or a sequence complementary thereto, is identified by providing a cell, treating the cell with a test reagent, determining the level of expression in the cell of a polynucleotide comprising a polynucleotide selected from SEQ ID NO: 1 to 26 and 53 to 75 or encoding a polypeptide comprising a polypeptide selected from SEQ ID NO: 27 to 52 and 76 to 98 or a sequence complementary thereto, and comparing the level of expression of the polynucleotide in the treated cell with the level of expression of the polynucleotide in an untreated cell, wherein a change in the level of expression of the polynucleotide in the treated cell relative to the level of expression of the polynucleotide in the untreated cell is indicative of an agent which alters the level of expression of the polynucleotide in a cell.

[0032] The invention further provides a pharmaceutical composition comprising a reagent identified by this method.

[0033] Another embodiment of the invention is a pharmaceutical composition which includes a polypeptide comprising a polypeptide selected from SEQ ID NO: 27 to 52 and 76 to 98 or which is encoded by a polynucleotide comprising a polynucleotide selected from SEQ ID NO: 1 to 26 and 53 to 75.

[0034] A further embodiment of the invention is a pharmaceutical composition comprising a polynucleotide including a sequence which hybridizes under stringent conditions to a polynucleotide comprising a polynucleotide selected from SEQ ID NO: 1 to 26 and 53 to 75 and encoding a polypeptide exhibiting the same biological function as given for the respective polynucleotide in Table 2 or 3, or encoding a polypeptide comprising a polypeptide selected from SEQ ID NO: 27 to 52 and 76 to 98. Pharmaceutical compositions, useful in the present invention may further include fusion proteins comprising a polypeptide comprising a polynucleotide selected from SEQ ID NO: 27 to 52 and 76 to 98, or a fragment thereof, antibodies, or antibody fragments

BRIEF DESCRIPTION OF THE DRAWINGS

[0035]

Fig. 1 shows a sketch of the chromosome 17 with G-banding pattern and cytogenetic positions. In the blow out at the lower part of the figure a detailed view of the chromosomal area of the long arm of chromosome 17 (17q12-21.1) is provided. Each vertical rectangle depicted in medium gray, represents a gene as labeled below or above the individual position. The order of genes depicted in this graph has been deduced from experiments questioning the amplification an over expression and from public available data (e.g. UCSC, NCBI or Ensemble).

Fig. 2 shows the same region as depicted before in Fig. 1 and a cluster representation of the individual expression values measured by DNA-chip hybridization. The gene representing squares are indicated by a dotted line. In the upper part of the cluster representation 4 tumor cell lines, of which two harbor a known HER-2/neu over expression (SKBR3 and AU565), are depicted with their individual expression profiles. Not only the HER-2/neu gene shows a clear over expression but as provided by this invention several other genes with in the surrounding. In the middle part of the cluster representation expression data obtained from immune histochemically characterized tumor samples are presented. Two of the depicted probes show a significant over

expression of genes marked by the white rectangles. For additional information and comparison expression profiles of several non diseased human tissues (RNAs obtained from Clontech Inc.) are provided. Closest relation to the expression profile of HER-2/neu positive tumors displays human brain and neural tissue.

Fig. 3 provides data from DNA amplification measurements by qPCR (e.g. TaqMan). Data indicates that in several analyzed breast cancer cell lines harbor amplification of genes which were located in the previously described region (ARCHEON). Data were displayed for each gene on the x-axis and 40-Ct at the y-axis. Data were normalized to the expression level of GAPDH as seen in the first group of columns.

Fig. 4 represents a graphical overview on the amplified regions and provides information on the length of the individual amplification and over expression in the analyzed tumor cell lines. The length of the amplification and the composition of genes has a significant impact on the nature of the cancer cell and on the responsiveness on certain drugs, as described elsewhere.

DETAILED DESCRIPTION OF THE INVENTION

Definitions

[0036] "Differential expression", as used herein, refers to both quantitative as well as qualitative differences in the genes' expression patterns depending on differential development and/or tumor growth. Differentially expressed genes may represent "marker genes," and/or "target genes". The expression pattern of a differentially expressed gene disclosed herein may be utilized as part of a prognostic or diagnostic breast cancer evaluation. Alternatively, a differentially expressed gene disclosed herein may be used in methods for identifying reagents and compounds and uses of these reagents and compounds for the treatment of breast cancer as well as methods of treatment.

[0037] "Biological activity" or "bioactivity" or "activity" or "biological function", which are used interchangeably, herein mean an effector or antigenic function that is directly or indirectly performed by a polypeptide (whether in its native or denatured conformation), or by any fragment thereof *in vivo* or *in vitro*. Biological activities include but are not limited to binding to polypeptides, binding to other proteins or molecules, enzymatic activity, signal transduction, activity as a DNA binding protein, as a transcription regulator, ability to bind damaged DNA, etc. A bioactivity can be modulated by directly affecting the subject polypeptide. Alternatively, a bioactivity can be altered by modulating the level of the polypeptide, such as by modulating expression of the corresponding gene.

[0038] The term "marker" or "biomarker" refers a biological molecule, e.g., a nucleic acid, peptide, hormone, etc., whose presence or concentration can be detected and correlated with a known condition, such as a disease state.

[0039] "Marker gene," as used herein, refers to a differentially expressed gene which expression pattern may be utilized as part of predictive, prognostic or diagnostic malignant neoplasia or breast cancer evaluation, or which, alternatively, may be used in methods for identifying compounds useful for the treatment or prevention of malignant neoplasia and breast cancer in particular. A marker gene may also have the characteristics of a target gene.

[0040] "Target gene", as used herein, refers to a differentially expressed gene involved in breast cancer in a manner by which modulation of the level of target gene expression or of target gene product activity may act to ameliorate symptoms of malignant neoplasia and breast cancer in particular. A target gene may also have the characteristics of a marker gene.

[0041] The term "biological sample", as used herein, refers to a sample obtained from an organism or from components (e.g., cells) of an organism. The sample may be of any biological tissue or fluid. Frequently the sample will be a "clinical sample" which is a sample derived from a patient. Such samples include, but are not limited to, sputum, blood, blood cells (e.g., white cells), tissue or fine needle biopsy samples, cell-containing bodyfluids, free floating nucleic acids, urine, peritoneal fluid, and pleural fluid, or cells therefrom. Biological samples may also include sections of tissues such as frozen sections taken for histological purposes.

[0042] By "array" or "matrix" is meant an arrangement of addressable locations or "addresses" on a device. The locations can be arranged in two dimensional arrays, three dimensional arrays, or other matrix formats. The number of locations can range from several to at least hundreds of thousands. Most importantly, each location represents a totally independent reaction site. Arrays include but are not limited to nucleic acid arrays, protein arrays and antibody arrays. A "nucleic acid array" refers to an array containing nucleic acid probes, such as oligonucleotides, polynucleotides or larger portions of genes. The nucleic acid on the array is preferably single stranded. Arrays wherein the probes are oligonucleotides are referred to as "oligonucleotide arrays" or "oligonucleotide chips." A "microarray," herein also refers to a "biochip" or "biological chip", an array of regions having a density of discrete regions of at least about 100/cm², and preferably at least about 1000/cm². The regions in a microarray have typical dimensions, e.g., diameters, in the range of between about 10-250 μ m, and are separated from other regions in the array by about the same distance. A "protein array" refers to an array containing polypeptide probes or protein probes which can be in native form or

denatured. An "antibody array" refers to an array containing antibodies which include but are not limited to monoclonal antibodies (e.g. from a mouse), chimeric antibodies, humanized antibodies or phage antibodies and single chain antibodies as well as fragments from antibodies.

[0043] The term "agonist", as used herein, is meant to refer to an agent that mimics or upregulates (e.g., potentiates or supplements) the bioactivity of a protein. An agonist can be a wild-type protein or derivative thereof having at least one bioactivity of the wild-type protein. An agonist can also be a compound that upregulates expression of a gene or which increases at least one bioactivity of a protein. An agonist can also be a compound which increases the interaction of a polypeptide with another molecule, e.g., a target peptide or nucleic acid.

[0044] The term "antagonist" as used herein is meant to refer to an agent that downregulates (e.g., suppresses or inhibits) at least one bioactivity of a protein. An antagonist can be a compound which inhibits or decreases the interaction between a protein and another molecule, e.g., a target peptide, a ligand or an enzyme substrate. An antagonist can also be a compound that downregulates expression of a gene or which reduces the amount of expressed protein present.

[0045] "Small molecule" as used herein, is meant to refer to a composition, which has a molecular weight of less than about 5 kD and most preferably less than about 4 kD. Small molecules can be nucleic acids, peptides, polypeptides, peptidomimetics, carbohydrates, lipids or other organic (carbon-containing) or inorganic molecules. Many pharmaceutical companies have extensive libraries of chemical and/or biological mixtures, often fungal, bacterial, or algal extracts, which can be screened with any of the assays of the invention to identify compounds that modulate a bioactivity.

[0046] The terms "modulated" or "modulation" or "regulated" or "regulation" and "differentially regulated" as used herein refer to both upregulation (i.e., activation or stimulation (e.g., by agonizing or potentiating) and down regulation [i.e., inhibition or suppression (e.g., by antagonizing, decreasing or inhibiting)].

[0047] "Transcriptional regulatory unit" refers to DNA sequences, such as initiation signals, enhancers, and promoters, which induce or control transcription of protein coding sequences with which they are operably linked. In preferred embodiments, transcription of one of the genes is under the control of a promoter sequence (or other transcriptional regulatory sequence) which controls the expression of the recombinant gene in a cell-type in which expression is intended. It will also be understood that the recombinant gene can be under the control of transcriptional regulatory sequences which are the same or which are different from those sequences which control transcription of the naturally occurring forms of the polypeptide.

[0048] The term "derivative" refers to the chemical modification of a polypeptide sequence, or a polynucleotide sequence. Chemical modifications of a polynucleotide sequence can include, for example, replacement of hydrogen by an alkyl, acyl, or amino group. A derivative polynucleotide encodes a polypeptide which retains at least one biological or immunological function of the natural molecule. A derivative polypeptide is one modified by glycosylation, pegylation, or any similar process that retains at least one biological or immunological function of the polypeptide from which it was derived.

[0049] The term "nucleotide analog" refers to oligomers or polymers being at least in one feature different from naturally occurring nucleotides, oligonucleotides or polynucleotides, but exhibiting functional features of the respective naturally occurring nucleotides (e.g. base pairing, hybridization, coding information) and that can be used for said compositions. The nucleotide analogs can consist of non-naturally occurring bases or polymer backbones, examples of which are LNAs, PNAs and Morpholinos. The nucleotide analog has at least one molecule different from its naturally occurring counterpart or equivalent.

[0050] "BREAST CANCER GENES" or "BREAST CANCER GENE" as used herein refers to the polynucleotides of SEQ ID NO: 1 to 26 and 53 to 75, as well as derivatives, fragments, analogs and homologues thereof, the polypeptides encoded thereby, the polypeptides of SEQ ID NO: 27 to 52 and 76 to 98 as well as derivatives, fragments, analogs and homologues thereof and the corresponding genomic transcription units which can be derived or identified with standard techniques well known in the art using the information disclosed in Tables 1 to 5 and Figures 1 to 4. The GenBank, Locuslink ID and the UniGene accession numbers of the polynucleotide sequences of the SEQ ID NO: 1 to 26 and 53 to 75 and the polypeptides of the SEQ ID NO: 27 to 52 and 76 to 98 are shown in Table 1, the gene description, gene function and subcellular localization is given in Tables 2 and 3.

[0051] The term "chromosomal region" as used herein refers to a consecutive DNA stretch on a chromosome which can be defined by cytogenetic or other genetic markers such as e.g. restriction length polymorphisms (RFLPs), single nucleotide polymorphisms (SNPs), expressed sequence tags (ESTs), sequence tagged sites (STSs), micro-satellites, variable number of tandem repeats (VNTRs) and genes. Typically a chromosomal region consists of up to 2 Megabases (MB), up to 4 MB, up to 6 MB, up to 8 MB, up to 10 MB, up to 20 MB or even more MB.

[0052] The term "altered chromosomal region" or "abberant chromosomal region" refers to a structural change of the chromosomal composition and DNA sequence, which can occur by the following events: amplifications, deletions, inversions, insertions, translocations and/or viral integrations. A trisomy, where a given cell harbors more than two copies of a chromosome, is within the meaning of the term "amplification" of a chromosome or chromosomal region.

[0053] The present invention provides polynucleotide sequences and proteins encoded thereby, as well as probes

derived from the polynucleotide sequences, antibodies directed to the encoded proteins, and predictive, preventive, diagnostic, prognostic and therapeutic uses for individuals which are at risk for or which have malignant neoplasia and breast cancer in particular. The sequences disclosure herein have been found to be differentially expressed in samples from breast cancer.

[0054] The present invention is based on the identification of 43 genes that are differentially regulated (up- or down-regulated) in tumor biopsies of patients with clinical evidence of breast cancer. The identification of 43 human genes which were not known to be differentially regulated in breast cancer states and their significance for the disease is described in the working examples herein. The characterization of the co-expression of these genes provides newly identified roles in breast cancer. The gene names, the database accession numbers (GenBank and UniGene) as well as the putative or known functions of the encoded proteins and their subcellular localization are given in Tables 1 to 4. The primer sequences used for the gene amplification are shown in Table 5.

[0055] In either situation, detecting expression of these genes in excess or in with lower level as compared to normal expression provides the basis for the diagnosis of malignant neoplasia and breast cancer. Furthermore, in testing the efficacy of compounds during clinical trials, a decrease in the level of the expression of these genes corresponds to a return from a disease condition to a normal state, and thereby indicates a positive effect of the compound.

[0056] Another aspect of the present invention is based on the observation that neighboring genes within defined genomic regions functionally interact and influence each others function directly or indirectly. A genomic region encoding functionally interacting genes that are co-amplified and co-expressed in neoplastic lesions has been defined as an "ARCHEON". (ARCHEON = **A**ltered **R**egion of **C**hanged **C**hromosomal **E**xpression **O**bserved in **N**eoplasms). Chromosomal alterations often affect more than one gene. This is true for amplifications, duplications, insertions, integrations, inversions, translocations, and deletions. These changes can have influence on the expression level of single or multiple genes. Most commonly in the field of cancer diagnostics and treatment the changes of expression levels have been investigated for single, putative relevant target genes such as MLVI2 (5p14), NRASL3 (6p12), EGFR (7p12), c-myc (8q23), Cyclin D1 (11q13), IGF1R (15q25), HER-2/neu (17q12), PCNA (20q12). However, the altered expression level and interaction of multiple (i.e. more than two) genes within one genomic region with each other has not been addressed. Genes of an ARCHEON form gene clusters with tissue specific expression patterns. The mode of interaction of individual genes within such a gene cluster suspected to represent an ARCHEON can be either protein-protein or protein-nucleic acid interaction, which may be illustrated but not limited by the following examples: ARCHEON gene interaction may be in the same signal transduction pathway, may be receptor to ligand binding, receptor kinase and SH2 or SH3 binding, transcription factor to promoter binding, nuclear hormone receptor to transcription factor binding, phosphogroup donation (e.g. kinases) and acceptance (e.g. phosphoprotein), mRNA stabilizing protein binding and transcriptional processes. The individual activity and specificity of a pair genes and or the proteins encoded thereby or of a group of such in a higher order, may be readily deduced from literature, published or deposited within public databases by the skilled person. However in the context of an ARCHEON the interaction of members being part of an ARCHEON will potentiate, exaggerate or reduce their singular functions. This interaction is of importance in defined normal tissues in which they are normally co-expressed. Therefore, these clusters have been commonly conserved during evolution. The aberrant expression of members of these ARCHEON in neoplastic lesions, however, (especially within tissues in which they are normally not expressed) has influence on tumor characteristics such as growth, invasiveness and drug responsiveness. Due to the interaction of these neighboring genes it is of importance to determine the members of the ARCHEON which are involved in the deregulation events. In this regard amplification and deletion events in neoplastic lesions are of special interest.

[0057] The invention relates to a method for the detection of chromosomal alterations by (a) determining the relative mRNA abundance of individual mRNA species or (b) determining the copy number of one or more chromosomal region (s) by quantitative PCR. In one embodiment information on the genomic organization and spatial regulation of chromosomal regions is assessed by bioinformatic analysis of the sequence information of the human genome (UCSC, NCBI) and then combined with RNA expression data from GeneChip™ DNA-Arrays (Affymetrix) and/or quantitative PCR (TaqMan) from RNA-samples or genomic DNA.

[0058] In a further embodiment the functional relationship of genes located on a chromosomal region which is altered (amplified or deleted) is established. The altered chromosomal region is defined as an ARCHEON if genes located on that region functionally interact.

[0059] The 17q 12 locus was investigated as one model system, harboring the HER-2/neu gene. By establishing a high-resolution assay to detect amplification events in neighboring genes, 43 genes that are commonly co-amplified in breast cancer cell lines and patient samples were identified. By gene array technologies and immunological methods their co-overexpression in tumor samples was demonstrated. Surprisingly, by clustering tissue samples with HER-2/neu positive Tumor samples, it was found that the expression pattern of this larger genomic region (consisting of 43 genes) is very similar to control brain tissue. HER-2/neu negative breast tumor tissue did not show a similar expression pattern. Indeed, some of the genes within these cluster are important for neural development (HER-2/neu, THRA) in mouse model systems or are described to be expressed in neural cells (NeuroD2). Moreover, by searching similar

gene combinations in the human and rodent genome additional homologous chromosomal regions on chromosome 3p21 and 12q13 harboring several isoforms of the respective genes (see below) were found. There was a strong evidence for multiple interactions between the 43 candidate genes, as being part of identical pathways (HER-2, neu, GRB7, CrkRS, CDC6), influencing the expression of each other (HER-2/neu, THRA, RARA), interacting with each other (PPARGBP, THRA, RARA, NR1D1 or HER-2/neu, GRB7) or expressed in defined tissues (CACNB1, PPARGBP, etc.). Interestingly, the genomic regions of the ARCHEONs that were identified are amplified in acquired Tamoxifen resistance of HER-2/neu negative cells (MCF7), which are normally sensitive to Tamoxifen treatment [Achuthan et al., 2001,(2)].

[0060] Moreover, altered responsiveness to treatment due to the alterations of the genes within these ARCHEONs was observed. Surprisingly, genes within the ARCHEONs are of importance even in the absence of HER-2/neu homologues. Some of the genes within the ARCHEONs, do not only serve as marker genes for prognostic purposes, but have already been known as targets for therapeutic intervention. For example TOP2 alpha is a target of anthracyclins. THRA and RARA can be targeted by hormones and hormone analogs (e.g. T3, rT3, RA). Due to their high affinity binding sites and available screening assays (reporter assays based on their transcriptional potential) the hormone receptors which are shown to be linked to neoplastic pathophysiology for the first time herein are ideal targets for drug screening and treatment of malignant neoplasia and breast cancer in particular. In this regard it is essential to know which members of the ARCHEON are altered in the neoplastic lesions. Particularly it is important to know the nature, number and extent to which the ARCHEON genes are amplified or deleted. The ARCHEONs are flanked by similar, endogenous retroviruses (e.g. HERV-K= "human endogenous retrovirus"), some of which are activated in breast cancer. These viruses may have also been involved in the evolutionary duplication of the ARCHEONs.

[0061] The analysis of the 17q12 region proved data obtained by IHC and identified several additional genes being co-amplified with the HER-2/neu gene. Comparative Analysis of RNA-based quantitative RT-PCR (TaqMan) with DNA-based qPCR from tumor cell lines identified the same amplified region. Genes at the 17q11.2 -21. region are offered by way of illustration not by way of limitation. A graphical display of the described chromosomal region is provided in Figure 1.

Biological relevance of the genes which are part of the 17q12 ARCHEON

MLN50

[0062] By differential screening of cDNAs from breast cancer-derived metastatic axillary lymph nodes, TRAF4 and 3 other novel genes (MLN51, MLN62, MLN64) were identified that are overexpressed in breast cancer [Tomasetto et al., 1995, (3)]. One gene, which they designated MLN50, was mapped to 17q11-q21.3 by radioactive in situ hybridization. In breast cancer cell lines, overexpression of the 4 kb MLN50 mRNA was correlated with amplification of the gene and with amplification and overexpression of ERBB2, which maps to the same region. The authors suggested that the 2 genes belong to the same amplicon. Amplification of chromosomal region 17q11-q21 is one of the most common events occurring in human breast cancers. They reported that the predicted 261-amino acid MLN50 protein contains an N-terminal LIM domain and a C-terminal SH3 domain. They renamed the protein LASP1, for 'LIM and SH3 protein.' Northern blot analysis revealed that LASP1 mRNA was expressed at a basal level in all normal tissues examined and overexpressed in 8% of primary breast cancers. In most of these cancers, LASP1 and ERBB2 were simultaneously overexpressed.

MLLT6

[0063] The MLLT6 (AF17) gene encodes a protein of 1,093 amino acids, containing a leucine-zipper dimerization motif located 3-prime of the fusion point and a cysteine-rich domain at the end terminus. AF17 was found to contain stretches of amino acids previously associated with domains involved in transcriptional repression or activation.

[0064] Chromosome translocations involving band 11q23 are associated with approximately 10% of patients with acute lymphoblastic leukemia (ALL) and more than 5% of patients with acute myeloid leukemia (AML). The gene at 11q23 involved in the translocations is variously designated ALL1, HRX, MLL, and TAX1. The partner gene in one of the rarer translocations, t(11;17)(q23;q21), designated MLLT6 on 17q12.

ZNF144 (Mel18)

[0065] Mel18 cDNA encodes a novel cys-rich zinc finger motif. The gene is expressed strongly in most tumor cell lines, but its normal tissue expression was limited to cells of neural origin and was especially abundant in fetal neural cells. It belongs to a RING-finger motif family which includes BMI1. The MEL18/BMI1 gene family represents a mammalian homolog of the Drosophila 'polycomb' gene group, thereby belonging to a memory mechanism involved in

maintaining the the expression pattern of key regulatory factors such as Hox genes. Bmi1, Mel18 and M33 genes, as representative examples of mouse Pc-G genes. Common phenotypes observed in knockout mice mutant for each of these genes indicate an important role for Pc-G genes not only in regulation of Hox gene expression and axial skeleton development but also in control of proliferation and survival of haematopoietic cell lineages. This is in line with the observed proliferative deregulation observed in lymphoblastic leukemia. The MEL18 gene is conserved among vertebrates. Its mRNA is expressed at high levels in placenta, lung, and kidney, and at lower levels in liver, pancreas, and skeletal muscle. Interestingly, cervical and lumbo-sacral-HOX gene expression is altered in several primary breast cancers with respect to normal breast tissue with the HoxB gene cluster being present on 17q distal to the 17q12 locus. Moreover, delay of differentiation with persistent nests of proliferating cells was found in endothelial cells cocultured with HOXB7-transduced SkBr3 cells, which exhibit a 17q12 amplification. Tumorigenicity of these cells has been evaluated in vivo. Xenograft in athymic nude mice showed that SkBr3/HOXB7 cells developed tumors with an increased number of blood vessels, either irradiated or not, whereas parental SkBr3 cells did not show any tumor take unless mice were sublethally irradiated. As part of this invention, we have found MEL18 to be overexpressed specifically in tumors bearing Her-2/neu gene amplification, which can be critical for Hox expression.

PHOSPHATIDYLINOSITOL-4-PHOSPHATE 5-KINASE, TYPE II, BETA; PIP5K2B

[0066] Phosphoinositide kinases play central roles in signal transduction. Phosphatidylinositol-4-phosphate 5-kinases (PIP5Ks) phosphorylate phosphatidylinositol 4-phosphate, giving rise to phosphatidylinositol 4,5-bisphosphate. The PIP5K enzymes exist as multiple isoforms that have various immunoreactivities, kinetic properties, and molecular masses. They are unique in that they possess almost no homology to the kinase motifs present in other phosphatidylinositol, protein, and lipid kinases. By screening a human fetal brain cDNA library with the PIP5K2B EST the full length gene could be isolated. The deduced 416-amino acid protein is 78% identical to PIP5K2A. Using SDS-PAGE, the authors estimated that bacterially expressed PIP5K2B has a molecular mass of 47 kD. Northern blot analysis detected a 6.3-kb PIP5K2B transcript which was abundantly expressed in several human tissues. PIP5K2B interacts specifically with the juxtamembrane region of the p55 TNF receptor (TNFR1) and PIP5K2B activity is increased in mammalian cells by treatment with TNF-alpha. A modeled complex with membrane-bound substrate and ATP shows how a phosphoinositide kinase can phosphorylate its substrate in situ at the membrane interface. The substrate-binding site is open on 1 side, consistent with dual specificity for phosphatidylinositol 3- and 5-phosphates. Although the amino acid sequence of PIP5K2A does not show homology to known kinases, recombinant PIP5K2A exhibited kinase activity. PIP5K2A contains a putative Src homology 3 (SH3) domain-binding sequence. Overexpression of mouse PIP5K1B in COS7 cells induced an increase in short actin fibers and a decrease in actin stress fibers.

TEM7

[0067] Using serial analysis of gene expression (SAGE) a partial cDNAs corresponding to several tumor endothelial markers (TEMs) that displayed elevated expression during tumor angiogenesis could be identified. Among the genes identified was TEM7. Using database searches and 5-prime RACE the entire TEM7 coding region, which encodes a 500-amino acid type I transmembrane protein, has been described. The extracellular region of TEM7 contains a plexin-like domain and has weak homology to the ECM protein nidogen. The function of these domains, which are usually found in secreted and extracellular matrix molecules, is unknown. Nidogen itself belongs to the entactin protein family and helps to determine pathways of migrating axons by switching from circumferential to longitudinal migration. Entactin is involved in cell migration, as it promotes trophoblast outgrowth through a mechanism mediated by the RGD recognition site, and plays an important role during invasion of the endometrial basement membrane at implantation. As entactin promotes thymocyte adhesion but affects thymocyte migration only marginally, it is suggested that entactin may plays a role in thymocyte localization during T cell development.

[0068] In situ hybridization analysis of human colorectal cancer demonstrated that TEM7 was expressed clearly in the endothelial cells of the tumor stroma but not in the endothelial cells of normal colonic tissue. Using in situ hybridization to assay expression in various normal adult mouse tissues, they observed that TEM7 was largely undetectable in mouse tissues or tumors, but was abundantly expressed in mouse brain.

ZNFN1A3

[0069] By screening a B-cell cDNA library with a mouse Aiolos N-terminal cDNA probe, a cDNA encoding human Aiolos, or ZNFN1A3, was obtained. The deduced 509-amino acid protein, which is 86% identical to its mouse counterpart, has 4 DNA-binding zinc fingers in its N terminus and 2 zinc fingers that mediate protein dimerization in its C terminus. These domains are 100% and 96% homologous to the corresponding domains in the mouse protein, respectively. Northern blot analysis revealed strong expression of a major 11.0- and a minor 4.4-kb ZNFN1A3 transcript in

peripheral blood leukocytes, spleen, and thymus, with lower expression in liver, small intestine, and lung.

[0070] Ikaros (ZNFN1A1), a hemopoietic zinc finger DNA-binding protein, is a central regulator of lymphoid differentiation and is implicated in leukemogenesis. The execution of normal function of Ikaros requires sequence-specific DNA binding, transactivation, and dimerization domains. Mice with a mutation in a related zinc finger protein, Aiolos, are prone to B-cell lymphoma. In chemically induced murine lymphomas allelic losses on markers surrounding the Znf1al gene were detected in 27% of the tumors analyzed. Moreover specific Ikaros expression was in primary mouse hormone-producing anterior pituitary cells and substantial for Fibroblast growth factor receptor 4 (FGFR4) expression, which itself is implicated in a multitude of endocrine cell hormonal and proliferative properties with FGFR4 being differentially expressed in normal and neoplastic pituitary. Moreover Ikaros binds to chromatin remodelling complexes containing SWI/SNF proteins, which antagonize Polycomb function. Interestingly at the telomeric end of the disclosed ARCHEON the SWI/SNF complex member SMARCE1 (= SWI/SNF-related, matrix-associated, actin-dependent regulators of chromatin) is located and part of the described amplification. Due to the related binding specificities of Ikaros and Palindrom Binding Protein (PBP) it is suggestive, that ZNFN1A3 is able to regulate the Her-2/neu enhancer.

PPP1R1B

[0071] Midbrain dopaminergic neurons play a critical role in multiple brain functions, and abnormal signaling through dopaminergic pathways has been implicated in several major neurologic and psychiatric disorders. One well-studied target for the actions of dopamine is DARPP32. In the densely dopamine- and glutamate-innervated rat caudate-putamen, DARPP32 is expressed in medium-sized spiny neurons that also express dopamine D1 receptors. The function of DARPP32 seems to be regulated by receptor stimulation. Both dopaminergic and glutamatergic (NMDA) receptor stimulation regulate the extent of DARPP32 phosphorylation, but in opposite directions.

[0072] The human DARPP32 was isolated from a striatal cDNA library. The 204-amino acid DARPP32 protein shares 88% and 85% sequence identity, respectively, with bovine and rat DARPP32 proteins. The DARPP32 sequence is particularly conserved through the N terminus, which represents the active portion of the protein. Northern blot analysis demonstrated that the 2.1-kb DARPP32 mRNA is more highly expressed in human caudate than in cortex. In situ hybridization to postmortem human brain showed a low level of DARPP32 expression in all neocortical layers, with the strongest hybridization in the superficial layers. CDK5 phosphorylated DARPP32 in vitro and in intact brain cells. Phospho-thr75 DARPP32 inhibits PKA in vitro by a competitive mechanism. Decreasing phospho-thr75 DARPP32 in striatal cells either by a CDK5-specific inhibitor or by using genetically altered mice resulted in increased dopamine-induced phosphorylation of PKA substrates and augmented peak voltage-gated calcium currents. Thus, DARPP32 is a bifunctional signal transduction molecule which, by distinct mechanisms, controls a serine/threonine kinase and a serine/threonine phosphatase.

[0073] DARPP32 and t-DARPP are overexpressed in gastric cancers. It's suggested that overexpression of these 2 proteins in gastric cancers may provide an important survival advantage to neoplastic cells. It could be demonstrated that Darpp32 is an obligate intermediate in progesterone-facilitated sexual receptivity in female rats and mice. The facilitative effect of progesterone on sexual receptivity in female rats was blocked by antisense oligonucleotides to Darpp32. Homozygous mice carrying a null mutation for the Darpp32 gene exhibited minimal levels of progesterone-facilitated sexual receptivity when compared to their wildtype littermates, and progesterone significantly increased hypothalamic cAMP levels and cAMP-dependent protein kinase activity.

CACNB1

[0074] In 1991 a cDNA clone encoding a protein with high homology to the beta subunit of the rabbit skeletal muscle dihydropyridine-sensitive calcium channel from a rat brain cDNA library [Pragnell et al., 1991, (4)]. This rat brain beta-subunit cDNA hybridized to a 3.4-kb message that was expressed in high levels in the cerebral hemispheres and hippocampus and much lower levels in cerebellum. The open reading frame encodes 597 amino acids with a predicted mass of 65,679 Da which is 82% homologous with the skeletal muscle beta subunit. The corresponding human beta-subunit gene was localized to chromosome 17 by analysis of somatic cell hybrids. The authors suggested that the encoded brain beta subunit, which has a primary structure highly similar to its isoform in skeletal muscle, may have a comparable role as an integral regulatory component of a neuronal calcium channel.

RPL19

[0075] The ribosome is the only organelle conserved between prokaryotes and eukaryotes. In eukaryotes, this organelle consists of a 60S large subunit and a 40S small subunit. The mammalian ribosome contains 4 species of RNA and approximately 80 different ribosomal proteins, most of which appear to be present in equimolar amounts. In mammalian cells, ribosomal proteins can account for up to 15% of the total cellular protein, and the expression of the different

ribosomal protein genes, which can account for up to 7 to 9% of the total cellular mRNAs, is coordinately regulated to meet the cell's varying requirements for protein synthesis. The mammalian ribosomal protein genes are members of multigene families, most of which are composed of multiple processed pseudogenes and a single functional intron-containing gene. The presence of multiple pseudogenes hampered the isolation and study of the functional ribosomal protein genes. By study of somatic cell hybrids, it has been elucidated that DNA sequences complementary to 6 mam-
 5 mammalian ribosomal protein cDNAs could be assigned to chromosomes 5, 8, and 17. Ten fragments mapped to 3 chromosomes [Nakamichi et al., 1986, (5)]. These are probably a mixture of functional (expressed) genes and pseudogenes. One that maps to 5q23-q33 rescues Chinese hamster emetine-resistance mutations in interspecies hybrids and is therefore the transcriptionally active RPS14 gene. In 1989 a PCR-based strategy for the detection of intron-containing
 10 genes in the presence of multiple pseudogenes was described. This technique was used to identify the intron-containing PCR products of 7 human ribosomal protein genes and to map their chromosomal locations by hybridization to human/rodent somatic cell hybrids [Feo et al., 1992, (6)]. All 7 ribosomal protein genes were found to be on different chromosomes: RPL19 on 17p12-q11; RPL30 on 8; RPL35A on 18; RPL36A on 14; RPS6 on 9pter-p13; RPS11 on 19cen-qter; and RPS17 on 11pter-p13. These are also different sites from the chromosomal location of previously mapped ribosomal protein genes S14 on chromosome 5, S4 on Xq and Yp, and RP117A on 9q3-q34. By fluorescence in situ hybridization the position of the RPL19 gene was mapped to 17q11 [Davies et al., 1989, (7)].

PPARBP, PBP, CRSP1, CRSP200, TRIP2, TRAP220, RB18A, DRIP230

[0076] The thyroid hormone receptors (TRs) are hormone-dependent transcription factors that regulate expression of a variety of specific target genes. They must specifically interact with a number of proteins as they progress from their initial translation and nuclear translocation to heterodimerization with retinoid X receptors (RXRs), functional interactions with other transcription factors and the basic transcriptional apparatus, and eventually, degradation. To help
 20 elucidate the mechanisms that underlie the transcriptional effects and other potential functions of TRs, the yeast interaction trap, a version of the yeast 2-hybrid system, was used to identify proteins that specifically interact with the ligand-binding domain of rat TR-beta-1 (THRB) [Lee et al., 1995, (8)]. The authors isolated HeLa cell cDNAs encoding several different TR-interacting proteins (TRIPs), including TRIP2. TRIP2 interacted with rat Thrb only in the presence of thyroid hormone. It showed a ligand-independent interaction with RXR-alpha, but did not interact with the glucocorticoid receptor (NR3C1) under any condition. By immunoscreening a human B-lymphoma cell cDNA expression library with
 25 the anti-p53 monoclonal antibody PAb1801, PPARBP was identified, which was called RB18A for 'recognized by PAb1801 monoclonal antibody' [Drane et al., 1997, (9)]. The predicted 1,566-amino acid RB 18A protein contains several potential nuclear localization signals, 13 potential N-glycosylation sites, and a high number of potential phosphorylation sites. Despite sharing common antigenic determinants with p53, RB18A does not show significant nucleotide or amino acid sequence similarity with p53. Whereas the calculated molecular mass of RB18A is 166 kD, the
 30 apparent mass of recombinant RB18A was 205 kD by SDS-PAGE analysis. The authors demonstrated that RB18A shares functional properties with p53, including DNA binding, p53 binding, and self-oligomerization. Furthermore, RB18A was able to activate the sequence-specific binding of p53 to DNA, which was induced through an unstable interaction between both proteins. Northern blot analysis of human tissues detected an 8.5-kb RB18A transcript in all tissues examined except kidney, with highest expression in heart. Moreover mouse Pparbp, which was called Pbp for
 35 'Ppar-binding protein,' as a protein that interacts with the Ppar-gamma (PPARG) ligand-binding domain in a yeast 2-hybrid system was identified [Zhu et al., 1997, (10)]. The authors found that Pbp also binds to PPAR-alpha (PPARA), RAR-alpha (RARA), RXR, and TR-beta-1 in vitro. The binding of Pbp to these receptors increased in the presence of specific ligands. Deletion of the last 12 amino acids from the C terminus of PPAR-gamma resulted in the abolition of interaction between Pbp and PPAR-gamma. Pbp modestly increased the transcriptional activity of PPAR-gamma, and
 40 a truncated form of Pbp acted as a dominant-negative repressor, suggesting that Pbp is a genuine transcriptional co-activator for PPAR. The predicted 1,560-amino acid Pbp protein contains 2 LXXLL motifs, which are considered necessary and sufficient for the binding of several co-activators to nuclear receptors. Northern blot analysis detected Pbp expression in all mouse tissues examined, with higher levels in liver, kidney, lung, and testis. In situ hybridization showed that Pbp is expressed during mouse ontogeny, suggesting a possible role for Pbp in cellular proliferation and
 45 differentiation. In adult mouse, in situ hybridization detected Pbp expression in liver, bronchial epithelium in the lung, intestinal mucosa, kidney cortex, thymic cortex, splenic follicles, and seminiferous epithelium in testis. Lateron PPARBP was identified, which was called TRAP220, from an immunopurified TR-alpha (THRA)-TRAP complex [Yuan et al., 1998, (11)]. The authors cloned Jurkat cell cDNAs encoding TRAP220. The predicted 1,581-amino acid TRAP220 protein contains LXXLL domains, which are found in other nuclear receptor-interacting proteins. TRAP220 is nearly
 50 identical to RB18A, with these proteins differing primarily by an extended N terminus on TRAP220. In the absence of TR-alpha, TRAP220 appears to reside in a single complex with other TRAPs. TRAP220 showed a direct ligand-dependent interaction with TR-alpha, which was mediated through the C terminus of TR-alpha and, at least in part, the LXXLL domains of TRAP220. TRAP220 also interacted with other nuclear receptors, including vitamin D receptor,

RARA, RXRA, PPARA, PPARG, and estrogen receptor-alpha (ESR1; 133430), in a ligand-dependent manner. TRAP220 moderately stimulated human TR-alpha-mediated transcription in transfected cells, whereas a fragment containing the LXXLL motifs acted as a dominant-negative inhibitor of nuclear receptor-mediated transcription both in transfected cells and in cell-free transcription systems. Further studies indicated that TRAP220 plays a major role in anchoring other TRAPs to TR-alpha during the function of the TR-alpha-TRAP complex and that TRAP220 may be a global co-activator for the nuclear receptor superfamily. PBP, a nuclear receptor co-activator, interacts with estrogen receptor-alpha (ESR1) in the absence of estrogen. This interaction was enhanced in the presence of estrogen, but was reduced in the presence of the anti-estrogen Tamoxifen. Transfection of PBP into cultured cells resulted in enhancement of estrogen-dependent transcription, indicating that PBP serves as a co-activator in estrogen receptor signaling. To examine whether overexpression of PBP plays a role in breast cancer because of its co-activator function in estrogen receptor signaling, the levels of PBP expression in breast tumors was determined [Zhu et al., 1999, (12)]. High levels of PBP expression were detected in approximately 50% of primary breast cancers and breast cancer cell lines by ribonuclease protection analysis, in situ hybridization, and immunoperoxidase staining. By using FISH, the authors mapped the PBP gene to 17q12, a region that is amplified in some breast cancers. They found PBP gene amplification in approximately 24% (6 of 25) of breast tumors and approximately 30% (2 of 6) of breast cancer cell lines, implying that PBP gene overexpression can occur independent of gene amplification. They determined that the PBP gene comprises 17 exons that together span more than 37 kb. Their findings, in particular PBP gene amplification, suggested that PBP, by its ability to function as an estrogen receptor-alpha co-activator, may play a role in mammary epithelial differentiation and in breast carcinogenesis.

NEUROD2

[0077] Basic helix-loop-helix (bHLH) proteins are transcription factors involved in determining cell type during development. In 1995 a bHLH protein was described, termed NeuroD (for 'neurogenic differentiation'), that functions during neurogenesis [Lee et al., 1995, (13)]. The human NEUROD gene maps to chromosome 2q32. The cloning and characterization of 2 additional NEUROD genes, NEUROD2 and NEUROD3 was described in 1996 [McCormick et al., 1996, (14)]. Sequences for the mouse and human homologues were presented. NEUROD2 shows a high degree of homology to the bHLH region of NEUROD, whereas NEUROD3 is more distantly related. The authors found that mouse neuroD2 was initially expressed at embryonic day 11, with persistent expression in the adult nervous system. Similar to neuroD, neuroD2 appears to mediate neuronal differentiation. The human NEUROD2 was mapped to 17q12 by fluorescence in situ hybridization and the mouse homologue to chromosome 11 [Tamimi et al., 1997, (15)].

TELETHONIN

[0078] Telethonin is a sarcomeric protein of 19 kD found exclusively in striated and cardiac muscle. It appears to be localized to the Z disc of adult skeletal muscle and cultured myocytes. Telethonin is a substrate of titin, which acts as a molecular 'ruler' for the assembly of the sarcomere by providing spatially defined binding sites for other sarcomeric proteins. After activation by phosphorylation and calcium/calmodulin binding, titin phosphorylates the C-terminal domain of telethonin in early differentiating myocytes. The telethonin gene has been mapped to 17q12, adjacent to the phenylethanolamine N-methyltransferase gene [Valle et al., 1997, (16)].

PENT, PNMT

[0079] Phenylethanolamine N-methyltransferase catalyzes the synthesis of epinephrine from norepinephrine, the last step of catecholamine biosynthesis. The cDNA clone was first isolated in 1998 for bovine adrenal medulla PNMT using mixed oligodeoxyribonucleotide probes whose synthesis was based on the partial amino acid sequence of tryptic peptides from the bovine enzyme [Kaneda et al., 1988, (17)]. Using a bovine cDNA as a probe, the authors screened a human pheochromocytoma cDNA library and isolated a cDNA clone with an insert of about 1.0 kb, which contained a complete coding region of the enzyme. Northern blot analysis of human pheochromocytoma polyadenylated RNA using this cDNA insert as the probe demonstrated a single RNA species of about 1,000 nucleotides, suggesting that this clone is a full-length cDNA. The nucleotide sequence showed that human PNMT has 282 amino acid residues with a predicted molecular weight of 30,853, including the initial methionine. The amino acid sequence was 88% homologous to that of bovine enzyme. The PNMT gene was found to consist of 3 exons and 2 introns spanning about 2,100 basepairs. It was demonstrated that in transgenic mice the gene is expressed in adrenal medulla and retina. A hybrid gene consisting of 2 kb of the PNMT 5-prime-flanking region fused to the simian virus 40 early region also resulted in tumor antigen mRNA expression in adrenal glands and eyes; furthermore, immunocytochemistry showed that the tumor antigen was localized in nuclei of adrenal medullary cells and cells of the inner nuclear cell layer of the retina, both prominent sites of epinephrine synthesis. The results indicate that the enhancer(s) for appropriate expres-

sion of the gene in these cell types are in the 2-kb 5-prime-flanking region of the gene.

[0080] Kaneda et al., 1988 (17), assigned the human PNMT gene to chromosome 17 by Southern blot analysis of DNA from mouse-human somatic cell hybrids. In 1992 the localization was narrowed down to 17q21-q22 by linkage analysis using RFLPs related to the PNMT gene and several 17q DNA markers [Hoehe et al., 1992, (18)]. The findings are of interest in light of the description of a genetic locus associated with blood pressure regulation in the stroke-prone spontaneously hypertensive rat (SHR-SP) on rat chromosome 10 in a conserved linkage synteny group corresponding to human chromosome 17q22-q24. See essential hypertension .

MGC9753

[0081] This gene maps on chromosome 17, at 17q12 according to RefSeq. It is expressed at very high level. It is defined by cDNA clones and produces, by alternative splicing, 7 different transcripts can be obtained (SEQ ID NO:60 to 66 and 83 to 89, Table 1), altogether encoding 7 different protein isoforms. Of specific interest is the putatively secreted isoform g, encoded by a mRNA of 2.55 kb. It's premessenger covers 16.94 kb on the genome. It has a very long 3' UTR. . The protein (226 aa, MW 24.6 kDa, pI 8.5) contains no Pfam motif. The MGC9753 gene produces, by alternative splicing, 7 types of transcripts, predicted to encode 7 distinct proteins. It contains 13 confirmed introns, 10 of which are alternative. Comparison to the genome sequence shows that 11 introns follow the consensual [gt-ag] rule, 1 is atypical with good support [tg_cg]. The six most abundant isoforms are designated by a) to i) and code for proteins as follows:

a) This mRNA is 3.03 kb long, its premessenger covers 16.95 kb on the genome. It has a very long 3' UTR. The protein (190 aa, MW 21.5 kDa, pI 7.2) contains no Pfam motif. It is predicted to localise in the endoplasmic reticulum.

c) This mRNA is 1.17 kb long, its premessenger covers 16.93 kb on the genome. It may be incomplete at the N terminus. The protein (368 aa, MW 41.5 kDa, pI 7.3) contains no Pfam motif.

d) This mRNA is 3.17 kb long, its premessenger covers 16.94 kb on the genome. It has a very long 3' UTR and 5'p UTR. . The protein (190 aa, MW 21.5 kDa, pI 7.2) contains no Pfam motif. It is predicted to localise in the endoplasmic reticulum.

g) This mRNA is 2.55 kb long, its premessenger covers 16.94 kb on the genome. It has a very long 3' UTR. . The protein (226 aa, MW 24.6 kDa, pI 8.5) contains no Pfam motif. It is predicted to be secreted.

h) This mRNA is 2.68 kb long, its premessenger covers 16.94 kb on the genome. It has a very long 3' UTR. . The protein (320 aa, MW 36.5 kDa, pI 6.8) contains no Pfam motif. It is predicted to localise in the endoplasmic reticulum.

i) This mRNA is 2.34 kb long, its premessenger covers 16.94 kb on the genome. It may be incomplete at the N terminus. It has a very long 3' UTR. . The protein (217 aa, MW 24.4 kDa, pI 5.9) contains no Pfam motif.

[0082] The MCG9753 gene may be homologue to the CAB2 gene located on chromosome 17q12. The CAB2, a human homologue of the yeast COS16 required for the repair of DNA double-strand breaks was cloned. Autofluorescence analysis of cells transfected with its GFP fusion protein demonstrated that CAB2 translocates into vesicles, suggesting that overexpression of CAB2 may decrease intercellular Mn-(2+) by accumulating it in the vesicles, in the same way as yeast.

Her-2/neu, ERBB2, NGL, TKR1

[0083] The oncogene originally called NEU was derived from rat neuro/glioblastoma cell lines. It encodes a tumor antigen, p185, which is serologically related to EGFR, the epidermal growth factor receptor. EGFR maps to chromosome 7. In 1985 it was found, that the human homologue, which they designated NGL (to avoid confusion with neuraminidase, which is also symbolized NEU), maps to 17q12-q22 by in situ hybridization and to 17q21-qter in somatic cell hybrids [Yang-Feng et al., 1985, (19)]. Thus, the SRO is 17q21-q22. Moreover, in 1985 a potential cell surface receptor of the tyrosine kinase gene family was identified and characterized by cloning the gene [Coussens et al., 1985, (20)]. Its primary sequence is very similar to that of the human epidermal growth factor receptor. Because of the seemingly close relationship to the human EGF receptor, the authors called the gene HER2. By Southern blot analysis of somatic cell hybrid DNA and by in situ hybridization, the gene was assigned to 17q21-q22. This chromosomal location of the gene is coincident with the NEU oncogene, which suggests that the 2 genes may in fact be the same; indeed, sequencing indicates that they are identical. In 1988 a correlation between overexpression of NEU protein and the

large-cell, comedo growth type of ductal carcinoma was found [van de Vijver et al., 1988, (21)]. The authors found no correlation, however, with lymph-node status or tumor recurrence. The role of HER2/NEU in breast and ovarian cancer was described in 1989, which together account for one-third of all cancers in women and approximately one-quarter of cancer-related deaths in females [Slamon et al., 1989, (22)].

[0084] An ERBB-related gene that is distinct from the ERBB gene, called ERBB1 was found in 1985. ERBB2 was not amplified in vulva carcinoma cells with EGFR amplification and did not react with EGF receptor mRNA. About 30-fold amplification of ERBB2 was observed in a human adenocarcinoma of the salivary gland. By chromosome sorting combined with velocity sedimentation and Southern hybridization, the ERBB2 gene was assigned to chromosome 17 [Fukushige et al., 1986, (23)]. By hybridization to sorted chromosomes and to metaphase spreads with a genomic probe, they mapped the ERBB2 locus to 17q21. This is the chromosome 17 breakpoint in acute promyelocytic leukemia (APL). Furthermore, they observed amplification and elevated expression of the ERBB2 gene in a gastric cancer cell line. Antibodies against a synthetic peptide corresponding to 14 amino acid residues at the COOH-terminus of a protein deduced from the ERBB2 nucleotide sequence were raised in 1986. With these antibodies, the ERBB2 gene product from adenocarcinoma cells was precipitated and demonstrated to be a 185-kD glycoprotein with tyrosine kinase activity. A cDNA probe for ERBB2 and by in situ hybridization to APL cells with a 15;17 chromosome translocation located the gene to the proximal side of the breakpoint [Kaneko et al., 1987, (24)]. The authors suggested that both the gene and the breakpoint are located in band 17q21.1 and, further, that the ERBB2 gene is involved in the development of leukemia. In 1987 experiments indicated that NEU and HER2 are both the same as ERBB2 [Di Fiore et al., 1987, (25)]. The authors demonstrated that overexpression alone can convert the gene for a normal growth factor receptor, namely, ERBB2, into an oncogene. The ERBB2 to 17q11-q21 by in situ hybridization [Popescu et al., 1989, (26)]. By in situ hybridization to chromosomes derived from fibroblasts carrying a constitutional translocation between 15 and 17, they showed that the ERBB2 gene was relocated to the derivative chromosome 15; the gene can thus be localized to 17q12-q21.32. By family linkage studies using multiple DNA markers in the 17q12-q21 region the ERBB2 gene was placed on the genetic map of the region.

[0085] Interleukin-6 is a cytokine that was initially recognized as a regulator of immune and inflammatory responses, but also regulates the growth of many tumor cells, including prostate cancer. Overexpression of ERBB2 and ERBB3 has been implicated in the neoplastic transformation of prostate cancer. Treatment of a prostate cancer cell line with IL6 induced tyrosine phosphorylation of ERBB2 and ERBB3, but not ERBB1/EGFR. The ERBB2 forms a complex with the gp130 subunit of the IL6 receptor in an IL6-dependent manner. This association was important because the inhibition of ERBB2 activity resulted in abrogation of IL6-induced MAPK activation. Thus, ERBB2 is a critical component of IL6 signaling through the MAP kinase pathway [Qiu et al., 1998, (27)]. These findings showed how a cytokine receptor can diversify its signaling pathways by engaging with a growth factor receptor kinase.

[0086] Overexpression of ERBB2 confers Taxol resistance in breast cancers. Overexpression of ERBB2 inhibits Taxol-induced apoptosis [Yu et al., 1998, (28)]. Taxol activates CDC2 kinase in MDA-MB-435 breast cancer cells, leading to cell cycle arrest at the G2/M phase and, subsequently, apoptosis. A chemical inhibitor of CDC2 and a dominant-negative mutant of CDC2 blocked Taxol-induced apoptosis in these cells. Overexpression of ERBB2 in MDA-MB-435 cells by transfection transcriptionally upregulates CDKN1A which associates with CDC2, inhibits Taxol-mediated CDC2 activation, delays cell entrance to G2/M phase, and thereby inhibits Taxol-induced apoptosis. In CDKN1A anti-sense-transfected MDA-MB-435 cells or in p21^{-/-} MEF cells, ERBB2 was unable to inhibit Taxol-induced apoptosis. Therefore, CDKN1A participates in the regulation of a G2/M checkpoint that contributes to resistance to Taxol-induced apoptosis in ERBB2-overexpressing breast cancer cells.

[0087] A secreted protein of approximately 68 kD was described, designated herstatin, as the product of an alternative ERBB2 transcript that retains intron 8 [Doherty et al., 1999, (29)]. This alternative transcript specifies 340 residues identical to subdomains I and II from the extracellular domain of p185ERBB2, followed by a unique C-terminal sequence of 79 amino acids encoded by intron 8. The recombinant product of the alternative transcript specifically bound to ERBB2-transfected cells and was chemically crosslinked to p185ERBB2, whereas the intron-encoded sequence alone also bound with high affinity to transfected cells and associated with p185 solubilized from cell extracts. The herstatin mRNA was expressed in normal human fetal kidney and liver, but was at reduced levels relative to p185ERBB2 mRNA in carcinoma cells that contained an amplified ERBB2 gene. Herstatin appears to be an inhibitor of p185ERBB2, because it disrupts dimers, reduces tyrosine phosphorylation of p185, and inhibits the anchorage-independent growth of transformed cells that overexpress ERBB2. The HER2 gene is amplified and HER2 is overexpressed in 25 to 30% of breast cancers, increasing the aggressiveness of the tumor. Finally, it was found that a recombinant monoclonal antibody against HER2 increased the clinical benefit of first-line chemotherapy in metastatic breast cancer that overexpresses HER2 [Slamon et al., 2001, (30)].

GRB7

[0088] Growth factor receptor tyrosine kinases (GF-RTKs) are involved in activating the cell cycle. Several substrates

of GF-RTKs contain Src-homology 2 (SH2) and SH3 domains. SH2 domain-containing proteins are a diverse group of molecules important in tyrosine kinase signaling. Using the CORT (cloning of receptor targets) method to screen a high expression mouse library, the gene for murine Grb7, which encodes a protein of 535 amino acids, was isolated [Margolis et al., 1992, (31)]. GRB7 is homologous to ras-GAP (ras-GTPase-activating protein). It contains an SH2 domain and is highly expressed in liver and kidney. This gene defines the GRB7 family, whose members include the mouse gene Grb 10 and the human gene GRB14.

[0089] A putative GRB7 signal transduction molecule and a GRB7V novel splice variant from an invasive human esophageal carcinoma was isolated [Tanaka et al., 1998, (32)]. Although both GRB7 isoforms shared homology with the Mig-10 cell migration gene of *Caenorhabditis elegans*, the GRB7V isoform lacked 88 basepairs in the C terminus; the resultant frameshift led to substitution of an SH2 domain with a short hydrophobic sequence. The wildtype GRB7 protein, but not the GRB7V isoform, was rapidly tyrosyl phosphorylated in response to EGF stimulation in esophageal carcinoma cells. Analysis of human esophageal tumor tissues and regional lymph nodes with metastases revealed that GRB7V was expressed in 40% of GRB7-positive esophageal carcinomas. GRB7V expression was enhanced after metastatic spread to lymph nodes as compared to the original tumor tissues. Transfection of an antisense GRB7 RNA expression construct lowered endogenous GRB7 protein levels and suppressed the invasive phenotype exhibited by esophageal carcinoma cells. These findings suggested that GRB7 isoforms are involved in cell invasion and metastatic progression of human esophageal carcinomas. By sequence analysis, The GRB7 gene was mapped to chromosome 17q21-q22, near the topoisomerase-2 gene [Dong et al., 1997, (33)]. GRB-7 is amplified in concert with HER2 in several breast cancer cell lines and that GRB-7 is overexpressed in both cell lines and breast tumors. GRB-7, through its SH2 domain, binds tightly to HER2 such that a large fraction of the tyrosine phosphorylated HER2 in SKBR-3 cells is bound to GRB-7 [Stein et al., 1994, (34)].

GCSF, CSF3

[0090] Granulocyte colony-stimulating factor (or colony stimulating factor-3) specifically stimulates the proliferation and differentiation of the progenitor cells for granulocytes. The partial amino acid sequence of purified GCSF protein was determined, and by using oligonucleotides as probes, several GCSF cDNA clones were isolated from a human squamous carcinoma cell line cDNA library [Nagata et al., 1986, (35)]. Cloning of human GCSF cDNA shows that a single gene codes for a 177- or 180-amino acid mature protein of molecular weight 19,600. The authors found that the GCSF gene has 4 introns and that 2 different polypeptides are synthesized from the same gene by differential splicing of mRNA. The 2 polypeptides differ by the presence or absence of 3 amino acids. Expression studies indicate that both have authentic GCSF activity. A stimulatory activity from a glioblastoma multiform cell line being biologically and biochemically indistinguishable from GCSF produced by a bladder cell line was found in 1987. By somatic cell hybridization and in situ chromosomal hybridization, the GCSF gene was mapped to 17q11 in the region of the breakpoint in the 15;17 translocation characteristic of acute promyelocytic leukemia [Le Beau et al., 1987, (36)]. Further studies indicated that the gene is proximal to the said breakpoint and that it remains on the rearranged chromosome 17. Southern blot analysis using both conventional and pulsed field gel electrophoresis showed no rearranged restriction fragments. By use of a full-length cDNA clone as a hybridization probe in human-mouse somatic cell hybrids and in flow-sorted human chromosomes, the gene for GCSF was mapped to 17q21-q22 lateron

THRA, THRA1, ERBA, EAR7, ERBA2, ERBA3

[0091] Both human and mouse DNA have been demonstrated to have two distantly related classes of ERBA genes and that in the human genome multiple copies of one of the classes exist [Jansson et al., 1983, (37)]. A cDNA was isolated derived from rat brain messenger RNA on the basis of homology to the human thyroid receptor gene [Thompson et al., 1987, (38)]. Expression of this cDNA produced a high-affinity binding protein for thyroid hormones. Messenger RNA from this gene was expressed in tissue-specific fashion, with highest levels in the central nervous system and no expression in the liver. An increasing body of evidence indicated the presence of multiple thyroid hormone receptors. The authors suggested that there may be as many as 5 different but related loci. Many of the clinical and physiologic studies suggested the existence of multiple receptors. For example, patients had been identified with familial thyroid hormone resistance in which peripheral response to thyroid hormones is lost or diminished while neuronal functions are maintained. Thyroidologists recognize a form of cretinism in which the nervous system is severely affected and another form in which the peripheral functions of thyroid hormone are more dramatically affected.

[0092] The cDNA encoding a specific form of thyroid hormone receptor expressed in human liver, kidney, placenta, and brain was isolated [Nakai et al., 1988, (39)]. Identical clones were found in human placenta. The cDNA encodes a protein of 490 amino acids and molecular mass of 54,824. Designated thyroid hormone receptor type alpha-2 (THRA2), this protein is represented by mRNAs of different size in liver and kidney, which may represent tissue-specific processing of the primary transcript.

[0093] The THRA gene contains 10 exons spanning 27 kb of DNA. The last 2 exons of the gene are alternatively spliced. A 5-kb THRA1 mRNA encodes a predicted 410-amino acid protein; a 2.7-kb THRA2 mRNA encodes a 490-amino acid protein. A third isoform, TR-alpha-3, is derived by alternative splicing. The proximal 39 amino acids of the TH-alpha-2 specific sequences are deleted in TR-alpha-3. A second gene, THRB on chromosome 3, encodes 2 isoforms of TR-beta by alternative splicing. In 1989 the structure and function of the EAR1 and EAR7 genes was elucidated, both located on 17q21 [Miyajima et al., 1989, (40)]. The authors determined that one of the exons in the EAR7 coding sequence overlaps an exon of EAR1, and that the 2 genes are transcribed from opposite DNA strands. In addition, the EAR7 mRNA generates 2 alternatively spliced isoforms, referred to as EAR71 and EAR72, of which the EAR71 protein is the human counterpart of the chicken c-erbA protein.

[0094] The thyroid hormone receptors, beta, alpha-1, and alpha-2 3 mRNAs are expressed in all tissues examined and the relative amounts of the three mRNAs were roughly parallel. None of the 3 mRNAs was abundant in liver, which is the major thyroid hormone-responsive organ. This led to the assumption that another thyroid hormone receptor may be present in liver. It was found that ERBA, which potentiates ERBB, has an amino acid sequence different from that of other known oncogene products and related to those of the carbonic anhydrases [Debuire et al., 1984, (41)]. ERBA potentiates ERBB by blocking differentiation of erythroblasts at an immature stage. Carbonic anhydrases participate in the transport of carbon dioxide in erythrocytes. In 1986 it was shown that the ERBA protein is a high-affinity receptor for thyroid hormone. The cDNA sequence indicates a relationship to steroid-hormone receptors, and binding studies indicate that it is a receptor for thyroid hormones. It is located in the nucleus, where it binds to DNA and activates transcription.

[0095] Maternal thyroid hormone is transferred to the fetus early in pregnancy and is postulated to regulate brain development. The ontogeny of TR isoforms and related splice variants in 9 first-trimester fetal brains by semi-quantitative RT-PCR analysis has been investigated. Expression of the TR-beta-1, TR-alpha-1, and TR-alpha-2 isoforms was detected from 8.1 weeks' gestation. An additional truncated species was detected with the TR-alpha-2 primer set, consistent with the TR-alpha-3 splice variant described in the rat. All TR-alpha-derived transcripts were coordinately expressed and increased approximately 8-fold between 8.1 and 13.9 weeks' gestation. A more complex ontogenic pattern was observed for TR-beta-1, suggestive of a nadir between 8.4 and 12.0 weeks' gestation. The authors concluded that these findings point to an important role for the TR-alpha-1 isoform in mediating maternal thyroid hormone action during first-trimester fetal brain development.

[0096] The identification of the several types of thyroid hormone receptor may explain the normal variation in thyroid hormone responsiveness of various organs and the selective tissue abnormalities found in the thyroid hormone resistance syndromes. Members of sibships, who were resistant to thyroid hormone action, had retarded growth, congenital deafness, and abnormal bones, but had normal intellect and sexual maturation, as well as augmented cardiovascular activity. In this family abnormal T3 nuclear receptors in blood cells and fibroblasts have been demonstrated. The availability of cDNAs encoding the various thyroid hormone receptors was considered useful in determining the underlying genetic defect in this family.

[0097] The ERBA oncogene has been assigned to chromosome 17. The ERBA locus remains on chromosome 17 in the t(15;17) translocation of acute promyelocytic leukemia (APL). The thymidine kinase locus is probably translocated to chromosome 15; study of leukemia with t(17;21) and apparently identical breakpoint showed that TK was on 21q+. By in situ hybridization of a cloned DNA probe of c-erb-A to meiotic pachytene spreads obtained from uncultured spermatocytes it has been concluded that ERBA is situated at 17q21.33-17q22, in the same region as the break that generated the t(15;17) seen in APL. Because most of the grains were seen in 17q22, they suggested that ERBA is probably in the proximal region of 17q22 or at the junction between 17q22 and 17q21.33. By in situ hybridization it has been demonstrated, that that ERBA remains at 17q11-q12 in APL, whereas TP53, at 17q21-q22, is translocated to chromosome 15. Thus, ERBA must be at 17q11.2 just proximal to the breakpoint in the APL translocation and just distal to it in the constitutional translocation.

[0098] The aberrant THRA expression in nonfunctioning pituitary tumors has been hypothesized to reflect mutations in the receptor coding and regulatory sequences. They screened THRA mRNA and THRB response elements and ligand-binding domains for sequence anomalies. Screening THRA mRNA from 23 tumors by RNase mismatch and sequencing candidate fragments identified 1 silent and 3 missense mutations, 2 in the common THRA region and 1 that was specific for the alpha-2 isoform. No THRB response element differences were detected in 14 nonfunctioning tumors, and no THRB ligand-binding domain differences were detected in 23 nonfunctioning tumors. Therefore it has been suggested that the novel thyroid receptor mutations may be of functional significance in terms of thyroid receptor action, and further definition of their functional properties may provide insight into the role of thyroid receptors in growth control in pituitary cells.

RAR-alpha

[0099] A cDNA encoding a protein that binds retinoic acid with high affinity has been cloned [Petkovich et al., 1987,

(42)]. The protein was found to be homologous to the receptors for steroid hormones, thyroid hormones, and vitamin D3, and appeared to be a retinoic acid-inducible transacting enhancer factor. Thus, the molecular mechanisms of the effect of vitamin A on embryonic development, differentiation and tumor cell growth may be similar to those described for other members of this nuclear receptor family. In general, the DNA-binding domain is most highly conserved, both within and between the 2 groups of receptors (steroid and thyroid); Using a cDNA probe, the RAR-alpha gene has been mapped to 17q21 by in situ hybridization [Mattei et al., 1988, (43)]. Evidence has been presented for the existence of 2 retinoic acid receptors, RAR-alpha and RAR-beta, mapping to chromosome 17q21.1 and 3p24, respectively. The alpha and beta forms of RAR were found to be more homologous to the 2 closely related thyroid hormone receptors alpha and beta, located on 17q11.2 and 3p25-p21, respectively, than to any other members of the nuclear receptor family. These observations suggest that the thyroid hormone and retinoic acid receptors evolved by gene, and possibly chromosome, duplications from a common ancestor, which itself diverged rather early in evolution from the common ancestor of the steroid receptor group of the family. They noted that the counterparts of the human RARA and RARB genes are present in both the mouse and chicken. The involvement of RARA at the APL breakpoint may explain why the use of retinoic acid as a therapeutic differentiation agent in the treatment of acute myeloid leukemias is limited to APL. Almost all patients with APL have a chromosomal translocation t(15;17)(q22;q21). Molecular studies reveal that the translocation results in a chimeric gene through fusion between the PML gene on chromosome 15 and the RARA gene on chromosome 17. A hormone-dependent interaction of the nuclear receptors RARA and RXRA with CLOCK and MOP4 has been presented.

CDC18 L, CDC 6

[0100] In yeasts, Cdc6 (*Saccharomyces cerevisiae*) and Cdc18 (*Schizosaccharomyces pombe*) associate with the origin recognition complex (ORC) proteins to render cells competent for DNA replication. Thus, Cdc6 has a critical regulatory role in the initiation of DNA replication in yeast. cDNAs encoding *Xenopus* and human homologues of yeast CDC6 have been isolated [Williams et al., 1997, (44)]. They designated the human and *Xenopus* proteins p62(cdc6). Independently, in a yeast 2-hybrid assay using PCNA as bait, cDNAs encoding the human CDC6/Cdc 18 homologue have been isolated [Saha et al, 1998, (45)]. These authors reported that the predicted 560-amino acid human protein shares approximately 33% sequence identity with the 2 yeast proteins. On Western blots of HeLa cell extracts, human CDC6/Cdc18 migrates as a 66-kD protein. Although Northern blots indicated that CDC6/Cdc18 mRNA levels peak at the onset of S phase and diminish at the onset of mitosis in HeLa cells, the authors found that total CDC6/Cdc18 protein level is unchanged throughout the cell cycle. Immunofluorescent analysis of epitope-tagged protein revealed that human CDC6/Cdc18 is nuclear in G1 and cytoplasmic in S-phase cells, suggesting that DNA replication may be regulated by either the translocation of this protein between the nucleus and cytoplasm or by selective degradation of the protein in the nucleus. Immunoprecipitation studies showed that human CDC6/Cdc18 associates in vivo with cyclin A, CDK2, and ORC1. The association of cyclin-CDK2 with CDC6/Cdc18 was specifically inhibited by a factor present in mitotic cell extracts. Therefore it has been suggested that if the interaction between CDC6/Cdc18 with the S phase-promoting factor cyclin-CDK2 is essential for the initiation of DNA replication, the mitotic inhibitor of this interaction could prevent a premature interaction until the appropriate time in G1 Cdc6 is expressed selectively in proliferating but not quiescent mammalian cells, both in culture and within tissues in intact animals [Yan et al., 1998, (46)]. During the transition from a growth-arrested to a proliferative state, transcription of mammalian Cdc6 is regulated by E2F proteins, as revealed by a functional analysis of the human Cdc6 promoter and by the ability of exogenously expressed E2F proteins to stimulate the endogenous Cdc6 gene. Immunodepletion of Cdc6 by microinjection of anti-Cdc6 antibody blocked initiation of DNA replication in a human tumor cell line. The authors concluded that expression of human Cdc6 is regulated in response to mitogenic signals through transcriptional control mechanisms involving E2F proteins, and that Cdc6 is required for initiation of DNA replication in mammalian cells.

[0101] Using a yeast 2-hybrid system, co-purification of recombinant proteins, and immunoprecipitation, it has been demonstrated lateron that an N-terminal segment of CDC6 binds specifically to PR48, a regulatory subunit of protein phosphatase 2A (PP2A). The authors hypothesized that dephosphorylation of CDC6 by PP2A, mediated by a specific interaction with PR48 or a related B-double prime protein, is a regulatory event controlling initiation of DNA replication in mammalian cells. By analysis of somatic cell hybrids and by fluorescence in situ hybridization the human p62(cdc6) gene has been to 17q21.3.

TOP2A, TOP2

[0102] DNA topoisomerases are enzymes that control and alter the topologic states of DNA in both prokaryotes and eukaryotes. Topoisomerase II from eukaryotic cells catalyzes the relaxation of supercoiled DNA molecules, catenation, decatenation, knotting, and unknotting of circular DNA. It appears likely that the reaction catalyzed by topoisomerase II involves the crossing-over of 2 DNA segments. It has been estimated that there are about 100,000 molecules of

topoisomerase II per HeLa cell nucleus, constituting about 0.1% of the nuclear extract. Since several of the abnormal characteristics of ataxia-telangiectasia appear to be due to defects in DNA processing, screening for these enzyme activities in 5 AT cell lines has been performed [Singh et al., 1988, (47)]. In comparison to controls, the level of DNA topoisomerase II, determined by unknotting of P4 phage DNA, was reduced substantially in 4 of these cell lines and to a lesser extent in the fifth. DNA topoisomerase I, assayed by relaxation of supercoil DNA, was found to be present at normal levels.

[0103] The entire coding sequence of the human TOP2 gene has been determined [Tsai-Pflugfelder et al., 1988, (48)].

[0104] In addition human cDNAs that had been isolated by screening a cDNA library derived from a mechlorethamine-resistant Burkitt lymphoma cell line (Raji-HN2) with a Drosophila Topo II cDNA had been sequenced [Chung et al., 1989, (49)]. The authors identified 2 classes of sequence representing 2 TOP2 isoenzymes, which have been named TOP2A and TOP2B. The sequence of 1 of the TOP2A cDNAs is identical to that of an internal fragment of the TOP2 cDNA isolated by Tsai-Pflugfelder et al., 1988 (48). Southern blot analysis indicated that the TOP2A and TOP2B cDNAs are derived from distinct genes. Northern blot analysis using a TOP2A-specific probe detected a 6.5-kb transcript in the human cell line U937. Antibodies against a TOP2A peptide recognized a 170-kD protein in U937 cell lysates. Therefore it was concluded that their data provide genetic and immunochemical evidence for 2 TOP2 isozymes. The complete structures of the TOP2A and TOP2B genes has been reported [Lang et al., 1998, (50)]. The TOP2A gene spans approximately 30 kb and contains 35 exons.

[0105] Tsai-Pflugfelder et al., 1988 (48) showed that the human enzyme is encoded by a single-copy gene which they mapped to 17q21-q22 by a combination of in situ hybridization of a cloned fragment to metaphase chromosomes and by Southern hybridization analysis with a panel of mouse-human hybrid cell lines. The assignment to chromosome 17 has been confirmed by the study of somatic cell hybrids. Because of co-amplification in an adenocarcinoma cell line, it was concluded that the TOP2A and ERBB2 genes may be closely linked on chromosome 17 [Keith et al., 1992, (51)]. Using probes that detected RFLPs at both the TOP2A and TOP2B loci, the demonstrated heterozygosity at a frequency of 0.17 and 0.37 for the alpha and beta loci, respectively. The mouse homologue was mapped to chromosome 11 [Kingsmore et al., 1993, (52)]. The structure and function of type II DNA topoisomerases has been reviewed [Watt et al., 1994, (53)]. DNA topoisomerase II-alpha is associated with the pol II holoenzyme and is a required component of chromatin-dependent co-activation. Specific inhibitors of topoisomerase II blocked transcription on chromatin templates, but did not affect transcription on naked templates. Addition of purified topoisomerase II-alpha reconstituted chromatin-dependent activation activity in reactions with core pol II. Therefore the transcription on chromatin templates seems to result in the accumulation of superhelical tension, making the relaxation activity of topoisomerase II essential for productive RNA synthesis on nucleosomal DNA.

IGFBP4

[0106] Six structurally distinct insulin-like growth factor binding proteins have been isolated and their cDNAs cloned: IGFBP1, IGFBP2, IGFBP3, IGFBP4, IGFBP5 and IGFBP6. The proteins display strong sequence homologies, suggesting that they are encoded by a closely related family of genes. The IGFBPs contain 3 structurally distinct domains each comprising approximately one-third of the molecule. The N-terminal domain 1 and the C-terminal domain 3 of the 6 human IGFBPs show moderate to high levels of sequence identity including 12 and 6 invariant cysteine residues in domains 1 and 3, respectively (IGFBP6 contains 10 cysteine residues in domain 1), and are thought to be the IGF binding domains. Domain 2 is defined primarily by a lack of sequence identity among the 6 IGFBPs and by a lack of cysteine residues, though it does contain 2 cysteines in IGFBP4. Domain 3 is homologous to the thyroglobulin type I repeat unit. Recombinant human insulin-like growth factor binding proteins 4, 5, and 6 have been characterized by their expression in yeast as fusion proteins with ubiquitin [Kiefer et al., 1992, (54)]. Results of the study suggested to the authors that the primary effect of the 3 proteins is the attenuation of IGF activity and suggested that they contribute to the control of IGF-mediated cell growth and metabolism.

[0107] Based on peptide sequences of a purified insulin-like growth factor-binding protein (IGFBP) rat IGFBP4 has been cloned by using PCR [Shimasaki et al., 1990, (55)]. They used the rat cDNA to clone the human ortholog from a liver cDNA library. Human IGFBP4 encodes a 258-amino acid polypeptide, which includes a 21-amino acid signal sequence. The protein is very hydrophilic, which may facilitate its ability as a carrier protein for the IGFs in blood. Northern blot analysis of rat tissues revealed expression in all tissues examined, with highest expression in liver. It was stated that IGFBP4 acts as an inhibitor of IGF-induced bone cell proliferation. The genomic region containing the IGFBP gene. The gene consists of 4 exons spanning approximately 15 kb of genomic DNA has been examined [Zazzi et al., 1998, (56)]. The upstream region of the gene contains a TATA box and a cAMP-responsive promoter.

[0108] By in situ hybridization, the IGFBP4 gene was mapped to 17q12-q21 [Bajalica et al., 1992, (57)]. Because the hereditary breast-ovarian cancer gene BRCA1 had been mapped to the same region, it has been investigated whether IGFBP4 is a candidate gene by linkage analysis of 22 BRCA1 families; the finding of genetic recombination suggested that it is not the BRCA1 gene [Tonin et al., 1993, (58)].

EBI 1, CCR7, CMKBR7

[0109] Using PCR with degenerate oligonucleotides, a lymphoid-specific member of the G protein-coupled receptor family has been identified and mapped mapped to 17q12-q21.2 by analysis of human/mouse somatic cell hybrid DNAs and fluorescence in situ hybridization. It has been shown that this receptor had been independently identified as the Epstein-Barr-induced cDNA (symbol EBI1) [Birkenbach et al., 1993, (59)]. EBI1 is expressed in normal lymphoid tissues and in several B- and T-lymphocyte cell lines. While the function and the ligand for EBI1 remains unknown, its sequence and gene structure suggest that it is related to receptors that recognize chemoattractants, such as interleukin-8, RANTES, C5a, and fMet-Leu-Phe. Like the chemoattractant receptors, EBI1 contains intervening sequences near its 5-prime end; however, EBI1 is unique in that both of its introns interrupt the coding region of the first extracellular domain. Mouse Ebi1 cDNA has been isolated and found to encode a protein with 86% identity to the human homologue.

[0110] Subsets of murine CD4⁺ T cells localize to different areas of the spleen after adoptive transfer. Naive and T helper-1 (TH1) cells, which express CCR7, home to the periarteriolar lymphoid sheath, whereas activated TH2 cells, which lack CCR7, form rings at the periphery of the T-cell zones near B-cell follicles. It has been found that retroviral transduction of TH2 cells with CCR7 forced them to localize in a TH1-like pattern and inhibited their participation in B-cell help in vivo but not in vitro. Apparently differential expression of chemokine receptors results in unique cellular migration patterns that are important for effective immune responses.

[0111] CCR7 expression divides human memory T cells into 2 functionally distinct subsets. CCR7-memory cells express receptors for migration to inflamed tissues and display immediate effector function. In contrast, CCR7⁺ memory cells express lymph node homing receptors and lack immediate effector function, but efficiently stimulate dendritic cells and differentiate into CCR7⁻ effector cells upon secondary stimulation. The CCR7⁺ and CCR7⁻ T cells, named central memory (T-CM) and effector memory (T-EM), differentiate in a step-wise fashion from naive T cells, persist for years after immunization, and allow a division of labor in the memory response.

[0112] CCR7 expression in memory CD8⁺ T lymphocyte responses to HIV and to cytomegalovirus (CMV) tetramers has been evaluated. Most memory T lymphocytes express CD45RO, but a fraction express instead the CD45RA marker. Flow cytometric analyses of marker expression and cell division identified 4 subsets of HIV- and CMV-specific CD8⁺ T cells, representing a lineage differentiation pattern: CD45RA⁺CCR7⁺ (double-positive); CD45RA⁺CCR7⁻; CD45RA⁻CCR7⁻ (double-negative); CD45RA⁻CCR7⁺. The capacity for cell division, as measured by 5-(and 6)-carboxyl-fluorescein diacetate, succinimidyl ester, and intracellular staining for the Ki67 nuclear antigen, is largely confined to the CCR7⁺ subsets and occurred more rapidly in cells that are also CD45RA⁺. Although the double-negative cells did not divide or expand after stimulation, they did revert to positivity for either CD45RA or CCR7 or both. The CD45RA⁺CCR7⁻ cells, considered to be terminally differentiated, fail to divide, but do produce interferon-gamma and express high levels of perforin. The representation of subsets specific for CMV and for HIV is distinct. Approximately 70% of HIV-specific CD8⁺ memory T cells are double-negative or preterminally differentiated compared to 40% of CMV-specific cells. Approximately 50% of the CMV-specific CD8⁺ memory T cells are terminally differentiated compared to fewer than 10% of the HIV-specific cells. It has been proposed that terminally differentiated CMV-specific cells are poised to rapidly intervene, while double-positive precursor cells remain for expansion and replenishment of the effector cell pool. Furthermore, high-dose antigen tolerance and the depletion of HIV-specific CD4⁺ helper T-cell activity may keep the HIV-specific memory CD8⁺ T cells at the double-negative stage, unable to differentiate to the terminal effector state. B lymphocytes recirculate between B cell-rich compartments (follicles or B zones) in secondary lymphoid organs, surveying for antigen. After antigen binding, B cells move to the boundary of B and T zones to interact with T-helper cells. Furthermore it has been demonstrated that antigen-engaged B cells have increased expression of CCR7, the receptor for the T-zone chemokines CCL19 (also known as ELC) and CCL21, and that they exhibit increased responsiveness to both chemoattractants. In mice lacking lymphoid CCL19 and CCL21 chemokines, or with B cells that lack CCR7, antigen engagement fails to cause movement to the T zone. Using retroviral-mediated gene transfer, the authors demonstrated that increased expression of CCR7 is sufficient to direct B cells to the T zone. Reciprocally, overexpression of CXCR5, the receptor for the B-zone chemokine CXCL13, is sufficient to overcome antigen-induced B-cell movement to the T zone. This points toward a mechanism of B-cell relocalization in response to antigen, and established that cell position in vivo can be determined by the balance of responsiveness to chemoattractants made in separate but adjacent zones.

BAF57, SMARCE 1

[0113] The SWI/SNF complex in *S. cerevisiae* and *Drosophila* is thought to facilitate transcriptional activation of specific genes by antagonizing chromatin-mediated transcriptional repression. The complex contains an ATP-dependent nucleosome disruption activity that can lead to enhanced binding of transcription factors. The BRG1/brm-associated factors, or BAF, complex in mammals is functionally related to SWI/SNF and consists of 9 to 12 subunits, some of which are homologous to SWI/SNF subunits. A 57-kD BAF subunit, BAF57, is present in higher eukaryotes, but not in

yeast. Partial coding sequence has been obtained from purified BAF57 from extracts of a human cell line [Wang et al., 1998, (60)]. Based on the peptide sequences, they identified cDNAs encoding BAF57. The predicted 411-amino acid protein contains an HMG domain adjacent to a kinesin-like region. Both recombinant BAF57 and the whole BAF complex bind 4-way junction (4WJ) DNA, which is thought to mimic the topology of DNA as it enters or exits the nucleosome. The BAF57 DNA-binding activity has characteristics similar to those of other HMG proteins. It was found that complexes with mutations in the BAF57 HMG domain retain their DNA-binding and nucleosome-disruption activities. They suggested that the mechanism by which mammalian SWI/SNF-like complexes interact with chromatin may involve recognition of higher-order chromatin structure by 2 or more DNA-binding domains. RNase protection studies and Western blot analysis revealed that BAF57 is expressed ubiquitously. Several lines of evidence point toward the involvement of SWI/SNF factors in cancer development [Klochendler-Yeivin et al., 2002, (61)]. Moreover, SWI/SNF related genes are assigned to chromosomal regions that are frequently involved in somatic rearrangements in human cancers [Ring et al., 1998, (62)]. In this respect it is interesting that some of the SWI/SNF family members (i.e. SMARCC1, SMARCC2, SMARCD1 and SMARCD22 are neighboring 3 of the eucaryotic ARCHEONs we have identified (i.e. 3p21-p24, 12q13-q14 and 17q respectively) and which are part of the present invention. In this invention we could also map SMARCE1/BAF57 to the 17q12 region by PCR karyotyping.

KRT 10, K10

[0114] Keratin 10 is an intermediate filament (IF) chain which belongs to the acidic type I family and is expressed in terminally differentiated epidermal cells. Epithelial cells almost always co-express pairs of type I and type II keratins, and the pairs that are co-expressed are highly characteristic of a given epithelial tissue. For example, in human epidermis, 3 different pairs of keratins are expressed: keratins 5 (type II) and 14 (type I), characteristic of basal or proliferative cells; keratins 1 (type II) and 10 (type I), characteristic of suprabasal terminally differentiating cells; and keratins 6 (type II) and 16 (type I) (and keratin 17 [type II]), characteristic of cells induced to hyper-proliferate by disease or injury, and epithelial cells grown in cell culture. The nucleotide sequence of a 1,700 bp cDNA encoding human epidermal keratin 10 (56.5 kD) [Darmon et al., 1987, (63)] has been published as well as the complete amino acid sequence of human keratin 10 [Zhou et al., 1988, (64)]. Polymorphism of the KRT10 gene, restricted to insertions and deletions of the glycine-richquasipeptide repeats that form the glycine-loop motif in the C-terminal domain, have been extensively described [Korge et al., 1992, (65)].

[0115] By use of specific cDNA clones in conjunction with somatic cell hybrid analysis and in situ hybridization, KRT10 gene has been mapped to 17q12-q21 in a region proximal to the breakpoint at 17q21 that is involved in a t(17;21)(q21;q22) translocation associated with a form of acute leukemia. KRT10 appeared to be telomeric to 3 other loci that map in the same region: CSF3, ERBA1, and HER2 [Lessin et al., 1988, (66)]. NGFR and HOX2 are distal to K9. It has been demonstrated that the KRT10, KRT13, and KRT15 genes are located in the same large pulsed field gel electrophoresis fragment [Romano et al., 1991, (67)]. A correlation of assignments of the 3 genes makes 17q21-q22 the likely location of the cluster. Transgenic mice expressing a mutant keratin 10 gene have the phenotype of epidermolytic hyperkeratosis, thus suggesting that a genetic basis for the human disorder resides in mutations in genes encoding suprabasal keratins KRT1 or KRT10 [Fuchs et al 1992, (68)]. The authors also showed that stimulation of basal cell proliferation can result from a defect in suprabasal cells and that distortion of nuclear shape or alterations in cytokinesis can occur when an intermediate filament network is perturbed. In a family with keratosis palmaris et plantaris without blistering either spontaneously or in response to mild mechanical or thermal stress and with no involvement of the skin and parts of the body other than the palms and soles, a tight linkage to an insertion-deletion polymorphism in the C-terminal coding region of the KRT10 gene (maximum lod score = 8.36 at theta = 0.00) was found [Rogaev et al., 1993, (69)]. It is noteworthy that it was a rare, high molecular weight allele of the KRT10 polymorphism that segregated with the disorder. The allele was observed once in 96 independent chromosomes from unaffected Caucasians. The KRT10 polymorphism arose from the insertion/deletion of imperfect (CCG)_n repeats within the coding region and gave rise to a variable glycine loop motif in the C-terminal tail of the keratin 10 protein. It is possible that there was a pathogenic role for the expansion of the imperfect trinucleotide repeat.

KRT12,K12

[0116] Keratins are a group of water-insoluble proteins that form 10 nm intermediate filaments in epithelial cells. Approximately 30 different keratin molecules have been identified. They can be divided into acidic and basic-neutral subfamilies according to their relative charges, immunoreactivity, and sequence homologies to types I and II wool keratins, respectively. In vivo, a basic keratin usually is co-expressed and 'paired' with a particular acidic keratin to form a heterodimer. The expression of various keratin pairs is tissue specific, differentiation dependent, and developmentally regulated. The presence of specific keratin pairs is essential for the maintenance of the integrity of epithelium. For example, mutations in human K14/K5 pair and the K10/K1 pair underlie the skin diseases, epidermolysis bullosa

simplex and epidermolytic hyperkeratosis, respectively. Expression of the K3 and K12 keratin pair have been found in the cornea of a wide number of species, including human, mouse, and chicken, and is regarded as a marker for corneal-type epithelial differentiation. The murine Krt12 (Krt1.12) gene and demonstrated that its expression is corneal epithelial cell specific, differentiation dependent, and developmentally regulated [Liu et al., 1993, (70)]. The corneal-specific nature of keratin 12 gene expression signifies keratin 12 plays a unique role in maintaining normal corneal epithelial function. Nevertheless, the exact function of keratin 12 remains unknown and no hereditary human corneal epithelial disorder has been linked directly to the mutation in the keratin 12 gene. As part of a study of the expression profile of human corneal epithelial cells, a cDNA with an open reading frame highly homologous to the cornea-specific mouse keratin 12 gene has been isolated [Nishida et al., 1996, (71)]. To elucidate the function of keratin 12 knockout mice lacking the Krt1.12 gene have been created by gene targeting techniques. The heterozygous mice appeared normal. Homozygous mice developed normally and suffered mild corneal epithelial erosion. The corneal epithelia were fragile and could be removed by gentle rubbing of the eyes or brushing. The corneal epithelium of the homozygotes did not express keratin 12 as judged by immunohistochemistry, Western immunoblot analysis with epitope-specific anti-keratin 12 antibodies, Northern hybridization, and in situ hybridization with an antisense keratin 12 riboprobe. The KRT12 gene has been mapped to 17q by study of radiation hybrids and localized it to the type I keratin cluster in the interval between D17S800 and D17S930 (17q12-q21) [Nishida et al., 1997, (72)]. The authors presented the exon-intron boundary structure of the KRT12 gene and mapped the gene to 17q12 by fluorescence in situ hybridization. The gene contains 7 introns, defining 8 exons that cover the coding sequence. Together the exons and introns span approximately 6 kb of genomic DNA.

[0117] Meesmann corneal dystrophy is an autosomal dominant disorder causing fragility of the anterior corneal epithelium, where the cornea-specific keratins K3 and K12 are expressed. Dominant-negative mutations in these keratins might be the cause of Meesmann corneal dystrophy. Indeed, linkage of the disorder to the K12 locus in Meesmann's original German kindred [Meesmann and Wilke, 1939, (73)] with $Z(\text{max}) = 7.53$ at $\theta = 0.0$ has been found. In 2 pedigrees from Northern Ireland, they found that the disorder co-segregated with K12 in one pedigree and K3 in the other. Heterozygous missense mutations in K3 or in K12 (R135T, V143L,) in each family have been identified. All these mutations occurred in highly conserved keratin helix boundary motifs, where dominant mutations in other keratins have been found to compromise cytoskeletal function severely, leading to keratinocyte fragility.

[0118] The regions of the human KRT12 gene have been sequenced to enable mutation detection for all exons using genomic DNA as a template [Corden et al., 2000, (74)]. The authors found that the human genomic sequence spans 5,919 bp and consists of 8 exons. A microsatellite dinucleotide repeat was identified within intron 3, which was highly polymorphic and which they developed for use in genotype analysis. In addition, 2 mutations in the helix initiation motif of K12 were found in families with Meesmann corneal dystrophy. In an American kindred, a missense M129T mutation was found in the KRT12 gene. They stated that a total of 8 mutations in the KRT12 gene had been reported.

Genetic interactions within ARCHEONs

[0119] Genes involved in genomic alterations (amplifications, insertions, translocations, deletions, etc.) exhibit changes in their expression pattern. Of particular interest are gene amplifications, which account for gene copy numbers >2 per cell or deletions accounting for gene copy numbers <2 per cell. Gene copy number and gene expression of the respective genes do not necessarily correlate. Transcriptional overexpression needs an intact transcriptional context, as determined by regulatory regions at the chromosomal locus (promotor, enhancer and silencer), and sufficient amounts of transcriptional regulators being present in effective combinations. This is especially true for genomic regions, which expression is tightly regulated in specific tissues or during specific developmental stages. ARCHEONs are specified by gene clusters of more than two genes being directly neighboured or in chromosomal order, interspersed by a maximum of 10, preferably 7, more preferably 5 or at least 1 gene. The interspersed genes are also co-amplified but do not directly interact with the ARCHEON. Such an ARCHEON may spread over a chromosomal region of a maximum of 20, more preferably 10 or at least 6 Megabases. The nature of an ARCHEON is characterized by the simultaneous amplification and/or deletion and the correlating expression (i.e. upregulation or downregulation respectively) of the encompassed genes in a specific tissue, cell type, cellular or developmental state or time point. Such ARCHEONs are commonly conserved during evolution, as they play critical roles during cellular development. In case of these ARCHEONs whole gene clusters are overexpressed upon amplification as they harbor self-regulatory feedback loops, which stabilize gene expression and/or biological effector function even in abnormal biological settings, or are regulated by very similar transcription factor combinations, reflecting their simultaneous function in specific tissues at certain developmental stages. Therefore, the gene copy numbers correlates with the expression level especially for genes in gene clusters functioning as ARCHEONs. In case of abnormal gene expressions in neoplastic lesions it is of great importance to know whether the self-regulatory feedback loops have been conserved as they determine the biological activity of the ARCHEON gene members.

[0120] The intensive interaction between genes in ARCHEONs is described for the 17q12 ARCHEON (Fig. 1) by

way of illustration not by limitation. In one embodiment the presence or absence of alterations of genes within distinct genomic regions are correlated with each other, as exemplified for breast cancer cell lines (Fig.3 and Fig. 4). This confers to the discovery of the present invention, that multiple interactions of said gene products of defined chromosomal localizations happen, that according to their respective alterations in abnormal tissue have predictive, diagnostic, prognostic and/or preventive and therapeutic value. These interactions are mediated directly or indirectly, due to the fact that the respective genes are part of interconnected or independent signaling networks or regulate cellular behavior (differentiation status, proliferative and /or apoptotic capacity, invasiveness, drug responsiveness, immune modulatory activities) in a synergistic, antagonistic or independent fashion. The order of functionally important genes within the ARCHEONs has been conserved during evolution (e.g. the ARCHEON on human chromosome 17q12 is present on mouse chromosome 11). Moreover, it has been found that the 17q12 ARCHEON is also present on human chromosome 3p21 and 12q13, both of which are also involved in amplification events and in tumor development. Most probably these homologous ARCHEONs were formed by duplications and rearrangements during vertebrate evolution. Homologous ARCHEONs consist of homologous genes and/or isoforms of specific gene families (e.g. RARA or RARB or RARG, THRA or THRB, TOP2A or TOP2B, RABSA or RABSB, BAF170 or BAF 155, BAF60A or BAF60B, WNT5A or WNT5B, IGFBP4 or IGFBP6). Moreover these regions are flanked by homologous chromosomal gene clusters (e.g. CACN, SCYA, HOX, Keratins). These ARCHEONs have diverged during evolution to fulfill their respective functions in distinct tissues (e.g. the 17q12 ARCHEON has one of its main functions in the central nervous system). Due to their tissue specific function extensive regulatory loops control the expression of the members of each ARCHEON. During tumor development these regulations become critical for the characteristics of the abnormal tissues with respect to differentiation, proliferation, drug responsiveness, invasiveness. It has been found that the co-amplification of genes within ARCHEONs can lead to co-expression of the respective gene products. Some of said genes also exhibit additional mutations or specific patterns of polymorphisms, which are substantial for the oncogenic capacities of these ARCHEONs. It is one of the critical features of such amplicons, which members of the ARCHEON have been conserved during tumor formation (e.g. during amplification and deletion events), thereby defining these genes as diagnostic marker genes. Moreover, the expression of the certain genes within the ARCHEON can be influenced by other members of the ARCHEON, thereby defining the regulatory and regulated genes as target genes for therapeutic intervention. It was also observed, that the expression of certain members of the ARCHEON is sensitive to drug treatment (e.g. TOPO2 alpha, RARA, THRA, HER-2) which defines these genes as "marker genes". Moreover several other genes are suitable for therapeutic intervention by antibodies (CACNB1, EBI1), ligands (CACNB1) or drugs like e.g. kinase inhibitors (CrkRS, CDC6). The following examples of interactions between members of ARCHEONs are offered by way of illustration, not by way of limitation.

[0121] EBI1/CCR7 is lymphoid-specific member of the G protein-coupled receptor family. EBI1 recognizes chemotactants, such as interleukin-8, SCYAs, Rantes, C5a, and fMet-Leu-Phe. The capacity for cell division is largely confined to the CCR7⁺ subsets in lymphocytes. Double-negative cells did not divide or expand after stimulation. CCR7⁺ cells, considered to be terminally differentiated, fail to divide, but do produce interferon-gamma and express high levels of perforin. EBI1 is induced by viral activities such as the Epstein-Barr-Virus. Therefore, EBI1 is associated with transformation events in lymphocytes. A functional role of EBI1 during tumor formation in non-lymphoid tissues has been investigated in this invention. Interestingly, also ERBA and ERBB, located in the same genomic region, are associated with lymphocyte transformation. Moreover, ligands of the receptor (i.e. SCYA5/Rantes) are in genomic proximity on 17q. Abnormal expression of both of these factors in lymphoid and non-lymphoid tissues establishes an autotranslational feedback loop, inducing signaling events within the respective cells. Expression of lymphoid factors has effect on immune cells and modulates cellular behavior. This is of particular interest with regard to abnormal breast tissue being infiltrated by lymphocytes. In line with this, another immunomodulatory and proliferation factor is located nearby on 17q12. Granulocyte colony-stimulating factor (GCSF3) specifically stimulates the proliferation and differentiation of the progenitor cells for granulocytes. A stimulatory activity from a glioblastoma multiforme cell line being biologically and biochemically indistinguishable from GCSF produced by a bladder cell line has also been found. Colony-stimulating factors not only affects immune cells, but also induce cellular responses of non-immune cells, indicating possible involvement in tumor development upon abnormal expression. In addition several other genes of the 17q12 ARCHEON are involved in proliferation, survival, differentiation of immune cells and/or lymphoblastic leukemia, such as MLLT6, ZNF144 and ZNFN1A3, again demonstrating the related functions of the gene products in interconnected key processes within specific cell types. Aberrant expression of more than one of these genes in non-immune cells constitutes signalling activities, that contribute to the oncogenic activities that derive solely from overexpression of the Her-2/neu gene.

[0122] PPARBP has been found in complex with the tumorsuppressor gene of the p53 family. Moreover, PPARBP also binds to PPAR-alpha (PPARA), RAR-alpha (RARA), RXR, THRA and TR-beta-1. Due to it's ability to bind to thyroid hormone receptors it has been named TRIP2 and TRAP220. In this complexes PPARBP affects gene regulatory activities. Interestingly, PPARBP is located in genomic proximity to its interaction partners THRA and RARA. We have found PPARBP to be co-amplified with THRA and RARA in tumor tissue. THRA has been isolated from avian eryth-

roblastosis virus in conjunction with ERBB and therefore was named ERBA. ERBA potentiates ERBB by blocking differentiation of erythroblasts at an immature stage. ERBA has been shown to influence ERBB expression. In this setting deletions of C-terminal portions of the THRA gene product are of influence. Aberrant THRA expression has also been found in nonfunctioning pituitary tumors, which has been hypothesized to reflect mutations in the receptor coding and regulatory sequences. THRA function promotes tumor cell development by regulating gene expression of regulatory genes and by influencing metabolic activities (e.g. of key enzymes of alternative metabolic pathways in tumors such as malic enzyme and genes responsible for lipogenesis). The observed activities of nuclear receptors not only reflect their transactivating potential, but are also due to posttranscriptional activities in the absence or presence of ligands. Co-amplification of THRA /ERBA and ERBB has been shown, but its influence on tumor development has been doubted as no overexpression could be demonstrated in breast tumors [van de Vijver et al., 1987, (75)]. THRA and RARA are part of nuclear receptor family whose function can be mediated as monomers, homodimers or heterodimers. RARA regulates differentiation of a broad spectrum of cells. Interactions of hormones with ERBB expression has been investigated. Ligands of RARA can inhibit the expression of amplified ERBB genes in breast tumors [Offterdinger et al., 1998, (76)]. As being part of this invention co-amplification and co-expression of THRA and RARA could be shown. It was also found that multiple genes, which are regulated by members of the thyroid hormone receptor - and retinoic acid receptor family, are differentially expressed in tumor samples, corresponding to their genomic alterations (amplification, mutation, deletion). These hormone receptor genes and respective target genes are useful to discriminate patient samples with respect to clinical features.

[0123] By expression analysis of multiple normal tissues, tumor samples and tumor cell lines and subsequent clustering of the 17q12 region, it was found that the expression profile of Her-2/neu positive tumor cells and tumor samples exhibits similarities with the expression pattern of tissue from the central nervous system (Fig. 2). This is in line with the observed malformations in the central nervous system of Her-2/neu and THRA knock-out mice. Moreover, it was found that NEUROD2, a nuclear factor involved specifically in neurogenesis, is commonly expressed in the respective samples. This led to the definition of the 17q12 Locus as being an "ARCHEON", whose primary function in normal organ development is defined to the central nervous system. Surprisingly, the expression of NEUROD2 was affected by therapeutic intervention. Strikingly, also ZNF144, TEM7, PIP5K and PPP1R1B are expressed in neuronal cells, where they display diverse tissue specific functions.

[0124] In addition Her-2/neu is often co-amplified with GRB7, a downstream member of the signaling cascade being involved in invasive properties of tumors. Surprisingly, we have found another member of the Her-2/neu signaling cascade being overexpressed in primary breast tumors TOB1 (= "Transducer of ERBB signaling"). Strong overexpression of TOB1 corellated with weaker overexpression of Her-2/neu, already indicating its involvement in oncogenic signaling activities. Amplification of Her-2/neu has been assigned to enhanced proliferative capacity, due to the identified downstream components of the signaling cascade (e.g. Ras-Raf-MAPK). In this respect it was surprising that some cdc genes, which are cell cycle dependent kinases, are part of the amplicons, which upon altered expression have great impact on cell cycle progression.

[0125] According to the observations described above the following examples of genes at 3q21-26 are offered by way of illustration, not by way of limitation.

→ WNT5A, CACNA1D, THRB, RARB, TOP2B, RAB5B, SMARCC1 (BAF155), RAF, WNT7A

[0126] The following examples of genes at 12q13 are offered by way of illustration, not by way of limitation.

→ CACNB3, Keratins, NR4A1, RAB5/13, RARGamma, STAT6, WNT10B, (GCN5), (SAS: Sarcoma Amplified Sequence), SMARCC2 (BAF170), SMARCD1 (BAF60A), (GAS41: Glioma Amplified Sequence), (CHOP), Her3, KRTHB, HOX C, IGFBP6, WNT5B

[0127] There is cross-talk between the amplified ARCHEONS described above and some other highly amplified genomic regions locate approximately at 1p13, 1q32, 2p16, 2q21, 3p12, 5p13, 6p12, 7p12, 7q21, 8q23, 11q13, 13q12, 19q13, 20q13 and 21q11. The above mentioned chromosomal regions are described by way of illustration not by way of limitation, as the amplified regions often span larger and/or overlapping positions at these chromosomal positions.

[0128] Additional alterations of non-transcribed genes, pseudogenes or intergenic regions of said chromosomal locations can be measured for prediction, diagnosis, prognosis, prevention and treatment of malignant neoplasia and breast cancer in particular. Some of the genes or genomic regions have no direct influence on the members of the ARCHEONS or the genes within distinct chromosomal regions but still retain marker gene function due to their chromosomal positioning in the neighborhood of functionally critical genes (e.g. Telethonin neighboring the Her-2/neu gene).

[0129] The invention further relates to the use of:

a) a polynucleotide comprising at least one of the sequences of SEQ ID NO: 1 to 26 or 53 to 75;

b) a polynucleotide which hybridizes under stringent conditions to a polynucleotide specified in (a) encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 2 or 3

c) a polynucleotide the sequence of which deviates from the polynucleotide specified in (a) and (b) due to the generation of the genetic code encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 2 or 3

d) a polynucleotide which represents a specific fragment, derivative or allelic variation of a polynucleotide sequence specified in (a) to (c)

e) an antisense molecule targeting specifically one of the polynucleotide sequences specified in (a) to (d);

f) a purified polypeptide encoded by a polynucleotide sequence specified in (a) to (d)

g) a purified polypeptide comprising at least one of the sequences of SEQ ID NO: 27 to 52 or 76 to 98;

h) an antibody capable of binding to one of the polynucleotide specified in (a) to (d) or a polypeptide specified in (f) and (g)

i) a reagent identified by any of the methods of claim 14 to 16 that modulates the amount or activity of a polynucleotide sequence specified in (a) to (d) or a polypeptide specified in (f) and (g)

in the preparation of a composition for the prevention, prediction, diagnosis, prognosis or a medicament for the treatment of malignant neoplasia and breast cancer in particular.

Polynucleotides

[0130] A "BREAST CANCER GENE" polynucleotide can be single- or double-stranded and comprises a coding sequence or the complement of a coding sequence for a "BREAST CANCER GENE" polypeptide. Degenerate nucleotide sequences encoding human "BREAST CANCER GENE" polypeptides, as well as homologous nucleotide sequences which are at least about 50, 55, 60, 65, 70, preferably about 75, 90, 96, or 98% identical to the nucleotide sequences of SEQ ID NO: 1 to 26 or 53 to 75 also are "BREAST CANCER GENE" polynucleotides. Percent sequence identity between the sequences of two polynucleotides is determined using computer programs such as ALIGN which employ the FASTA algorithm, using an affine gap search with a gap open penalty of -12 and a gap extension penalty of -2. Complementary DNA (cDNA) molecules, species homologues, and variants of "BREAST CANCER GENE" polynucleotides which encode biologically active "BREAST CANCER GENE" polypeptides also are "BREAST CANCER GENE" polynucleotides.

Preparation of Polynucleotides

[0131] A naturally occurring "BREAST CANCER GENE" polynucleotide can be isolated free of other cellular components such as membrane components, proteins, and lipids. Polynucleotides can be made by a cell and isolated using standard nucleic acid purification techniques, or synthesized using an amplification technique, such as the polymerase chain reaction (PCR), or by using an automatic synthesizer. Methods for isolating polynucleotides are routine and are known in the art. Any such technique for obtaining a polynucleotide can be used to obtain isolated "BREAST CANCER GENE" polynucleotides. For example, restriction enzymes and probes can be used to isolate polynucleotide fragments which comprises "BREAST CANCER GENE" nucleotide sequences. Isolated polynucleotides are in preparations which are free or at least 70, 80, or 90% free of other molecules.

[0132] "BREAST CANCER GENE" cDNA molecules can be made with standard molecular biology techniques, using "BREAST CANCER GENE" mRNA as a template. Any RNA isolation technique which does not select against the isolation of mRNA may be utilized for the purification of such RNA samples. See, for example, Sambrook et al., 1989, (77); and Ausubel, F. M. et al., 1989, (78), both of which are incorporated herein by reference in their entirety. Additionally, large numbers of tissue samples may readily be processed using techniques well known to those of skill in the art, such as, for example, the single-step RNA isolation process of Chomczynski, P. (1989, U.S. Pat. No. 4,843,155), which is incorporated herein by reference in its entirety.

[0133] "BREAST CANCER GENE" cDNA molecules can thereafter be replicated using molecular biology techniques known in the art and disclosed in manuals such as Sambrook et al., 1989, (77). An amplification technique, such as PCR, can be used to obtain additional copies of polynucleotides of the invention, using either human genomic DNA or

cDNA as a template.

[0134] Alternatively, synthetic chemistry techniques can be used to synthesize "BREAST CANCER GENE" polynucleotides. The degeneracy of the genetic code allows alternate nucleotide sequences to be synthesized which will encode a "BREAST CANCER GENE" polypeptide or a biologically active variant thereof.

Identification of differential expression

[0135] Transcripts within the collected RNA samples which represent RNA produced by differentially expressed genes may be identified by utilizing a variety of methods which are well known to those of skill in the art. For example, differential screening [Tedder, T. F. et al., 1988, (79)], subtractive hybridization [Hedrick, S. M. et al., 1984, (80); Lee, S. W. et al., 1984, (81)], and, preferably, differential display (Liang, P., and Pardee, A. B., 1993, U.S. Pat. No. 5,262,311, which is incorporated herein by reference in its entirety), may be utilized to identify polynucleotide sequences derived from genes that are differentially expressed.

[0136] Differential screening involves the duplicate screening of a cDNA library in which one copy of the library is screened with a total cell cDNA probe corresponding to the mRNA population of one cell type while a duplicate copy of the cDNA library is screened with a total cDNA probe corresponding to the mRNA population of a second cell type. For example, one cDNA probe may correspond to a total cell cDNA probe of a cell type derived from a control subject, while the second cDNA probe may correspond to a total cell cDNA probe of the same cell type derived from an experimental subject. Those clones which hybridize to one probe but not to the other potentially represent clones derived from genes differentially expressed in the cell type of interest in control versus experimental subjects.

[0137] Subtractive hybridization techniques generally involve the isolation of mRNA taken from two different sources, e.g., control and experimental tissue, the hybridization of the mRNA or single-stranded cDNA reverse-transcribed from the isolated mRNA, and the removal of all hybridized, and therefore double-stranded, sequences. The remaining non-hybridized, single-stranded cDNAs, potentially represent clones derived from genes that are differentially expressed in the two mRNA sources. Such single-stranded cDNAs are then used as the starting material for the construction of a library comprising clones derived from differentially expressed genes.

[0138] The differential display technique describes a procedure, utilizing the well known polymerase chain reaction (PCR); the experimental embodiment set forth in Mullis, K. B., 1987, U.S. Pat. No. 4,683,202 which allows for the identification of sequences derived from genes which are differentially expressed. First, isolated RNA is reverse-transcribed into single-stranded cDNA, utilizing standard techniques which are well known to those of skill in the art. Primers for the reverse transcriptase reaction may include, but are not limited to, oligo dT-containing primers, preferably of the reverse primer type of oligonucleotide described below. Next, this technique uses pairs of PCR primers, as described below, which allow for the amplification of clones representing a random subset of the RNA transcripts present within any given cell. Utilizing different pairs of primers allows each of the mRNA transcripts present in a cell to be amplified. Among such amplified transcripts may be identified those which have been produced from differentially expressed genes.

[0139] The reverse oligonucleotide primer of the primer pairs may contain an oligo dT stretch of nucleotides, preferably eleven nucleotides long, at its 5' end, which hybridizes to the poly(A) tail of mRNA or to the complement of a cDNA reverse transcribed from an mRNA poly(A) tail. Second, in order to increase the specificity of the reverse primer, the primer may contain one or more, preferably two, additional nucleotides at its 3' end. Because, statistically, only a subset of the mRNA derived sequences present in the sample of interest will hybridize to such primers, the additional nucleotides allow the primers to amplify only a subset of the mRNA derived sequences present in the sample of interest. This is preferred in that it allows more accurate and complete visualization and characterization of each of the bands representing amplified sequences.

[0140] The forward primer may contain a nucleotide sequence expected, statistically, to have the ability to hybridize to cDNA sequences derived from the tissues of interest. The nucleotide sequence may be an arbitrary one, and the length of the forward oligonucleotide primer may range from about 9 to about 13 nucleotides, with about 10 nucleotides being preferred. Arbitrary primer sequences cause the lengths of the amplified partial cDNAs produced to be variable, thus allowing different clones to be separated by using standard denaturing sequencing gel electrophoresis. PCR reaction conditions should be chosen which optimize amplified product yield and specificity, and, additionally, produce amplified products of lengths which may be resolved utilizing standard gel electrophoresis techniques. Such reaction conditions are well known to those of skill in the art, and important reaction parameters include, for example, length and nucleotide sequence of oligonucleotide primers as discussed above, and annealing and elongation step temperatures and reaction times. The pattern of clones resulting from the reverse transcription and amplification of the mRNA of two different cell types is displayed via sequencing gel electrophoresis and compared. Differences in the two banding patterns indicate potentially differentially expressed genes.

[0141] When screening for full-length cDNAs, it is preferable to use libraries that have been size-selected to include larger cDNAs. Randomly-primed libraries are preferable, in that they will contain more sequences which contain the

5' regions of genes. Use of a randomly primed library may be especially preferable for situations in which an oligo d(T) library does not yield a full-length cDNA. Genomic libraries can be useful for extension of sequence into 5' non-transcribed regulatory regions.

[0142] Commercially available capillary electrophoresis systems can be used to analyze the size or confirm the nucleotide sequence of PCR or sequencing products. For example, capillary sequencing can employ flowable polymers for electrophoretic separation, four different fluorescent dyes (one for each nucleotide) which are laser activated, and detection of the emitted wavelengths by a charge coupled device camera. Output/light intensity can be converted to electrical signal using appropriate software (e.g. GENOTYPER and Sequence NAVIGATOR, Perkin Elmer; ABI), and the entire process from loading of samples to computer analysis and electronic data display can be computer controlled.

Capillary electrophoresis is especially preferable for the sequencing of small pieces of DNA which might be present in limited amounts in a particular sample.

[0143] Once potentially differentially expressed gene sequences have been identified via bulk techniques such as, for example, those described above, the differential expression of such putatively differentially expressed genes should be corroborated. Corroboration may be accomplished via, for example, such well known techniques as Northern analysis and/or RT-PCR. Upon corroboration, the differentially expressed genes may be further characterized, and may be identified as target and/or marker genes, as discussed, below.

[0144] Also, amplified sequences of differentially expressed genes obtained through, for example, differential display may be used to isolate full length clones of the corresponding gene. The full length coding portion of the gene may readily be isolated, without undue experimentation, by molecular biological techniques well known in the art. For example, the isolated differentially expressed amplified fragment may be labeled and used to screen a cDNA library. Alternatively, the labeled fragment may be used to screen a genomic library.

[0145] An analysis of the tissue distribution of the mRNA produced by the identified genes may be conducted, utilizing standard techniques well known to those of skill in the art. Such techniques may include, for example, Northern analyses and RT-PCR. Such analyses provide information as to whether the identified genes are expressed in tissues expected to contribute to breast cancer. Such analyses may also provide quantitative information regarding steady state mRNA regulation, yielding data concerning which of the identified genes exhibits a high level of regulation in, preferably, tissues which may be expected to contribute to breast cancer.

[0146] Such analyses may also be performed on an isolated cell population of a particular cell type derived from a given tissue. Additionally, standard in situ hybridization techniques may be utilized to provide information regarding which cells within a given tissue express the identified gene. Such analyses may provide information regarding the biological function of an identified gene relative to breast cancer in instances wherein only a subset of the cells within the tissue is thought to be relevant to breast cancer.

Identification of co-amplified genes

[0147] Genes involved in genomic alterations (amplifications, insertions, translocations, deletions, etc.) are identified by PCR-based karyotyping in combination with database analysis. Of particular interest are gene amplifications, which account for gene copy numbers >2 per cell. Gene copy number and gene expression of the respective genes often correlates. Therefore clusters of genes being simultaneously overexpressed due to gene amplifications can be identified by expression analysis via DNA-chip technologies or quantitative RTPCR. For example, the altered expression of genes due to increased or decreased gene copy numbers can be determined by GeneArray™ technologies from Affymetrix or qRT-PCR with the TaqMan or iCycler Systems. Moreover combination of RNA with DNA analytic enables highly parallel and automated characterization of multiple genomic regions of variable length with high resolution in tissue or single cell samples. Furthermore these assays enable the correlation of gene transcription relative to gene copy number of target genes. As there is not necessarily a linear correlation of expression level and gene copy number and as there are synergistic or antagonistic effects in certain gene clusters, the identification on the RNA-level is easier and probably more relevant for the biological outcome of the alterations especially in tumor tissue.

Detection of co-amplified genes in malignant neoplasia

[0148] Chromosomal changes are commonly detected by FISH (=Fluorescence-In-Situ-Hybridization) and CGH (=Comparative Genomic Hybridization). For quantification of genomic regions genes or intergenic regions can be used. Such quantification measures the relative abundance of multiple genes with respect to each other (e.g. target gene vs. centromeric region or housekeeping genes). Changes in relative abundance can be detected in paraffin-embedded material even after extraction of RNA or genomic DNA. Measurement of genomic DNA has advantages compared to RNA-analysis due to the stability of DNA, which accounts for the possibility to perform also retrospective studies and offers multiple internal controls (genes not being altered, amplified or deleted) for standardization and exact calculations. Moreover, PCR-analysis of genomic DNA offers the advantage to investigate intergenic, highly variable regions

or combinations of SNP's (=Single Nucleotide Polymorphisms), RFLPs, VNTRs and STRs (in general polymorphic markers). Determination of SNPs or polymorphic markers within defined genomic regions (e.g. SNP analysis by "Pyrosequencing™") has impact on the phenotype of the genomic alterations. For example it is of advantage to determine combinations of polymorphisms or haplotypes in order to characterize the biological potential of genes being part of amplified alleles. Of particular interest are polymorphic markers in breakpoint regions, coding regions or regulatory regions of genes or intergenic regions. By determining predictive haplotypes with defined biological or clinical outcome it is possible to establish diagnostic and prognostic assays with non-tumor samples from patients. Depending on whether preferably one allele or both alleles to same extent are amplified (= linear or non-linear amplifications) haplotypes can be determined. Overrepresentation of specific polymorphic markers combinations in cells or tissues with gene amplifications facilitates haplotype determination, as e.g. combinations of heterozygous polymorphic markers in nucleic acids isolated from normal tissues, body fluids or biological samples of one patient become almost homozygous in neoplastic tissue of the very same patient. This "gain of homozygosity" corresponds to the measurement of altered genomic region due to amplification events and is suitable for identification of "gain of function"- alterations in tumors, which result in e.g. oncogenic or growth promoting activities. In contrast, the detection of "losses of heterozygosity" is used for identification of anti-oncogenes, gate keeper genes or checkpoint genes, that suppress oncogenic activities and negatively regulate cellular growth processes. This intrinsic difference clearly opposes the impact of the respective genomic regions for tumor development and emphasizes the significance of "gain of homozygosity" measurements disclosed in this invention. In addition to the analyses on SNPs, a comparative approach of blood leucocyte DNA and tumor DNA based on VNTR detection can reveal the existence of a formerly described ARCHEON. SNP and VNTR sequences and primer sets most suitable for detection of the ARCHEON at 17q11-21 are disclosed in Table 4 and Table 6. Detection, quantification and sizing of such polymorphic markers can be achieved by methods known to those with skill in the art. In one embodiment of this invention we disclose the comparative measurement of amount and size of any of the disclosed VNTRs (Table 6) by PCR amplification and capillary electrophoresis. PCR can be carried out by standard protocols favorably in a linear amplification range (low cycle number) and detection by CE should be carried out by suppliers protocols (e.g. Agilent). More favorably the detection of the VNTRs disclosed in Table 6 can be carried out in a multiplex fashion, utilizing a variety of labeled primers (e.g. fluorescent, radioactive, bioactive) and a suitable CE detection system (e.g. ABI 310). However the detection can also be performed on slab gels consisting of highly concentrated agarose or polyacrylamide with a monochromal DNA stain. Enhancement of resolution can be achieved by appropriate primer design and length variation to give best results in multiplex PCR.

[0149] It is also of interest to determine covalent modifications of DNA (e.g. methylation) or the associated chromatin (e.g. acetylation or methylation of associated proteins) within the altered genomic regions, that have impact on transcriptional activity of the genes. In general, by measuring multiple, short sequences (60-300 bp) these techniques enable high-resolution analysis of target regions, which cannot be obtained by conventional methods such as FISH analytic (2-100 kb). Moreover the PCR-based DNA analysis techniques offer advantages with regard to sensitivity, specificity, multiplexing, time consumption and low amount of patient material required. These techniques can be optimized by combination with microdissection or macrodissection to obtain purer starting material for analysis.

Extending Polynucleotides

[0150] In one embodiment of such a procedure for the identification and cloning of full length gene sequences, RNA may be isolated, following standard procedures, from an appropriate tissue or cellular source. A reverse transcription reaction may then be performed on the RNA using an oligonucleotide primer complementary to the mRNA that corresponds to the amplified fragment, for the priming of first strand synthesis. Because the primer is anti-parallel to the mRNA, extension will proceed toward the 5' end of the mRNA. The resulting RNA hybrid may then be "tailed" with guanines using a standard terminal transferase reaction, the hybrid may be digested with RNase H, and second strand synthesis may then be primed with a poly-C primer. Using the two primers, the 5' portion of the gene is amplified using PCR. Sequences obtained may then be isolated and recombined with previously isolated sequences to generate a full-length cDNA of the differentially expressed genes of the invention. For a review of cloning strategies and recombinant DNA techniques, see e.g., Sambrook et al., (77); and Ausubel et al., (78).

[0151] Various PCR-based methods can be used to extend the polynucleotide sequences disclosed herein to detect upstream sequences such as promoters and regulatory elements. For example, restriction site PCR uses universal primers to retrieve unknown sequence adjacent to a known locus [Sarkar, 1993, (82)]. Genomic DNA is first amplified in the presence of a primer to a linker sequence and a primer specific to the known region. The amplified sequences are then subjected to a second round of PCR with the same linker primer and another specific primer internal to the first one. Products of each round of PCR are transcribed with an appropriate RNA polymerase and sequenced using reverse transcriptase.

[0152] Inverse PCR also can be used to amplify or extend sequences using divergent primers based on a known region [Triglia et al., 1988, (83)]. Primers can be designed using commercially available software, such as OLIGO 4.06

Primer Analysis software (National Biosciences Inc., Plymouth, Minn.), to be e.g. 2230 nucleotides in length, to have a GC content of 50% or more, and to anneal to the target sequence at temperatures about 68-72°C. The method uses several restriction enzymes to generate a suitable fragment in the known region of a gene. The fragment is then circularized by intramolecular ligation and used as a PCR template.

[0153] Another method which can be used is capture PCR, which involves PCR amplification of DNA fragments adjacent to a known sequence in human and yeast artificial chromosome DNA [Lagerstrom et al., 1991, (84)]. In this method, multiple restriction enzyme digestions and ligations also can be used to place an engineered double-stranded sequence into an unknown fragment of the DNA molecule before performing PCR.

[0154] Additionally, PCR, nested primers, and PROMOTERFINDER libraries (CLONTECH, Palo Alto, Calif.) can be used to walk genomic DNA (CLONTECH, Palo Alto, Calif.). This process avoids the need to screen libraries and is useful in finding intron/exon junctions.

[0155] The sequences of the identified genes may be used, utilizing standard techniques, to place the genes onto genetic maps, e.g., mouse [Copeland & Jenkins, 1991, (85)] and human genetic maps [Cohen, et al., 1993, (86)]. Such mapping information may yield information regarding the genes' importance to human disease by, for example, identifying genes which map near genetic regions to which known genetic breast cancer tendencies map.

Identification of polynucleotide variants and homologues or splice variants

[0156] Variants and homologues of the "BREAST CANCER GENE" polynucleotides described above also are "BREAST CANCER GENE" polynucleotides. Typically, homologous "BREAST CANCER GENE" polynucleotide sequences can be identified by hybridization of candidate polynucleotides to known "BREAST CANCER GENE" polynucleotides under stringent conditions, as is known in the art. For example, using the following wash conditions: 2 X SSC (0.3 M NaCl, 0.03 M sodium citrate, pH 7.0), 0.1% SDS, room temperature twice, 30 minutes each; then 2 X SSC, 0.1% SDS, 50 EC once, 30 minutes; then 2 X SSC, room temperature twice, 10 minutes each homologous sequences can be identified which contain at most about 25-30% basepair mismatches. More preferably, homologous polynucleotide strands contain 15-25% basepair mismatches, even more preferably 5-15% basepair mismatches.

[0157] Species homologues of the "BREAST CANCER GENE" polynucleotides disclosed herein also can be identified by making suitable probes or primers and screening cDNA expression libraries from other species, such as mice, monkeys, or yeast. Human variants of "BREAST CANCER GENE" polynucleotides can be identified, for example, by screening human cDNA expression libraries. It is well known that the T_m of a double-stranded DNA decreases by 1-1.5°C with every 1% decrease in homology [Bonner et al., 1973, (87)]. Variants of human "BREAST CANCER GENE" polynucleotides or "BREAST CANCER GENE" polynucleotides of other species can therefore be identified by hybridizing a putative homologous "BREAST CANCER GENE" polynucleotide with a polynucleotide having a nucleotide sequence of one of the sequences of the SEQ ID NO: 1 to 26 or 53 to 75 or the complement thereof to form a test hybrid. The melting temperature of the test hybrid is compared with the melting temperature of a hybrid comprising polynucleotides having perfectly complementary nucleotide sequences, and the number or percent of basepair mismatches within the test hybrid is calculated.

[0158] Nucleotide sequences which hybridize to "BREAST CANCER GENE" polynucleotides or their complements following stringent hybridization and/or wash conditions also are "BREAST CANCER GENE" polynucleotides. Stringent wash conditions are well known and understood in the art and are disclosed, for example, in Sambrook et al., (77). Typically, for stringent hybridization conditions a combination of temperature and salt concentration should be chosen that is approximately 12-20°C below the calculated T_m of the hybrid under study. The T_m of a hybrid between a "BREAST CANCER GENE" polynucleotide having a nucleotide sequence of one of the sequences of the SEQ ID NO: 1 to 26 or 53 to 75 or the complement thereof and a polynucleotide sequence which is at least about 50, preferably about 75, 90, 96, or 98% identical to one of those nucleotide sequences can be calculated, for example, using the equation below [Bolton and McCarthy, 1962, (88)]:

$$T_m = 81.5^\circ\text{C} - 16.6(\log_{10}[\text{Na}^+]) + 0.41(\%G + C) - 0.63(\%\text{formamide}) - 600/1,$$

where 1 = the length of the hybrid in basepairs.

[0159] Stringent wash conditions include, for example, 4 X SSC at 65°C, or 50% formamide, 4 X SSC at 28°C, or 0.5 X SSC, 0.1% SDS at 65°C. Highly stringent wash conditions include, for example, 0.2 X SSC at 65°C.

[0160] The biological function of the identified genes may be more directly assessed by utilizing relevant *in vivo* and *in vitro* systems. *In vivo* systems may include, but are not limited to, animal systems which naturally exhibit breast cancer predisposition, or ones which have been engineered to exhibit such symptoms, including but not limited to the apoE-deficient malignant neoplasia mouse model [Plump et al., 1992, (89)].

[0161] Splice variants derived from the same genomic region, encoded by the same pre mRNA can be identified by

hybridization conditions described above for homology search. The specific characteristics of variant proteins encoded by splice variants of the same pre transcript may differ and can also be assayed as disclosed. A "BREAST CANCER GENE" polynucleotide having a nucleotide sequence of one of the sequences of the SEQ ID NO: 1 to 26 or 53 to 75 or the complement thereof may therefor differ in parts of the entire sequence as presented for SEQ ID NO: 60 and the encoded splice variants SEQ ID NO: 61 to 66. These refer to individual proteins SEQ ID NO: 83 to 89. The prediction of splicing events and the identification of the utilized acceptor and donor sites within the pre mRNA can be computed (e.g. Software Package GRAIL or GenomeSCAN) and verified by PCR method by those with skill in the art.

Antisense oligonucleotides

[0162] Antisense oligonucleotides are nucleotide sequences which are complementary to a specific DNA or RNA sequence. Once introduced into a cell, the complementary nucleotides combine with natural sequences produced by the cell to form complexes and block either transcription or translation. Preferably, an antisense oligonucleotide is at least 6 nucleotides in length, but can be at least 7, 8, 10, 12, 15, 20, 25, 30, 35, 40, 45, or 50 or more nucleotides long. Longer sequences also can be used. Antisense oligonucleotide molecules can be provided in a DNA construct and introduced into a cell as described above to decrease the level of "BREAST CANCER GENE" gene products in the cell.

[0163] Antisense oligonucleotides can be deoxyribonucleotides, ribonucleotides, peptide nucleic acids (PNAs; described in U.S. Pat. No. 5,714,331), locked nucleic acids (LNAs; described in WO 99/12826), or a combination of them. Oligonucleotides can be synthesized manually or by an automated synthesizer, by covalently linking the 5' end of one nucleotide with the 3' end of another nucleotide with non-phosphodiester internucleotide linkages such as alkylphosphonates, phosphorothioates, phosphorodithioates, alkylphosphonothioates, alkylphosphonates, phosphoramidates, phosphate esters, carbamates, acetamides, carboxymethyl esters, carbonates, and phosphate triesters [Brown, 1994, (126); Sonveaux, 1994, (127) and Uhlmann et al., 1990, (128)].

[0164] Modifications of "BREAST CANCER GENE" expression can be obtained by designing antisense oligonucleotides which will form duplexes to the control, 5', or regulatory regions of the "BREAST CANCER GENE". Oligonucleotides derived from the transcription initiation site, e.g., between positions 10 and +10 from the start site, are preferred. Similarly, inhibition can be achieved using "triple helix" base-pairing methodology. Triple helix pairing is useful because it causes inhibition of the ability of the double helix to open sufficiently for the binding of polymerases, transcription factors, or chaperons. Therapeutic advances using triplex DNA have been described in the literature [Gee et al., 1994, (129)]. An antisense oligonucleotide also can be designed to block translation of mRNA by preventing the transcript from binding to ribosomes.

[0165] Precise complementarity is not required for successful complex formation between an antisense oligonucleotide and the complementary sequence of a "BREAST CANCER GENE" polynucleotide. Antisense oligonucleotides which comprise, for example, 2, 3, 4, or 5 or more stretches of contiguous nucleotides which are precisely complementary to a "BREAST CANCER GENE" polynucleotide, each separated by a stretch of contiguous nucleotides which are not complementary to adjacent "BREAST CANCER GENE" nucleotides, can provide sufficient targeting specificity for "BREAST CANCER GENE" mRNA. Preferably, each stretch of complementary contiguous nucleotides is at least 4, 5, 6, 7, or 8 or more nucleotides in length. Non-complementary intervening sequences are preferably 1, 2, 3, or 4 nucleotides in length. One skilled in the art can easily use the calculated melting point of an antisense-sense pair to determine the degree of mismatching which will be tolerated between a particular antisense oligonucleotide and a particular "BREAST CANCER GENE" polynucleotide sequence.

[0166] Antisense oligonucleotides can be modified without affecting their ability to hybridize to a "BREAST CANCER GENE" polynucleotide. These modifications can be internal or at one or both ends of the antisense molecule. For example, inter-nucleoside phosphate linkages can be modified by adding cholesteryl or diamine moieties with varying numbers of carbon residues between the amino groups and terminal ribose. Modified bases and/or sugars, such as arabinose instead of ribose, or a 3', 5' substituted oligonucleotide in which the 3' hydroxyl group or the 5' phosphate group are substituted, also can be employed in a modified antisense oligonucleotide. These modified oligonucleotides can be prepared by methods well known in the art [Agrawal et al., 1992, (130); Uhlmann et al., 1987, (131) and Uhlmann et al., (128)].

Ribozymes

[0167] Ribozymes are RNA molecules with catalytic activity [Cech, 1987, (132); Cech, 1990, (133) and Couture & Stinchcomb, 1996, (134)]. Ribozymes can be used to inhibit gene function by cleaving an RNA sequence, as is known in the art (e.g., Haseloff et al., U.S. Patent 5,641,673). The mechanism of ribozyme action involves sequence-specific hybridization of the ribozyme molecule to complementary target RNA, followed by endonucleolytic cleavage. Examples include engineered hammerhead motif ribozyme molecules that can specifically and efficiently catalyze endonucleolytic cleavage of specific nucleotide sequences.

[0168] The transcribed sequence of a "BREAST CANCER GENE" can be used to generate ribozymes which will specifically bind to mRNA transcribed from a "BREAST CANCER GENE" genomic locus. Methods of designing and constructing ribozymes which can cleave other RNA molecules in trans in a highly sequence specific manner have been developed and described in the art [Haseloff et al., 1988, (135)]. For example, the cleavage activity of ribozymes can be targeted to specific RNAs by engineering a discrete "hybridization" region into the ribozyme. The hybridization region contains a sequence complementary to the target RNA and thus specifically hybridizes with the target [see, for example, Gerlach et al., EP 0 321201].

[0169] Specific ribozyme cleavage sites within a "BREAST CANCER GENE" RNA target can be identified by scanning the target molecule for ribozyme cleavage sites which include the following sequences: GUA, GUU, and GUC. Once identified, short RNA sequences of between 15 and 20 ribonucleotides corresponding to the region of the target RNA containing the cleavage site can be evaluated for secondary structural features which may render the target inoperable. Suitability of candidate "BREAST CANCER GENE" RNA targets also can be evaluated by testing accessibility to hybridization with complementary oligonucleotides using ribonuclease protection assays. Longer complementary sequences can be used to increase the affinity of the hybridization sequence for the target. The hybridizing and cleavage regions of the ribozyme can be integrally related such that upon hybridizing to the target RNA through the complementary regions, the catalytic region of the ribozyme can cleave the target.

[0170] Ribozymes can be introduced into cells as part of a DNA construct. Mechanical methods, such as microinjection, liposome-mediated transfection, electroporation, or calcium phosphate precipitation, can be used to introduce a ribozyme-containing DNA construct into cells in which it is desired to decrease "BREAST CANCER GENE" expression. Alternatively, if it is desired that the cells stably retain the DNA construct, the construct can be supplied on a plasmid and maintained as a separate element or integrated into the genome of the cells, as is known in the art. A ribozyme-encoding DNA construct can include transcriptional regulatory elements, such as a promoter element, an enhancer or UAS element, and a transcriptional terminator signal, for controlling transcription of ribozymes in the cells.

[0171] As taught in Haseloff et al., U.S. Pat. No. 5,641,673, ribozymes can be engineered so that ribozyme expression will occur in response to factors which induce expression of a target gene. Ribozymes also can be engineered to provide an additional level of regulation, so that destruction of mRNA occurs only when both a ribozyme and a target gene are induced in the cells.

Polypeptides

[0172] "BREAST CANCER GENE" polypeptides according to the invention comprise an polypeptide selected from SEQ ID NO: 27 to 52 and 76 to 98 or encoded by any of the polynucleotide sequences of the SEQ ID NO: 1 to 26 and 53 to 75 or derivatives, fragments, analogues and homologues thereof. A "BREAST CANCER GENE" polypeptide of the invention therefore can be a portion, a full-length, or a fusion protein comprising all or a portion of a "BREAST CANCER GENE" polypeptide.

Protein Purification

[0173] "BREAST CANCER GENE" polypeptides can be purified from any cell which expresses the enzyme, including host cells which have been transfected with "BREAST CANCER GENE" expression constructs. Breast tissue is an especially useful source of "BREAST CANCER GENE" polypeptides. A purified "BREAST CANCER GENE" polypeptide is separated from other compounds which normally associate with the "BREAST CANCER GENE" polypeptide in the cell, such as certain proteins, carbohydrates, or lipids, using methods well-known in the art. Such methods include, but are not limited to, size exclusion chromatography, ammonium sulfate fractionation, ion exchange chromatography, affinity chromatography, and preparative gel electrophoresis. A preparation of purified "BREAST CANCER GENE" polypeptides is at least 80% pure; preferably, the preparations are 90%, 95%, or 99% pure. Purity of the preparations can be assessed by any means known in the art, such as SDS-polyacrylamide gel electrophoresis.

Obtaining Polypeptides

[0174] "BREAST CANCER GENE" polypeptides can be obtained, for example, by purification from human cells, by expression of "BREAST CANCER GENE" polynucleotides, or by direct chemical synthesis.

Biologically Active Variants

[0175] "BREAST CANCER GENE" polypeptide variants which are biologically active, i.e., retain an "BREAST CANCER GENE" activity, also are "BREAST CANCER GENE" polypeptides. Preferably, naturally or non-naturally occurring "BREAST CANCER GENE" polypeptide variants have amino acid sequences which are at least about 60, 65, or 70,

preferably about 75, 80, 85, 90, 92, 94, 96, or 98% identical to the any of the amino acid sequences of the polypeptides of SEQ ID NO: 27 to 52 or 76 to 98 or the polypeptides encoded by any of the polynucleotides of SEQ ID NO: 1 to 26 or 53 to 75 or a fragment thereof. Percent identity between a putative "BREAST CANCER GENE" polypeptide variant and of the polypeptides of SEQ ID NO: 27 to 52 or 76 to 98 or the polypeptides encoded by any of the polynucleotides of SEQ ID NO: 1 to 26 or 53 to 75 or a fragment thereof is determined by conventional methods. [See, for example, 5 Altschul *et al.*, 1986, (90) and Henikoff & Henikoff, 1992, (91)]. Briefly, two amino acid sequences are aligned to optimize the alignment scores using a gap opening penalty of 10, a gap extension penalty of 1, and the "BLOSUM62" scoring matrix of Henikoff & Henikoff, (91).

[0176] Those skilled in the art appreciate that there are many established algorithms available to align two amino acid sequences. The "FASTA" similarity search algorithm of Pearson & Lipman is a suitable protein alignment method for examining the level of identity shared by an amino acid sequence disclosed herein and the amino acid sequence of a putative variant [Pearson & Lipman, 1988, (92), and Pearson, 1990, (93)]. Briefly, FASTA first characterizes sequence similarity by identifying regions shared by the query sequence (e.g., SEQ ID NO: 1 to 26 or 53 to 75) and a test sequence that have either the highest density of identities (if the ktup variable is 1) or pairs of identities (if ktup=2), 10 without considering conservative amino acid substitutions, insertions, or deletions. The ten regions with the highest density of identities are then rescored by comparing the similarity of all paired amino acids using an amino acid substitution matrix, and the ends of the regions are "trimmed" to include only those residues that contribute to the highest score. If there are several regions with scores greater than the "cutoff" value (calculated by a predetermined formula based upon the length of the sequence the ktup value), then the trimmed initial regions are examined to determine 20 whether the regions can be joined to form an approximate alignment with gaps. Finally, the highest scoring regions of the two amino acid sequences are aligned using a modification of the Needleman-Wunsch-Sellers algorithm [Needleman & Wunsch, 1970, (94), and Sellers, 1974, (95)], which allows for amino acid insertions and deletions. Preferred parameters for FASTA analysis are: ktup=1, gap opening penalty=10, gap extension penalty=1, and substitution matrix=BLOSUM62. These parameters can be introduced into a FASTA program by modifying the scoring matrix file ("SMATRIX"), as explained in Appendix 2 of Pearson, (93).

[0177] FASTA can also be used to determine the sequence identity of nucleic acid molecules using a ratio as disclosed above. For nucleotide sequence comparisons, the ktup value can range between one to six, preferably from three to six, most preferably three, with other parameters set as default.

[0178] Variations in percent identity can be due, for example, to amino acid substitutions, insertions, or deletions. Amino acid substitutions are defined as one for one amino acid replacements. They are conservative in nature when the substituted amino acid has similar structural and/or chemical properties. Examples of conservative replacements are substitution of a leucine with an isoleucine or valine, an aspartate with a glutamate, or a threonine with a serine.

[0179] Amino acid insertions or deletions are changes to or within an amino acid sequence. They typically fall in the range of about 1 to 5 amino acids. Guidance in determining which amino acid residues can be substituted, inserted, 35 or deleted without abolishing biological or immunological activity of a "BREAST CANCER GENE" polypeptide can be found using computer programs well known in the art, such as DNASTAR software. Whether an amino acid change results in a biologically active "BREAST CANCER GENE" polypeptide can readily be determined by assaying for "BREAST CANCER GENE" activity, as described for example, in the specific Examples, below. Larger insertions or deletions can also be caused by alternative splicing. Protein domains can be inserted or deleted without altering the 40 main activity of the protein.

Fusion Proteins

[0180] Fusion proteins are useful for generating antibodies against "BREAST CANCER GENE" polypeptide amino acid sequences and for use in various assay systems. For example, fusion proteins can be used to identify proteins which interact with portions of a "BREAST CANCER GENE" polypeptide. Protein affinity chromatography or library-based assays for protein-protein interactions, such as the yeast two-hybrid or phage display systems, can be used for this purpose. Such methods are well known in the art and also can be used as drug screens.

[0181] A "BREAST CANCER GENE" polypeptide fusion protein comprises two polypeptide segments fused together by means of a peptide bond. The first polypeptide segment comprises at least 25, 50, 75, 100, 150, 200, 300, 400, 500, 600, 700 or 750 contiguous amino acids of an amino acid sequence encoded by any polynucleotide sequences of the SEQ ID NO: 1 to 26 or 53 to 75 or of a biologically active variant, such as those described above. The first polypeptide segment also can comprise full-length "BREAST CANCER GENE".

[0182] The second polypeptide segment can be a full-length protein or a protein fragment. Proteins commonly used in fusion protein construction include β -galactosidase, β -glucuronidase, green fluorescent protein (GFP), autofluorescent proteins, including blue fluorescent protein (BFP), glutathione-S-transferase (GST), luciferase, horseradish peroxidase (HRP), and chloramphenicol acetyltransferase (CAT). Additionally, epitope tags are used in fusion protein constructions, including histidine (His) tags, FLAG tags, influenza hemagglutinin (HA) tags, Myc tags, VSV-G tags, 55

and thioredoxin (Trx) tags. Other fusion constructions can include maltose binding protein (MBP), S-tag, Lex a DNA binding domain (DBD) fusions, GAL4 DNA binding domain fusions, and herpes simplex virus (HSV) BP16 protein fusions. A fusion protein also can be engineered to contain a cleavage site located between the "BREAST CANCER GENE" polypeptide-encoding sequence and the heterologous protein sequence, so that the "BREAST CANCER GENE" polypeptide can be cleaved and purified away from the heterologous moiety.

[0183] A fusion protein can be synthesized chemically, as is known in the art. Preferably, a fusion protein is produced by covalently linking two polypeptide segments or by standard procedures in the art of molecular biology. Recombinant DNA methods can be used to prepare fusion proteins, for example, by making a DNA construct which comprises coding sequences selected from any of the polynucleotide sequences of the SEQ ID NO: 1 to 26 and 53 to 75 in proper reading frame with nucleotides encoding the second polypeptide segment and expressing the DNA construct in a host cell, as is known in the art. Many kits for constructing fusion proteins are available from companies such as Promega Corporation (Madison, WI), Stratagene (La Jolla, CA), CLONTECH (Mountain View, CA), Santa Cruz Biotechnology (Santa Cruz, CA), MBL International Corporation (MIC; Watertown, MA), and Quantum Biotechnologies (Montreal, Canada; 1-888-DNA-KITS).

Identification of Species Homologues

[0184] Species homologues of human a "BREAST CANCER GENE" polypeptide can be obtained using "BREAST CANCER GENE" polypeptide polynucleotides (described below) to make suitable probes or primers for screening cDNA expression libraries from other species, such as mice, monkeys, or yeast, identifying cDNAs which encode homologues of a "BREAST CANCER GENE" polypeptide, and expressing the cDNAs as is known in the art.

Expression of Polynucleotides

[0185] To express a "BREAST CANCER GENE" polynucleotide, the polynucleotide can be inserted into an expression vector which contains the necessary elements for the transcription and translation of the inserted coding sequence. Methods which are well known to those skilled in the art can be used to construct expression vectors containing sequences encoding "BREAST CANCER GENE" polypeptides and appropriate transcriptional and translational control elements. These methods include in vitro recombinant DNA techniques, synthetic techniques, and in vivo genetic recombination. Such techniques are described, for example, in Sambrook et al., (77) and in Ausubel et al., (78).

[0186] A variety of expression vector/host systems can be utilized to contain and express sequences encoding a "BREAST CANCER GENE" polypeptide. These include, but are not limited to, microorganisms, such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors, insect cell systems infected with virus expression vectors (e.g., baculovirus), plant cell systems transformed with virus expression vectors (e.g., cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or with bacterial expression vectors (e.g., Ti or pBR322 plasmids), or animal cell systems.

[0187] The control elements or regulatory sequences are those regions of the vector enhancers, promoters, 5' and 3' untranslated regions which interact with host cellular proteins to carry out transcription and translation. Such elements can vary in their strength and specificity. Depending on the vector system and host utilized, any number of suitable transcription and translation elements, including constitutive and inducible promoters, can be used. For example, when cloning in bacterial systems, inducible promoters such as the hybrid lacZ promoter of the BLUESCRIPT phagemid (Stratagene, LaJolla, Calif.) or pSPORT1 plasmid (Life Technologies) and the like can be used. The baculovirus polyhedrin promoter can be used in insect cells. Promoters or enhancers derived from the genomes of plant cells (e.g., heat shock, RUBISCO, and storage protein genes) or from plant viruses (e.g., viral promoters or leader sequences) can be cloned into the vector. In mammalian cell systems, promoters from mammalian genes or from mammalian viruses are preferable. If it is necessary to generate a cell line that contains multiple copies of a nucleotide sequence encoding a "BREAST CANCER GENE" polypeptide, vectors based on SV40 or EBV can be used with an appropriate selectable marker.

Bacterial and Yeast Expression Systems

[0188] In bacterial systems, a number of expression vectors can be selected depending upon the use intended for the "BREAST CANCER GENE" polypeptide. For example, when a large quantity of the "BREAST CANCER GENE" polypeptide is needed for the induction of antibodies, vectors which direct high level expression of fusion proteins that are readily purified can be used. Such vectors include, but are not limited to, multifunctional *E. coli* cloning and expression vectors such as BLUESCRIPT (Stratagene). In a BLUESCRIPT vector, a sequence encoding the "BREAST CANCER GENE" polypeptide can be ligated into the vector in frame with sequences for the amino terminal Met and the subsequent 7 residues of β -galactosidase so that a hybrid protein is produced. pIN vectors [Van Heeke & Schuster,

(17)] or pGEX vectors (Promega, Madison, Wis.) also can be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST).

[0189] In general, such fusion proteins are soluble and can easily be purified from lysed cells by adsorption to glutathione agarose beads followed by elution in the presence of free glutathione. Proteins made in such systems can be designed to include heparin, thrombin, or factor Xa protease cleavage sites so that the cloned polypeptide of interest can be released from the GST moiety at will.

[0190] In the yeast *Saccharomyces cerevisiae*, a number of vectors containing constitutive or inducible promoters such as alpha factor, alcohol oxidase, and PGH can be used. For reviews, see Ausubel et al., (4) and Grant et al., (18).

Plant and Insect Expression Systems

[0191] If plant expression vectors are used, the expression of sequences encoding "BREAST CANCER GENE" polypeptides can be driven by any of a number of promoters. For example, viral promoters such as the 35S and 19S promoters of CaMV can be used alone or in combination with the omega leader sequence from TMV [Takamatsu, 1987, (96)]. Alternatively, plant promoters such as the small subunit of RUBISCO or heat shock promoters can be used [Coruzzi et al., 1984, (97); Broglie et al., 1984, (98); Winter et al., 1991, (99)]. These constructs can be introduced into plant cells by direct DNA transformation or by pathogen-mediated transfection. Such techniques are described in a number of generally available reviews.

[0192] An insect system also can be used to express a "BREAST CANCER GENE" polypeptide. For example, in one such system *Autographa californica* nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign genes in *Spodoptera frugiperda* cells or in *Trichoplusia* larvae. Sequences encoding "BREAST CANCER GENE" polypeptides can be cloned into a nonessential region of the virus, such as the polyhedrin gene, and placed under control of the polyhedrin promoter. Successful insertion of "BREAST CANCER GENE" polypeptides will render the polyhedrin gene inactive and produce recombinant virus lacking coat protein. The recombinant viruses can then be used to infect *S. frugiperda* cells or *Trichoplusia* larvae in which "BREAST CANCER GENE" polypeptides can be expressed [Engelhard et al., 1994, (100)].

Mammalian Expression Systems

[0193] A number of viral-based expression systems can be used to express "BREAST CANCER GENE" polypeptides in mammalian host cells. For example, if an adenovirus is used as an expression vector, sequences encoding "BREAST CANCER GENE" polypeptides can be ligated into an adenovirus transcription/translation complex comprising the late promoter and tripartite leader sequence. Insertion in a nonessential E1 or E3 region of the viral genome can be used to obtain a viable virus which is capable of expressing a "BREAST CANCER GENE" polypeptide in infected host cells [Logan & Shenk, 1984, (101)]. If desired, transcription enhancers, such as the Rous sarcoma virus (RSV) enhancer, can be used to increase expression in mammalian host cells.

[0194] Human artificial chromosomes (HACs) also can be used to deliver larger fragments of DNA than can be contained and expressed in a plasmid. HACs of 6M to 10M are constructed and delivered to cells via conventional delivery methods (e.g., liposomes, polycationic amino polymers, or vesicles).

[0195] Specific initiation signals also can be used to achieve more efficient translation of sequences encoding "BREAST CANCER GENE" polypeptides. Such signals include the ATG initiation codon and adjacent sequences. In cases where sequences encoding a "BREAST CANCER GENE" polypeptide, its initiation codon, and upstream sequences are inserted into the appropriate expression vector, no additional transcriptional or translational control signals may be needed. However, in cases where only coding sequence, or a fragment thereof, is inserted, exogenous translational control signals (including the ATG initiation codon) should be provided. The initiation codon should be in the correct reading frame to ensure translation of the entire insert. Exogenous translational elements and initiation codons can be of various origins, both natural and synthetic. The efficiency of expression can be enhanced by the inclusion of enhancers which are appropriate for the particular cell system which is used [Scharf et al., 1994, (102)].

Host Cells

[0196] A host cell strain can be chosen for its ability to modulate the expression of the inserted sequences or to process the expressed "BREAST CANCER GENE" polypeptide in the desired fashion. Such modifications of the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation. Posttranslational processing which cleaves a "prepro" form of the polypeptide also can be used to facilitate correct insertion, folding and/or function. Different host cells which have specific cellular machinery and characteristic mechanisms for Post-translational activities (e.g., CHO, HeLa, MDCK, HEK293, and WI38), are available from the American Type Culture Collection (ATCC; 10801 University Boulevard, Manassas, VA 20110-2209) and can be chosen

to ensure the correct modification and processing of the foreign protein.

[0197] Stable expression is preferred for long-term, high-yield production of recombinant proteins. For example, cell lines which stably express "BREAST CANCER GENE" polypeptides can be transformed using expression vectors which can contain viral origins of replication and/or endogenous expression elements and a selectable marker gene on the same or on a separate vector. Following the introduction of the vector, cells can be allowed to grow for 12 days in an enriched medium before they are switched to a selective medium. The purpose of the selectable marker is to confer resistance to selection, and its presence allows growth and recovery of cells which successfully express the introduced "BREAST CANCER GENE" sequences. Resistant clones of stably transformed cells can be proliferated using tissue culture techniques appropriate to the cell type [Freshney et al., 1986, (103).

[0198] Any number of selection systems can be used to recover transformed cell lines. These include, but are not limited to, the herpes simplex virus thymidine kinase [Wigler et al., 1977, (104)] and adenine phosphoribosyltransferase [Lowy et al., 1980, (105)] genes which can be employed in tk- or apt⁻ cells, respectively. Also, antimetabolite, antibiotic, or herbicide resistance can be used as the basis for selection. For example, dhfr confers resistance to methotrexate [Wigler et al., 1980, (106)], npt confers resistance to the aminoglycosides, neomycin and G418 [Colbere-Garapin et al., 1981, (107)], and als and pat confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively. Additional selectable genes have been described. For example, trpB allows cells to utilize indole in place of tryptophan, or hisD, which allows cells to utilize histinol in place of histidine [Hartman & Mulligan, 1988, (108)]. Visible markers such as anthocyanins, β -glucuronidase and its substrate GUS, and luciferase and its substrate luciferin, can be used to identify transformants and to quantify the amount of transient or stable protein expression attributable to a specific vector system [Rhodes et al., 1995, (109)].

Detecting Expression and gene product

[0199] Although the presence of marker gene expression suggests that the "BREAST CANCER GENE" polynucleotide is also present, its presence and expression may need to be confirmed. For example, if a sequence encoding a "BREAST CANCER GENE" polypeptide is inserted within a marker gene sequence, transformed cells containing sequences which encode a "BREAST CANCER GENE" polypeptide can be identified by the absence of marker gene function. Alternatively, a marker gene can be placed in tandem with a sequence encoding a "BREAST CANCER GENE" polypeptide under the control of a single promoter. Expression of the marker gene in response to induction or selection usually indicates expression of the "BREAST CANCER GENE" polynucleotide.

[0200] Alternatively, host cells which contain a "BREAST CANCER GENE" polynucleotide and which express a "BREAST CANCER GENE" polypeptide can be identified by a variety of procedures known to those of skill in the art. These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridization and protein bioassay or immunoassay techniques which include membrane, solution, or chip-based technologies for the detection and/or quantification of polynucleotide or protein. For example, the presence of a polynucleotide sequence encoding a "BREAST CANCER GENE" polypeptide can be detected by DNA-DNA or DNA-RNA hybridization or amplification using probes or fragments or fragments of polynucleotides encoding a "BREAST CANCER GENE" polypeptide. Nucleic acid amplification-based assays involve the use of oligonucleotides selected from sequences encoding a "BREAST CANCER GENE" polypeptide to detect transformants which contain a "BREAST CANCER GENE" polynucleotide.

[0201] A variety of protocols for detecting and measuring the expression of a "BREAST CANCER GENE" polypeptide, using either polyclonal or monoclonal antibodies specific for the polypeptide, are known in the art. Examples include enzyme-linked immunosorbent assay (ELISA), radioimmunoassay (RIA), and fluorescence activated cell sorting (FACS). A two-site, monoclonal-based immunoassay using monoclonal antibodies reactive to two non-interfering epitopes on a "BREAST CANCER GENE" polypeptide can be used, or a competitive binding assay can be employed. These and other assays are described in Hampton et al., (110) and Maddox et al., (111).

[0202] A wide variety of labels and conjugation techniques are known by those skilled in the art and can be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting sequences related to polynucleotides encoding "BREAST CANCER GENE" polypeptides include oligo labeling, nick translation, end-labeling, or PCR amplification using a labeled nucleotide. Alternatively, sequences encoding a "BREAST CANCER GENE" polypeptide can be cloned into a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available, and can be used to synthesize RNA probes in vitro by addition of labeled nucleotides and an appropriate RNA polymerase such as T7, T3, or SP6. These procedures can be conducted using a variety of commercially available kits (Amersham Pharmacia Biotech, Promega, and US Biochemical). Suitable reporter molecules or labels which can be used for ease of detection include radionuclides, enzymes, and fluorescent, chemiluminescent, or chromogenic agents, as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

Expression and Purification of Polypeptides

[0203] Host cells transformed with nucleotide sequences encoding a "BREAST CANCER GENE" polypeptide can be cultured under conditions suitable for the expression and recovery of the protein from cell culture. The polypeptide produced by a transformed cell can be secreted or stored intracellular depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides which encode "BREAST CANCER GENE" polypeptides can be designed to contain signal sequences which direct secretion of soluble "BREAST CANCER GENE" polypeptides through a prokaryotic or eukaryotic cell membrane or which direct the membrane insertion of membrane-bound "BREAST CANCER GENE" polypeptide.

[0204] As discussed above, other constructions can be used to join a sequence encoding a "BREAST CANCER GENE" polypeptide to a nucleotide sequence encoding a polypeptide domain which will facilitate purification of soluble proteins. Such purification facilitating domains include, but are not limited to, metal chelating peptides such as histidine-tryptophan modules that allow purification on immobilized metals, protein A domains that allow purification on immobilized immunoglobulin, and the domain utilized in the FLAGS extension/affinity purification system (Immunex Corp., Seattle, Wash.). Inclusion of cleavable linker sequences such as those specific for Factor Xa or enterokinase (Invitrogen, San Diego, CA) between the purification domain and the "BREAST CANCER GENE" polypeptide also can be used to facilitate purification. One such expression vector provides for expression of a fusion protein containing a "BREAST CANCER GENE" polypeptide and 6 histidine residues preceding a thioredoxin or an enterokinase cleavage site. The histidine residues facilitate purification by IMAC (immobilized metal ion affinity chromatography [Porath et al., 1992, (112)]), while the enterokinase cleavage site provides a means for purifying the "BREAST CANCER GENE" polypeptide from the fusion protein. Vectors which contain fusion proteins are disclosed in Kroll et al., (113).

Chemical Synthesis

[0205] Sequences encoding a "BREAST CANCER GENE" polypeptide can be synthesized, in whole or in part, using chemical methods well known in the art (see Caruthers et al., (114) and Horn et al., (115). Alternatively, a "BREAST CANCER GENE" polypeptide itself can be produced using chemical methods to synthesize its amino acid sequence, such as by direct peptide synthesis using solid-phase techniques [Merrifield, 1963, (116) and Roberge et al., 1995, (117)]. Protein synthesis can be performed using manual techniques or by automation. Automated synthesis can be achieved, for example, using Applied Biosystems 431 A Peptide Synthesizer (Perkin Elmer). Optionally, fragments of "BREAST CANCER GENE" polypeptides can be separately synthesized and combined using chemical methods to produce a full-length molecule.

[0206] The newly synthesized peptide can be substantially purified by preparative high performance liquid chromatography [Creighton, 1983, (118)]. The composition of a synthetic "BREAST CANCER GENE" polypeptide can be confirmed by amino acid analysis or sequencing (e.g., the Edman degradation procedure; see Creighton, (118). Additionally, any portion of the amino acid sequence of the "BREAST CANCER GENE" polypeptide can be altered during direct synthesis and/or combined using chemical methods with sequences from other proteins to produce a variant polypeptide or a fusion protein.

Production of Altered Polypeptides

[0207] As will be understood by those of skill in the art, it may be advantageous to produce "BREAST CANCER GENE" polypeptide-encoding nucleotide sequences possessing non-natural occurring codons. For example, codons preferred by a particular prokaryotic or eukaryotic host can be selected to increase the rate of protein expression or to produce an RNA transcript having desirable properties, such as a half-life which is longer than that of a transcript generated from the naturally occurring sequence.

[0208] The nucleotide sequences disclosed herein can be engineered using methods generally known in the art to alter "BREAST CANCER GENE" polypeptide-encoding sequences for a variety of reasons, including but not limited to, alterations which modify the cloning, processing, and/or expression of the polypeptide or mRNA product. DNA shuffling by random fragmentation and PCR re-assembly of gene fragments and synthetic oligonucleotides can be used to engineer the nucleotide sequences. For example, site-directed mutagenesis can be used to insert new restriction sites, alter glycosylation patterns, change codon preference, produce splice variants, introduce mutations, and so forth.

Predictive, Diagnostic and Prognostic Assays

[0209] The present invention provides method for determining whether a subject is at risk for developing malignant neoplasia and breast cancer in particular by detecting one of the disclosed polynucleotide markers comprising any of

the polynucleotides sequences of the SEQ ID NO: 2 to 6, 8, 9, 11 to 16, 18, 19 or 21 to 26 or 53 to 75 and/or the polypeptide markers encoded thereby or polypeptide markers comprising any of the polypeptide sequences of the SEQ ID NO: 28 to 32, 34, 35, 37 to 42, 44, 45 or 47 to 52 or 76 to 98 or at least 2 of the disclosed polynucleotides selected from SEQ ID NO: 1 to 26 and 53 to 75 or the at least 2 of the disclosed polypeptides selected from SEQ ID

NO: 28 to 32 and 76 to 98 for malignant neoplasia and breast cancer in particular.
[0210] In clinical applications, biological samples can be screened for the presence and/or absence of the biomarkers identified herein. Such samples are for example needle biopsy cores, surgical resection samples, or body fluids like serum, thin needle nipple aspirates and urine. For example, these methods include obtaining a biopsy, which is optionally fractionated by cryostat sectioning to enrich diseased cells to about 80% of the total cell population. In certain

embodiments, polynucleotides extracted from these samples may be amplified using techniques well known in the art. The expression levels of selected markers detected would be compared with statistically valid groups of diseased and healthy samples.
[0211] In one embodiment the diagnostic method comprises determining whether a subject has an abnormal mRNA and/or protein level of the disclosed markers, such as by Northern blot analysis, reverse transcription-polymerase chain reaction (RT-PCR), in situ hybridization, immunoprecipitation, Western blot hybridization, or immunohistochemistry.

According to the method, cells are obtained from a subject and the levels of the disclosed biomarkers, protein or mRNA level, is determined and compared to the level of these markers in a healthy subject. An abnormal level of the biomarker polypeptide or mRNA levels is likely to be indicative of malignant neoplasia such as breast cancer.

[0212] In another embodiment the diagnostic method comprises determining whether a subject has an abnormal DNA content of said genes or said genomic loci, such as by Southern blot analysis, dot blot analysis, fluorescence or colorimetric In Situ hybridization, comparative genomic hybridization, genotyping by VNTR, STS-PCR or quantitative PCR. In general these assays comprise the usage of probes from representative genomic regions. The probes contain at least parts of said genomic regions or sequences complementary or analogous to said regions. In particular intra- or intergenic regions of said genes or genomic regions. The probes can consist of nucleotide sequences or sequences of analogous functions (e.g. PNAs, Morpholino oligomers) being able to bind to target regions by hybridization. In general genomic regions being altered in said patient samples are compared with unaffected control samples (normal tissue from the same or different patients, surrounding unaffected tissue, peripheral blood) or with genomic regions of the same sample that don't have said alterations and can therefore serve as internal controls. In a preferred embodiment regions located on the same chromosome are used. Alternatively, gonosomal regions and /or regions with defined varying amount in the sample are used. In one favored embodiment the DNA content, structure, composition or modification is compared that lie within distinct genomic regions. Especially favored are methods that detect the DNA content of said samples, where the amount of target regions are altered by amplification and or deletions. In another embodiment the target regions are analyzed for the presence of polymorphisms (e.g. Single Nucleotide Polymorphisms or mutations) that affect or predispose the cells in said samples with regard to clinical aspects, being of diagnostic, prognostic or therapeutic value. Preferably, the identification of sequence variations is used to define haplotypes that result in characteristic behavior of said samples with said clinical aspects.

[0213] The following examples of genes in 17q12-21.2 are offered by way of illustration, not by way of limitation.

[0214] One embodiment of the invention is a method for the prediction, diagnosis or prognosis of malignant neoplasia by the detection of at least 10, at least 5, or at least 4, or at least 3 and more preferably at least 2 markers whereby the markers are genes and fragments thereof and/or genomic nucleic acid sequences that are located on one chromosomal region which is altered in malignant neoplasia.

[0215] One further embodiment of the invention is method for the prediction, diagnosis or prognosis of malignant neoplasia by the detection of at least 10, at least 5, or at least 4, or at least 3 and more preferably at least 2 markers whereby the markers (a) are genes and fragments thereof and/or genomic nucleic acid sequences that are located on one or more chromosomal region(s) which is/are altered in malignant neoplasia and (b) functionally interact as (i) receptor and ligand or (ii) members of the same signal transduction pathway or (iii) members of synergistic signal transduction pathways or (iv) members of antagonistic signal transduction pathways or (v) transcription factor and transcription factor binding site.

[0216] In one embodiment, the method for the prediction, diagnosis or prognosis of malignant neoplasia and breast cancer in particular is done by the detection of:

(a) polynucleotide selected from the polynucleotides of the SEQ ID NO: 2 to 6, 8, 9, 11 to 16, 18, 19, 21 to 26 or 53 to 75;

(b) a polynucleotide which hybridizes under stringent conditions to a polynucleotide specified in (a) encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 2 or 3;

(c) a polynucleotide the sequence of which deviates from the polynucleotide specified in (a) and (b) due to the

generation of the genetic code encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 2 or 3;

(d) a polynucleotide which represents a specific fragment, derivative or allelic variation of a polynucleotide sequence specified in (a) to (c);

in a biological sample comprising the following steps: hybridizing any polynucleotide or analogous oligomer specified in (a) to (d) to a polynucleotide material of a biological sample, thereby forming a hybridization complex; and detecting said hybridization complex.

[0217] In another embodiment the method for the prediction, diagnosis or prognosis of malignant neoplasia is done as just described but, wherein before hybridization, the polynucleotide material of the biological sample is amplified.

[0218] In another embodiment the method for the diagnosis or prognosis of malignant neoplasia and breast cancer in particular is done by the detection of:

(a) a polynucleotide selected from the polynucleotides of the SEQ ID NO: 2 to 6, 8, 9, 11 to 16, 18, 19, 21 to 26 or 53 to 75;

(b) a polynucleotide which hybridizes under stringent conditions to a polynucleotide specified in (a) encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 2 or 3;

(c) a polynucleotide the sequence of which deviates from the polynucleotide specified in (a) and (b) due to the generation of the genetic code encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 2 or 3;

(d) a polynucleotide which represents a specific fragment, derivative or allelic variation of a polynucleotide sequence specified in (a) to (c);

(e) a polypeptide encoded by a polynucleotide sequence specified in (a) to (d)

(f) a polypeptide comprising any polypeptide of SEQ ID NO: 28 to 32, 34, 35, 37 to 42, 44, 45, 47 to 52 or 76 to 98;

comprising the steps of contacting a biological sample with a reagent which specifically interacts with the polynucleotide specified in (a) to (d) or the polypeptide specified in (e).

DNA array technology

[0219] In one embodiment, the present Invention also provides a method wherein polynucleotide probes are immobilized on a DNA chip in an organized array. Oligonucleotides can be bound to a solid Support by a variety of processes, including lithography. For example a chip can hold up to 4100,00 oligonucleotides (GeneChip, Affymetrix). The present invention provides significant advantages over the available tests for malignant neoplasia, such as breast cancer, because it increases the reliability of the test by providing an array of polynucleotide markers on a single chip.

[0220] The method includes obtaining a biopsy of an affected person, which is optionally fractionated by cryostat sectioning to enrich diseased cells to about 80% of the total cell population and the use of body fluids such as serum or urine, serum or cell containing liquids (e.g. derived from fine needle aspirates). The DNA or RNA is then extracted, amplified, and analyzed with a DNA chip to determine the presence or absence of the marker polynucleotide sequences. In one embodiment, the polynucleotide probes are spotted onto a substrate in a two-dimensional matrix or array. Samples of polynucleotides can be labeled and then hybridized to the probes. Double-stranded polynucleotides, comprising the labeled sample polynucleotides bound to probe polynucleotides, can be detected once the unbound portion of the sample is washed away.

[0221] The probe polynucleotides can be spotted on substrates including glass, nitrocellulose, etc. The probes can be bound to the Substrate by either covalent bonds or by non-specific interactions, such as hydrophobic interactions. The sample polynucleotides can be labeled using radioactive labels, fluorophores, chromophores, etc. Techniques for constructing arrays and methods of using these arrays are described in EP 0 799 897; WO 97/29212; WO 97/27317; EP 0 785 280; WO 97/02357; U.S. Pat. No. 5,593,839; U.S. Pat. No. 5,578,832; EP 0 728 520; U.S. Pat. No. 5,599,695; EP 0 721 016; U.S. Pat. No. 5,556,752; WO 95/22058; and U.S. Pat. No. 5,631,734.

[0222] Further, arrays can be used to examine differential expression of genes and can be used to determine gene function. For example, arrays of the instant polynucleotide sequences can be used to determine if any of the polynucleotide sequences are differentially expressed between normal cells and diseased cells, for example. High expression

of a particular message in a diseased sample, which is not observed in a corresponding normal sample, can indicate a breast cancer specific protein.

[0223] Accordingly, in one aspect, the invention provides probes and primers that are specific to the unique polynucleotide markers disclosed herein.

[0224] In one embodiment, the method comprises using a polynucleotide probe to determine the presence of malignant or breast cancer cells in particular in a tissue from a patient. Specifically, the method comprises:

1) providing a polynucleotide probe comprising a nucleotide sequence at least 12 nucleotides in length, preferably at least 15 nucleotides, more preferably, 25 nucleotides, and most preferably at least 40 nucleotides, and up to all or nearly all of the coding sequence which is complementary to a portion of the coding sequence of a polynucleotide selected from the polynucleotides of SEQ ID NO: 1 to 26 and 53 to 75 or a sequence complementary thereto and is

2) differentially expressed in malignant neoplasia, such as breast cancer;

3) obtaining a tissue sample from a patient with malignant neoplasia;

4) providing a second tissue sample from a patient with no malignant neoplasia;

5) contacting the polynucleotide probe under stringent conditions with RNA of each of said first and second tissue samples (e.g., in a Northern blot or in situ hybridization assay); and

6) comparing (a) the amount of hybridization of the probe with RNA of the first tissue sample, with (b) the amount of hybridization of the probe with RNA of the second tissue sample;

wherein a statistically significant difference in the amount of hybridization with the RNA of the first tissue sample as compared to the amount of hybridization with the RNA of the second tissue sample is indicative of malignant neoplasia and breast cancer in particular in the first tissue sample.

Data analysis methods

[0225] Comparison of the expression levels of one or more "BREAST CANCER GENES" with reference expression levels, e.g., expression levels in diseased cells of breast cancer or in normal counterpart cells, is preferably conducted using computer systems. In one embodiment, expression levels are obtained in two cells and these two sets of expression levels are introduced into a computer system for comparison. In a preferred embodiment, one set of expression levels is entered into a computer system for comparison with values that are already present in the computer system, or in computer-readable form that is then entered into the computer system.

[0226] In one embodiment, the invention provides a computer readable form of the gene expression profile data of the invention, or of values corresponding to the level of expression of at least one "BREAST CANCER GENE" in a diseased cell. The values can be mRNA expression levels obtained from experiments, e.g., microarray analysis. The values can also be mRNA levels normalised relative to a reference gene whose expression is constant in numerous cells under numerous conditions, e.g., GAPDH. In other embodiments, the values in the computer are ratios of, or differences between, normalized or non-normalized mRNA levels in different samples.

[0227] The gene expression profile data can be in the form of a table, such as an Excel table. The data can be alone, or it can be part of a larger database, e.g., comprising other expression profiles. For example, the expression profile data of the invention can be part of a public database. The computer readable form can be in a computer. In another embodiment, the invention provides a computer displaying the gene expression profile data.

[0228] In one embodiment, the invention provides a method for determining the similarity between the level of expression of one or more "BREAST CANCER GENES" in a first cell, e.g., a cell of a subject, and that in a second cell, comprising obtaining the level of expression of one or more "BREAST CANCER GENES" in a first cell and entering these values into a computer comprising a database including records comprising values corresponding to levels of expression of one or more "BREAST CANCER GENES" in a second cell, and processor instructions, e.g., a user interface, capable of receiving a selection of one or more values for comparison purposes with data that is stored in the computer. The computer may further comprise a means for converting the comparison data into a diagram or chart or other type of output.

[0229] In another embodiment, values representing expression levels of "BREAST CANCER GENES" are entered into a computer system, comprising one or more databases with reference expression levels obtained from more than one cell. For example, the computer comprises expression data of diseased and normal cells. Instructions are provided to the computer, and the computer is capable of comparing the data entered with the data in the computer to determine

whether the data entered is more similar to that of a normal cell or of a diseased cell.

[0230] In another embodiment, the computer comprises values of expression levels in cells of subjects at different stages of breast cancer, and the computer is capable of comparing expression data entered into the computer with the data stored, and produce results indicating to which of the expression profiles in the computer, the one entered is most similar, such as to determine the stage of breast cancer in the subject.

[0231] In yet another embodiment, the reference expression profiles in the computer are expression profiles from cells of breast cancer of one or more subjects, which cells are treated *in vivo* or *in vitro* with a drug used for therapy of breast cancer. Upon entering of expression data of a cell of a subject treated *in vitro* or *in vivo* with the drug, the computer is instructed to compare the data entered to the data in the computer, and to provide results indicating whether the expression data input into the computer are more similar to those of a cell of a subject that is responsive to the drug or more similar to those of a cell of a subject that is not responsive to the drug. Thus, the results indicate whether the subject is likely to respond to the treatment with the drug or unlikely to respond to it.

[0232] In one embodiment, the invention provides a system that comprises a means for receiving gene expression data for one or a plurality of genes; a means for comparing the gene expression data from each of said one or plurality of genes to a common reference frame; and a means for presenting the results of the comparison. This system may further comprise a means for clustering the data.

[0233] In another embodiment, the invention provides a computer program for analyzing gene expression data comprising (i) a computer code that receives as input gene expression data for a plurality of genes and (ii) a computer code that compares said gene expression data from each of said plurality of genes to a common reference frame.

[0234] The invention also provides a machine-readable or computer-readable medium including program instructions for performing the following steps: (i) comparing a plurality of values corresponding to expression levels of one or more genes characteristic of breast cancer in a query cell with a database including records comprising reference expression or expression profile data of one or more reference cells and an annotation of the type of cell; and (ii) indicating to which cell the query cell is most similar based on similarities of expression profiles. The reference cells can be cells from subjects at different stages of breast cancer. The reference cells can also be cells from subjects responding or not responding to a particular drug treatment and optionally incubated *in vitro* or *in vivo* with the drug.

[0235] The reference cells may also be cells from subjects responding or not responding to several different treatments, and the computer system indicates a preferred treatment for the subject. Accordingly, the invention provides a method for selecting a therapy for a patient having breast cancer, the method comprising: (i) providing the level of expression of one or more genes characteristic of breast cancer in a diseased cell of the patient; (ii) providing a plurality of reference profiles, each associated with a therapy, wherein the subject expression profile and each reference profile has a plurality of values, each value representing the level of expression of a gene characteristic of breast cancer; and (iii) selecting the reference profile most similar to the subject expression profile, to thereby select a therapy for said patient. In a preferred embodiment step (iii) is performed by a computer. The most similar reference profile may be selected by weighing a comparison value of the plurality using a weight value associated with the corresponding expression data.

[0236] The relative abundance of an mRNA in two biological samples can be scored as a perturbation and its magnitude determined (i.e., the abundance is different in the two sources of mRNA tested), or as not perturbed (i.e., the relative abundance is the same). In various embodiments, a difference between the two sources of RNA of at least a factor of about 25% (RNA from one source is 25% more abundant in one source than the other source), more usually about 50%, even more often by a factor of about 2 (twice as abundant), 3 (three times as abundant) or 5 (five times as abundant) is scored as a perturbation. Perturbations can be used by a computer for calculating and expression comparisons.

[0237] Preferably, in addition to identifying a perturbation as positive or negative, it is advantageous to determine the magnitude of the perturbation. This can be carried out, as noted above, by calculating the ratio of the emission of the two fluorophores used for differential labeling, or by analogous methods that will be readily apparent to those of skill in the art.

[0238] The computer readable medium may further comprise a pointer to a descriptor of a stage of breast cancer or to a treatment for breast cancer.

[0239] In operation, the means for receiving gene expression data, the means for comparing the gene expression data, the means for presenting, the means for normalizing, and the means for clustering within the context of the systems of the present invention can involve a programmed computer with the respective functionalities described herein, implemented in hardware or hardware and software; a logic circuit or other component of a programmed computer that performs the operations specifically identified herein, dictated by a computer program; or a computer memory encoded with executable instructions representing a computer program that can cause a computer to function in the particular fashion described herein.

[0240] Those skilled in the art will understand that the systems and methods of the present invention may be applied to a variety of systems, including IBM-compatible personal computers running MS-DOS or Microsoft Windows.

[0241] The computer may have internal components linked to external components. The internal components may include a processor element interconnected with a main memory. The computer system can be an Intel Pentium®-based processor of 200 MHz or greater clock rate and with 32 MB or more of main memory. The external component may comprise a mass storage, which can be one or more hard disks (which are typically packaged together with the processor and memory). Such hard disks are typically of 1 GB or greater storage capacity. Other external components include a user interface device, which can be a monitor, together with an inputting device, which can be a "mouse", or other graphic input devices, and/or a keyboard. A printing device can also be attached to the computer.

[0242] Typically, the computer system is also linked to a network link, which can be part of an Ethernet link to other local computer systems, remote computer systems, or wide area communication networks, such as the Internet. This network link allows the computer system to share data and processing tasks with other computer systems.

[0243] Loaded into memory during operation of this system are several software components, which are both standard in the art and special to the instant invention. These software components collectively cause the computer system to function according to the methods of this invention. These software components are typically stored on a mass storage. A software component represents the operating system, which is responsible for managing the computer system and its network interconnections. This operating system can be, for example, of the Microsoft Windows' family, such as Windows 95, Windows 98, or Windows NT. A software component represents common languages and functions conveniently present on this system to assist programs implementing the methods specific to this invention. Many high or low level computer languages can be used to program the analytic methods of this invention. Instructions can be interpreted during run-time or compiled. Preferred languages include C/C++, and JAVA®. Most preferably, the methods of this invention are programmed in mathematical software packages which allow symbolic entry of equations and high-level specification of processing, including algorithms to be used, thereby freeing a user of the need to procedurally program individual equations or algorithms. Such packages include Matlab from Mathworks (Natick, Mass.), Mathematica from Wolfram Research (Champaign, Ill.), or S-Plus from Math Soft (Cambridge, Mass.). Accordingly, a software component represents the analytic methods of this invention as programmed in a procedural language or symbolic package. In a preferred embodiment, the computer system also contains a database comprising values representing levels of expression of one or more genes characteristic of breast cancer. The database may contain one or more expression profiles of genes characteristic of breast cancer in different cells.

[0244] In an exemplary implementation, to practice the methods of the present invention, a user first loads expression profile data into the computer system. These data can be directly entered by the user from a monitor and keyboard, or from other computer systems linked by a network connection, or on removable storage media such as a CD-ROM or floppy disk or through the network. Next the user causes execution of expression profile analysis software which performs the steps of comparing and, e.g., clustering co-varying genes into groups of genes.

[0245] In another exemplary implementation, expression profiles are compared using a method described in U.S. Patent No. 6,203,987. A user first loads expression profile data into the computer system. Geneset profile definitions are loaded into the memory from the storage media or from a remote computer, preferably from a dynamic geneset database system, through the network. Next the user causes execution of projection software which performs the steps of converting expression profile to projected expression profiles. The projected expression profiles are then displayed.

[0246] In yet another exemplary implementation, a user first loads a projected profile into the memory. The user then causes the loading of a reference profile into the memory. Next, the user causes the execution of comparison software which performs the steps of objectively comparing the profiles.

Detection of variant polynucleotide sequence

[0247] In yet another embodiment, the invention provides methods for determining whether a subject is at risk for developing a disease, such as a predisposition to develop malignant neoplasia, for example breast cancer, associated with an aberrant activity of any one of the polypeptides encoded by any of the polynucleotides of the SEQ ID NO: 1 to 26 or 53 to 75, wherein the aberrant activity of the polypeptide is characterized by detecting the presence or absence of a genetic lesion characterized by at least one of these:

- (i) an alteration affecting the integrity of a gene encoding a marker polypeptides, or
- (ii) the misexpression of the encoding polynucleotide.

[0248] To illustrate, such genetic lesions can be detected by ascertaining the existence of at least one of these:

- I. a deletion of one or more nucleotides from the polynucleotide sequence
- II. an addition of one or more nucleotides to the polynucleotide sequence

III. a substitution of one or more nucleotides of the polynucleotide sequence

IV. a gross chromosomal rearrangement of the polynucleotide sequence

V. a gross alteration in the level of a messenger RNA transcript of the polynucleotide sequence

VI. aberrant modification of the polynucleotide sequence, such as of the methylation pattern of the genomic DNA

VII. the presence of a non-wild type splicing pattern of a messenger RNA transcript of the gene

VIII. a non-wild type level of the marker polypeptide

IX. allelic loss of the gene

X. allelic gain of the gene

XI. inappropriate post-translational modification of the marker polypeptide

[0249] The present Invention provides assay techniques for detecting mutations in the encoding polynucleotide sequence. These methods include, but are not limited to, methods involving sequence analysis, Southern blot hybridization, restriction enzyme site mapping, and methods involving detection of absence of nucleotide pairing between the polynucleotide to be analyzed and a probe.

[0250] Specific diseases or disorders, e.g., genetic diseases or disorders, are associated with specific allelic variants of polymorphic regions of certain genes, which do not necessarily encode a mutated protein. Thus, the presence of a specific allelic variant of a polymorphic region of a gene in a subject can render the subject susceptible to developing a specific disease or disorder. Polymorphic regions in genes, can be identified, by determining the nucleotide sequence of genes in populations of individuals. If a polymorphic region is identified, then the link with a specific disease can be determined by studying specific populations of individuals, e.g. individuals which developed a specific disease, such as breast cancer. A polymorphic region can be located in any region of a gene, e.g., exons, in coding or non coding regions of exons, introns, and promoter region.

[0251] In an exemplary embodiment, there is provided a polynucleotide composition comprising a polynucleotide probe including a region of nucleotide sequence which is capable of hybridising to a sense or antisense sequence of a gene or naturally occurring mutants thereof, or 5' or 3' flanking sequences or intronic sequences naturally associated with the subject genes or naturally occurring mutants thereof. The polynucleotide of a cell is rendered accessible for hybridization, the probe is contacted with the polynucleotide of the sample, and the hybridization of the probe to the sample polynucleotide is detected. Such techniques can be used to detect lesions or allelic variants at either the genomic or mRNA level, including deletions, substitutions, etc., as well as to determine mRNA transcript levels.

[0252] A preferred detection method is allele specific hybridization using probes overlapping the mutation or polymorphic site and having about 5, 10, 20, 25, or 30 nucleotides around the mutation or polymorphic region. In a preferred embodiment of the invention, several probes capable of hybridising specifically to allelic variants are attached to a solid phase support, e.g., a "chip". Mutation detection analysis using these chips comprising oligonucleotides, also termed "DNA probe arrays" is described e.g., in Cronin et al. (119). In one embodiment, a chip comprises all the allelic variants of at least one polymorphic region of a gene. The solid phase support is then contacted with a test polynucleotide and hybridization to the specific probes is detected. Accordingly, the identity of numerous allelic variants of one or more genes can be identified in a simple hybridization experiment.

[0253] In certain embodiments, detection of the lesion comprises utilizing the probe/primer in a polymerase chain reaction (PCR) (see, e.g. U.S. Patent Nos. 4,683,195 and 4,683,202), such as anchor PCR or RACE PCR, or, alternatively, in a ligase chain reaction (LCR) [Landegran et al., 1988, (120) and Nakazawa et al., 1994 (121)], the latter of which can be particularly useful for detecting point mutations in the gene; Abravaya et al., 1995, (122)]. In a merely illustrative embodiment, the method includes the steps of (i) collecting a sample of cells from a patient, (ii) isolating polynucleotide (e.g., genomic, mRNA or both) from the cells of the sample, (iii) contacting the polynucleotide sample with one or more primers which specifically hybridize to a polynucleotide sequence under conditions such that hybridization and amplification of the polynucleotide (if present) occurs, and (iv) detecting the presence or absence of an amplification product, or detecting the size of the amplification product and comparing the length to a control sample. It is anticipated that PCR and/or LCR may be desirable to use as a preliminary amplification step in conjunction with any of the techniques used for detecting mutations described herein.

[0254] Alternative amplification methods include: self sustained sequence replication [Guatelli, J.C. et al., 1990, (123)], transcriptional amplification system [Kwoh, D.Y. et al., 1989, (124)], Q-Beta replicase [Lizardi, P.M. et al., 1988,

(125)], or any other polynucleotide amplification method, followed by the detection of the amplified molecules using techniques well known to those of skill in the art. These detection schemes are especially useful for the detection of polynucleotide molecules if such molecules are present in very low numbers.

[0255] In a preferred embodiment of the subject assay, mutations in, or allelic variants, of a gene from a sample cell are identified by alterations in restriction enzyme cleavage patterns. For example, sample and control DNA is isolated, amplified (optionally), digested with one or more restriction endonucleases, and fragment length sizes are determined by gel electrophoresis. Moreover, the use of sequence specific ribozymes (see, for example, U.S. Patent No. 5,498,531) can be used to score for the presence of specific mutations by development or loss of a ribozyme cleavage site.

In situ hybridization

[0256] In one aspect, the method comprises *in situ* hybridization with a probe derived from a given marker polynucleotide, which sequence is selected from any of the polynucleotide sequences of the SEQ ID NO: 1 to 9, or 11 to 19 or 21 to 26 and 53 to 75 or a sequence complementary thereto. The method comprises contacting the labeled hybridization probe with a sample of a given type of tissue from a patient potentially having malignant neoplasia and breast cancer in particular as well as normal tissue from a person with no malignant neoplasia, and determining whether the probe labels tissue of the patient to a degree significantly different (e.g., by at least a factor of two, or at least a factor of five, or at least a factor of twenty, or at least a factor of fifty) than the degree to which normal tissue is labelled.

Polypeptide detection

[0257] The subject invention further provides a method of determining whether a cell sample obtained from a subject possesses an abnormal amount of marker polypeptide which comprises (a) obtaining a cell sample from the subject, (b) quantitatively determining the amount of the marker polypeptide in the sample so obtained, and (c) comparing the amount of the marker polypeptide so determined with a known standard, so as to thereby determine whether the cell sample obtained from the subject possesses an abnormal amount of the marker polypeptide. Such marker polypeptides may be detected by immunohistochemical assays, dot-blot assays, ELISA and the like.

Antibodies

[0258] Any type of antibody known in the art can be generated to bind specifically to an epitope of a "BREAST CANCER GENE" polypeptide. An antibody as used herein includes intact immunoglobulin molecules, as well as fragments thereof, such as Fab, F(ab)₂, and Fv, which are capable of binding an epitope of a "BREAST CANCER GENE" polypeptide. Typically, at least 6, 8, 10, or 12 contiguous amino acids are required to form an epitope. However, epitopes which involve non-contiguous amino acids may require more, e.g., at least 15, 25, or 50 amino acids.

[0259] An antibody which specifically binds to an epitope of a "BREAST CANCER GENE" polypeptide can be used therapeutically, as well as in immunochemical assays, such as Western blots, ELISAs, radioimmunoassays, immunohistochemical assays, immunoprecipitations, or other immunochemical assays known in the art. Various immunoassays can be used to identify antibodies having the desired specificity. Numerous protocols for competitive binding or immunoradiometric assays are well known in the art. Such immunoassays typically involve the measurement of complex formation between an immunogen and an antibody which specifically binds to the immunogen.

[0260] Typically, an antibody which specifically binds to a "BREAST CANCER GENE" polypeptide provides a detection signal at least 5-, 10-, or 20-fold higher than a detection signal provided with other proteins when used in an immunochemical assay. Preferably, antibodies which specifically bind to "BREAST CANCER GENE" polypeptides do not detect other proteins in immunochemical assays and can immunoprecipitate a "BREAST CANCER GENE" polypeptide from solution.

[0261] "BREAST CANCER GENE" polypeptides can be used to immunize a mammal, such as a mouse, rat, rabbit, guinea pig, monkey, or human, to produce polyclonal antibodies. If desired, a "BREAST CANCER GENE" polypeptide can be conjugated to a carrier protein, such as bovine serum albumin, thyroglobulin, and keyhole limpet hemocyanin. Depending on the host species, various adjuvants can be used to increase the immunological response. Such adjuvants include, but are not limited to, Freund's adjuvant, mineral gels (e.g., aluminum hydroxide), and surface active substances (e.g. lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, keyhole limpet hemocyanin, and dinutrophenol). Among adjuvants used in humans, BCG (bacilli Calmette-Guerin) and *Corynebacterium parvum* are especially useful.

[0262] Monoclonal antibodies which specifically bind to a "BREAST CANCER GENE" polypeptide can be prepared using any technique which provides for the production of antibody molecules by continuous cell lines in culture. These techniques include, but are not limited to, the hybridoma technique, the human B cell hybridoma technique, and the EBV hybridoma technique [Kohler et al., 1985, (136); Kozbor et al., 1985, (137); Cote et al., 1983, (138) and Cole et

al., 1984, (139)].

[0263] In addition, techniques developed for the production of chimeric antibodies, the splicing of mouse antibody genes to human antibody genes to obtain a molecule with appropriate antigen specificity and biological activity, can be used [Morrison et al., 1984, (140); Neuberger et al., 1984, (141); Takeda et al., 1985, (142)]. Monoclonal and other antibodies also can be humanized to prevent a patient from mounting an immune response against the antibody when it is used therapeutically. Such antibodies may be sufficiently similar in sequence to human antibodies to be used directly in therapy or may require alteration of a few key residues. Sequence differences between rodent antibodies and human sequences can be minimized by replacing residues which differ from those in the human sequences by site directed mutagenesis of individual residues or by grating of entire complementarity determining regions. Alternatively, humanized antibodies can be produced using recombinant methods, as described in GB2188638B. Antibodies which specifically bind to a "BREAST CANCER GENE" polypeptide can contain antigen binding sites which are either partially or fully humanized, as disclosed in U.S. Patent 5,565,332.

[0264] Alternatively, techniques described for the production of single chain antibodies can be adapted using methods known in the art to produce single chain antibodies which specifically bind to "BREAST CANCER GENE" polypeptides. Antibodies with related specificity, but of distinct idiotypic composition, can be generated by chain shuffling from random combinatorial immunoglobulin libraries [Burton, 1991, (143)].

[0265] Single-chain antibodies also can be constructed using a DNA amplification method, such as PCR, using hybridoma cDNA as a template [Thirion et al., 1996, (144)]. Single-chain antibodies can be mono- or bispecific, and can be bivalent or tetravalent. Construction of tetravalent, bispecific single-chain antibodies is taught, for example, in Coloma & Morrison, (145). Construction of bivalent, bispecific single-chain antibodies is taught in Mallender & Voss, (146).

[0266] A nucleotide sequence encoding a single-chain antibody can be constructed using manual or automated nucleotide synthesis, cloned into an expression construct using standard recombinant DNA methods, and introduced into a cell to express the coding sequence, as described below. Alternatively, single-chain antibodies can be produced directly using, for example, filamentous phage technology [Verhaar et al., 1995, (147); Nicholls et al., 1993, (148)].

[0267] Antibodies which specifically bind to "BREAST CANCER GENE" polypeptides also can be produced by inducing in vivo production in the lymphocyte population or by screening immunoglobulin libraries or panels of highly specific binding reagents as disclosed in the literature [Orlandi et al., 1989, (149) and Winter et al., 1991, (150)].

[0268] Other types of antibodies can be constructed and used therapeutically in methods of the invention. For example, chimeric antibodies can be constructed as disclosed in WO 93/03151. Binding proteins which are derived from immunoglobulins and which are multivalent and multispecific, such as the antibodies described in WO 94/13804, also can be prepared.

[0269] Antibodies according to the invention can be purified by methods well known in the art. For example, antibodies can be affinity purified by passage over a column to which a "BREAST CANCER GENE" polypeptide is bound. The bound antibodies can then be eluted from the column using a buffer with a high salt concentration.

[0270] Immunoassays are commonly used to quantify the levels of proteins in cell samples, and many other immunoassay techniques are known in the art. The invention is not limited to a particular assay procedure, and therefore is intended to include both homogeneous and heterogeneous procedures. Exemplary immunoassays which can be conducted according to the invention include fluorescence polarisation immunoassay (FPIA), fluorescence immunoassay (FIA), enzyme immunoassay (EIA), nephelometric inhibition immunoassay (NIA), enzyme linked immunosorbent assay (ELISA), and radioimmunoassay (RIA). An indicator moiety, or label group, can be attached to the subject antibodies and is selected so as to meet the needs of various uses of the method which are often dictated by the availability of assay equipment and compatible immunoassay procedures. General techniques to be used in performing the various immunoassays noted above are known to those of ordinary skill in the art.

[0271] In another embodiment, the level of at least one product encoded by any of the polynucleotide sequences of the SEQ ID NO: 2 to 6, 8, 9, 11 to 16, 18, 19 or 21 to 26 or 53 to 75 or of at least 2 products encoded by a polynucleotide selected from SEQ ID NO: 1 to 26 and 53 to 75 or a sequence complementary thereto, in a biological fluid (e.g., blood or urine) of a patient may be determined as a way of monitoring the level of expression of the marker polynucleotide sequence in cells of that patient. Such a method would include the steps of obtaining a sample of a biological fluid from the patient, contacting the sample (or proteins from the sample) with an antibody specific for a encoded marker polypeptide, and determining the amount of immune complex formation by the antibody, with the amount of immune complex formation being indicative of the level of the marker encoded product in the sample. This determination is particularly instructive when compared to the amount of immune complex formation by the same antibody in a control sample taken from a normal individual or in one or more samples previously or subsequently obtained from the same person.

[0272] In another embodiment, the method can be used to determine the amount of marker polypeptide present in a cell, which in turn can be correlated with progression of the disorder, e.g., plaque formation. The level of the marker polypeptide can be used predictively to evaluate whether a sample of cells contains cells which are, or are predisposed

towards becoming, plaque associated cells. The observation of marker polypeptide level can be utilized in decisions regarding, e.g., the use of more stringent therapies.

[0273] As set out above, one aspect of the present invention relates to diagnostic assays for determining, in the context of cells isolated from a patient, if the level of a marker polypeptide is significantly reduced in the sample cells. The term "significantly reduced" refers to a cell phenotype wherein the cell possesses a reduced cellular amount of the marker polypeptide relative to a normal cell of similar tissue origin. For example, a cell may have less than about 50%, 25%, 10%, or 5% of the marker polypeptide that a normal control cell. In particular, the assay evaluates the level of marker polypeptide in the test cells, and, preferably, compares the measured level with marker polypeptide detected in at least one control cell, e.g., a normal cell and/or a transformed cell of known phenotype.

[0274] Of particular importance to the subject invention is the ability to quantify the level of marker polypeptide as determined by the number of cells associated with a normal or abnormal marker polypeptide level. The number of cells with a particular marker polypeptide phenotype may then be correlated with patient prognosis. In one embodiment of the invention, the marker polypeptide phenotype of the lesion is determined as a percentage of cells in a biopsy which are found to have abnormally high/low levels of the marker polypeptide. Such expression may be detected by immunohistochemical assays, dot-blot assays, ELISA and the like.

Immunohistochemistry

[0275] Where tissue samples are employed, immunohistochemical staining may be used to determine the number of cells having the marker polypeptide phenotype. For such staining, a multiblock of tissue is taken from the biopsy or other tissue sample and subjected to proteolytic hydrolysis, employing such agents as protease K or pepsin. In certain embodiments, it may be desirable to isolate a nuclear fraction from the sample cells and detect the level of the marker polypeptide in the nuclear fraction.

[0276] The tissues samples are fixed by treatment with a reagent such as formalin, glutaraldehyde, methanol, or the like. The samples are then incubated with an antibody, preferably a monoclonal antibody, with binding specificity for the marker polypeptides. This antibody may be conjugated to a Label for subsequent detection of binding. samples are incubated for a time Sufficient for formation of the immuno-complexes. Binding of the antibody is then detected by virtue of a Label conjugated to this antibody. Where the antibody is unlabelled, a second labeled antibody may be employed, e.g., which is specific for the isotype of the anti-marker polypeptide antibody. Examples of labels which may be employed include radionuclides, fluorescence, chemiluminescence, and enzymes.

[0277] Where enzymes are employed, the Substrate for the enzyme may be added to the samples to provide a colored or fluorescent product. Examples of suitable enzymes for use in conjugates include horseradish peroxidase, alkaline phosphatase, malate dehydrogenase and the like. Where not commercially available, such antibody-enzyme conjugates are readily produced by techniques known to those skilled in the art.

[0278] In one embodiment, the assay is performed as a dot blot assay. The dot blot assay finds particular application where tissue samples are employed as it allows determination of the average amount of the marker polypeptide associated with a Single cell by correlating the amount of marker polypeptide in a cell-free extract produced from a predetermined number of cells.

[0279] In yet another embodiment, the invention contemplates using one or more antibodies which are generated against one or more of the marker polypeptides of this invention, which polypeptides are encoded by any of the polynucleotide sequences of the SEQ ID NO: 1 to 26 or 53 to 75. Such a panel of antibodies may be used as a reliable diagnostic probe for breast cancer. The assay of the present invention comprises contacting a biopsy sample containing cells, e.g., macrophages, with a panel of antibodies to one or more of the encoded products to determine the presence or absence of the marker polypeptides.

[0280] The diagnostic methods of the subject invention may also be employed as follow-up to treatment, e.g., quantification of the level of marker polypeptides may be indicative of the effectiveness of current or previously employed therapies for malignant neoplasia and breast cancer in particular as well as the effect of these therapies upon patient prognosis.

[0281] The diagnostic assays described above can be adapted to be used as prognostic assays, as well. Such an application takes advantage of the sensitivity of the assays of the Invention to events which take place at characteristic stages in the progression of plaque generation in case of malignant neoplasia. For example, a given marker gene may be up- or down-regulated at a very early stage, perhaps before the cell is developing into a foam cell, while another marker gene may be characteristically up or down regulated only at a much later stage. Such a method could involve the steps of contacting the mRNA of a test cell with a polynucleotide probe derived from a given marker polynucleotide which is expressed at different characteristic levels in breast cancer tissue cells at different stages of malignant neoplasia progression, and determining the approximate amount of hybridization of the probe to the mRNA of the cell, such amount being an indication of the level of expression of the gene in the cell, and thus an indication of the stage of disease progression of the cell; alternatively, the assay can be carried out with an antibody specific for the gene

product of the given marker polynucleotide, contacted with the proteins of the test cell. A battery of such tests will disclose not only the existence of a certain arteriosclerotic plaque, but also will allow the clinician to select the mode of treatment most appropriate for the disease, and to predict the likelihood of success of that treatment.

[0282] The methods of the invention can also be used to follow the clinical course of a given breast cancer predisposition. For example, the assay of the Invention can be applied to a blood sample from a patient; following treatment of the patient for BREAST CANCER, another blood sample is taken and the test repeated. Successful treatment will result in removal of demonstrate differential expression, characteristic of the breast cancer tissue cells, perhaps approaching or even surpassing normal levels.

Polypeptide activity

[0283] In one embodiment the present invention provides a method for screening potentially therapeutic agents which modulate the activity of one or more "BREAST CANCER GENE" polypeptides, such that if the activity of the polypeptide is increased as a result of the upregulation of the "BREAST CANCER GENE" in a subject having or at risk for malignant neoplasia and breast cancer in particular, the therapeutic substance will decrease the activity of the polypeptide relative to the activity of the some polypeptide in a subject not having or not at risk for malignant neoplasia or breast cancer in particular but not treated with the therapeutic agent. Likewise, if the activity of the polypeptide as a result of the downregulation of the "BREAST CANCER GENE" is decreased in a subject having or at risk for malignant neoplasia or breast cancer in particular, the therapeutic agent will increase the activity of the polypeptide relative to the activity of the same polypeptide in a subject not having or not at risk for malignant neoplasia or breast cancer in particular, but not treated with the therapeutic agent.

[0284] The activity of the "BREAST CANCER GENE" polypeptides indicated in Table 2 or 3 may be measured by any means known to those of skill in the art, and which are particular for the type of activity performed by the particular polypeptide. Examples of specific assays which may be used to measure the activity of particular polynucleotides are shown below.

a) G protein coupled receptors

[0285] In one embodiment, the "BREAST CANCER GENE" polynucleotide may encode a G protein coupled receptor. In one embodiment, the present invention provides a method of screening potential modulators (inhibitors or activators) of the G protein coupled receptor by measuring changes in the activity of the receptor in the presence of a candidate modulator.

1. G_i-coupled receptors

[0286] Cells (such as CHO cells or primary cells) are stably transfected with the relevant receptor and with an inducible CRE-luciferase construct. Cells are grown in 50% Dulbecco's modified Eagle medium / 50% F12 (DMEM/F12) supplemented with 10% FBS, at 37°C in a humidified atmosphere with 10% CO₂ and are routinely split at a ratio of 1:10 every 2 or 3 days. Test cultures are seeded into 384 - well plates at an appropriate density (e.g. 2000 cells / well in 35 µl cell culture medium) in DMEM/F12 with FBS, and are grown for 48 hours (range:~ 24 - 60 hours, depending on cell line). Growth medium is then exchanged against serum free medium (SFM; e.g. Ultra-CHO), containing 0,1% BSA. Test compounds dissolved in DMSO are diluted in SFM and transferred to the test cultures (maximal final concentration 10 µmolar), followed by addition of forskolin (~ 1 µmolar, final conc.) in SFM + 0,1% BSA 10 minutes later. In case of antagonist screening both, an appropriate concentration of agonist, and forskolin are added. The plates are incubated at 37°C in 10% CO₂ for 3 hours. Then the supernatant is removed, cells are lysed with lysis reagent (25 mmolar phosphate-buffer, pH 7,8, containing 2 mmolar DDT, 10% glycerol and 3% Triton X100). The luciferase reaction is started by addition of substrate-buffer (e.g. luciferase assay reagent, Promega) and luminescence is immediately determined (e.g. Berthold luminometer or Hamamatsu camera system).

2. G_s-coupled receptors

[0287] Cells (such as CHO cells or primary cells) are stably transfected with the relevant receptor and with an inducible CRE-luciferase construct. Cells are grown in 50% Dulbecco's modified Eagle medium / 50% F12 (DMEM/F12) supplemented with 10% FBS, at 37°C in a humidified atmosphere with 10% CO₂ and are routinely split at a ratio of 1:10 every 2 or 3 days. Test cultures are seeded into 384 - well plates at an appropriate density (e.g. 1000 or 2000 cells / well in 35 µl cell culture medium) in DMEM/F12 with FBS, and are grown for 48 hours (range:~ 24 - 60 hours, depending on cell line). The assay is started by addition of test-compounds in serum free medium (SFM; e.g. Ultra-CHO) containing 0,1% BSA: Test compounds are dissolved in DMSO, diluted in SFM and transferred to the test cultures (maximal final

concentration 10 μ molar, DMSO conc. < 0,6 %). In case of antagonist screening an appropriate concentration of agonist is added 5 - 10 minutes later. The plates are incubated at 37°C in 10% CO₂ for 3 hours. Then the cells are lysed with 10 μ l lysis reagent per well (25 mmolar phosphate-buffer, pH 7,8 , containing 2 mmolar DDT, 10% glycerol and 3% Triton X100) and the luciferase reaction is started by addition of 20 μ l substrate-buffer per well (e.g. luciferase assay reagent, Promega). Measurement of luminescence is started immediately (e.g. Berthold luminometer or Hamamatsu camera system).

3. G_q-coupled receptors

[0288] Cells (such as CHO cells or primary cells) are stably transfected with the relevant receptor. Cells expressing functional receptor protein are grown in 50% Dulbecco's modified Eagle medium / 50% F12 (DMEM/F12) supplemented with 10% FBS, at 37°C in a humidified atmosphere with 5% CO₂ and are routinely split at a cell line dependent ratio every 3 or 4 days. Test cultures are seeded into 384 - well plates at an appropriate density (e.g. 2000 cells / well in 35 μ l cell culture medium) in DMEM/F12 with FBS, and are grown for 48 hours (range: ~ 24 - 60 hours, depending on cell line). Growth medium is then exchanged against physiological salt solution (e.g. Tyrode solution). Test compounds dissolved in DMSO are diluted in Tyrode solution containing 0.1% BSA and transferred to the test cultures (maximal final concentration 10 molar). After addition of the receptor specific agonist the resulting G_q-mediated intracellular calcium increase is measured using appropriate read-out systems (e.g. calcium-sensitive dyes).

b) Ion channels

[0289] Ion channels are integral membrane proteins involved in electrical signaling, transmembrane signal transduction, and electrolyte and solute transport. By forming macromolecular pores through the membrane lipid bilayer, ion channels account for the flow of specific ion species driven by the electrochemical potential gradient for the permeating ion. At the single molecule level, individual channels undergo conformational transitions ("gating") between the 'open' (ion conducting) and 'closed' (non conducting) state. Typical single channel openings last for a few milliseconds and result in elementary transmembrane currents in the range of 10⁻⁹ - 10⁻¹² Ampere. Channel gating is controlled by various chemical and/or biophysical parameters, such as neurotransmitters and intracellular second messengers ('ligand-gated' channels) or membrane potential ('voltage-gated' channels). Ion channels are functionally characterized by their ion selectivity, gating properties, and regulation by hormones and pharmacological agents. Because of their central role in signaling and transport processes, ion channels present ideal targets for pharmacological therapeutics in various pathophysiological settings.

[0290] In one embodiment, the "BREAST CANCER GENE" may encode an ion channel. In one embodiment, the present invention provides a method of screening potential activators or inhibitors of channels activity of the "BREAST CANCER GENE" polypeptide. Screening for compounds interaction with ion channels to either inhibit or promote their activity can be based on (1.) binding and (2.) functional assays in living cells [Hille (183)].

1. For ligand-gated channels, e.g. ionotropic neurotransmitter/hormone receptors, assays can be designed detecting binding to the target by competition between the compound and a labeled ligand.

2. Ion channel function can be tested functionally in living cells. Target proteins are either expressed endogenously in appropriate reporter cells or are introduced recombinantly. Channel activity can be monitored by (2.1) concentration changes of the permeating ion (most prominently Ca²⁺ ions), (2.2) by changes in the transmembrane electrical potential gradient, and (2.3) by measuring a cellular response (e.g. expression of a reporter gene, secretion of a neurotransmitter) triggered or modulated by the target activity.

2.1 Channel activity results in transmembrane ion fluxes. Thus activation of ionic channels can be monitored by the resulting changes in intracellular ion concentrations using luminescent or fluorescent indicators. Because of its wide dynamic range and availability of suitable indicators this applies particularly to changes in intracellular Ca²⁺ ion concentration ([Ca²⁺]_i). [Ca²⁺]_i can be measured, for example, by aequorin luminescence or fluorescence dye technology (e.g. using Fluo-3, Indo-1, Fura-2). Cellular assays can be designed where either the Ca²⁺ flux through the target channel itself is measured directly or where modulation of the target channel affects membrane potential and thereby the activity of co-expressed voltage-gated Ca²⁺ channels.

2.2 Ion channel currents result in changes of electrical membrane potential (V_m) which can be monitored directly using potentiometric fluorescent probes. These electrically charged indicators (e.g. the anionic oxonol dye DiBAC₄(3)) redistribute between extra- and intracellular compartment in response to voltage changes. The equilibrium distribution is governed by the Nernst-equation. Thus changes in membrane potential results

in concomitant changes in cellular fluorescence. Again, changes in V_m might be caused directly by the activity of the target ion channel or through amplification and/or prolongation of the signal by channels co-expressed in the same cell.

2.3 Target channel activity can cause cellular Ca^{2+} entry either directly or through activation of additional Ca^{2+} channel (see 2.1). The resulting intracellular Ca^{2+} signals regulate a variety of cellular responses, e.g. secretion or gene transcription. Therefore modulation of the target channel can be detected by monitoring secretion of a known hormone/transmitter from the target-expressing cell or through expression of a reporter gene (e.g. luciferase) controlled by an Ca^{2+} -responsive promoter element (e.g. cyclic AMP/ Ca^{2+} -responsive elements; CRE).

c) DNA-binding proteins and transcription factors

[0291] In one embodiment, the "BREAST CANCER GENE" may encode a DNA-binding protein or a transcription factor. The activity of such a DNA-binding protein or a transcription factor may be measured, for example, by a promoter assay which measures the ability of the DNA-binding protein or the transcription factor to initiate transcription of a test sequence linked to a particular promoter. In one embodiment, the present invention provides a method of screening test compounds for its ability to modulate the activity of such a DNA-binding protein or a transcription factor by measuring the changes in the expression of a test gene which is regulated by a promoter which is responsive to the transcription factor.

d) Promotor assays

[0292] A promoter assay was set up with a human hepatocellular carcinoma cell HepG2 that was stably transfected with a luciferase gene under the control of a gene of interest (e.g. thyroid hormone) regulated promoter. The vector 2xIROluc, which was used for transfection, carries a thyroid hormone responsive element (TRE) of two 12 bp inverted palindromes separated by an 8 bp spacer in front of a tk minimal promoter and the luciferase gene. Test cultures were seeded in 96 well plates in serum - free Eagle's Minimal Essential Medium supplemented with glutamine, tricine, sodium pyruvate, non - essential amino acids, insulin, selen, transferrin, and were cultivated in a humidified atmosphere at 10 % CO_2 at 37°C. After 48 hours of incubation serial dilutions of test compounds or reference compounds (L-T3, L-T4 e.g.) and co-stimulator if appropriate (final concentration 1 nM) were added to the cell cultures and incubation was continued for the optimal time (e.g. another 4-72 hours). The cells were then lysed by addition of buffer containing Triton X100 and luciferin and the luminescence of luciferase induced by T3 or other compounds was measured in a luminometer. For each concentration of a test compound replicates of 4 were tested. EC_{50} values for each test compound were calculated by use of the Graph Pad Prism Scientific software.

Screening Methods

[0293] The invention provides assays for screening test compounds which bind to or modulate the activity of a "BREAST CANCER GENE" polypeptide or a "BREAST CANCER GENE" polynucleotide. A test compound preferably binds to a "BREAST CANCER GENE" polypeptide or polynucleotide. More preferably, a test compound decreases or increases "BREAST CANCER GENE" activity by at least about 10, preferably about 50, more preferably about 75, 90, or 100% relative to the absence of the test compound.

Test Compounds

[0294] Test compounds can be pharmacological agents already known in the art or can be compounds previously unknown to have any pharmacological activity. The compounds can be naturally occurring or designed in the laboratory. They can be isolated from microorganisms, animals, or plants, and can be produced recombinant, or synthesised by chemical methods known in the art. If desired, test compounds can be obtained using any of the numerous combinatorial library methods known in the art, including but not limited to, biological libraries, spatially addressable parallel solid phase or solution phase libraries, synthetic library methods requiring de-convolution, the one-bead one-compound library method, and synthetic library methods using affinity chromatography selection. The biological library approach is limited to polypeptide libraries, while the other four approaches are applicable to polypeptide, non-peptide oligomer, or small molecule libraries of compounds. [For review see Lam, 1997, (151)].

[0295] Methods for the synthesis of molecular libraries are well known in the art [see, for example, DeWitt et al., 1993, (152); Erb et al., 1994, (153); Zuckermann et al., 1994, (154); Cho et al., 1993, (155); Carell et al., 1994, (156) and Gallop et al., 1994, (157)]. Libraries of compounds can be presented in solution [see, e.g., Houghten, 1992, (158)],

or on beads [Lam, 1991, (159)], DNA-chips [Fodor, 1993, (160)], bacteria or spores (Ladner, U.S. Patent 5,223,409), plasmids [Cull et al., 1992, (161)], or phage [Scott & Smith, 1990, (162); Devlin, 1990, (163); Cwirla et al., 1990, (164); Felici, 1991, (165)].

5 High Throughput Screening

[0296] Test compounds can be screened for the ability to bind to "BREAST CANCER GENE" polypeptides or polynucleotides or to affect "BREAST CANCER GENE" activity or "BREAST CANCER GENE" expression using high throughput screening. Using high throughput screening, many discrete compounds can be tested in parallel so that
10 large numbers of test compounds can be quickly screened. The most widely established techniques utilize 96-well, 384-well or 1536-well microtiter plates. The wells of the microtiter plates typically require assay volumes that range from 5 to 500 μ l. In addition to the plates, many instruments, materials, pipettors, robotics, plate washers, and plate readers are commercially available to fit the microwell formats.

[0297] Alternatively, free format assays, or assays that have no physical barrier between samples, can be used. For
15 example, an assay using pigment cells (melanocytes) in a simple homogeneous assay for combinatorial peptide libraries is described by Jayawickreme et al., (166). The cells are placed under agarose in culture dishes, then beads that carry combinatorial compounds are placed on the surface of the agarose. The combinatorial compounds are partially released the compounds from the beads. Active compounds can be visualised as dark pigment areas because, as the compounds diffuse locally into the gel matrix, the active compounds cause the cells to change colors.

[0298] Another example of a free format assay is described by Chelsky, (167). Chelsky placed a simple homogenous
20 enzyme assay for carbonic anhydrase inside an agarose gel such that the enzyme in the gel would cause a color change throughout the gel.

[0299] Thereafter, beads carrying combinatorial compounds via a photolinker were placed inside the gel and the
25 compounds were partially released by UV light. Compounds that inhibited the enzyme were observed as local zones of inhibition having less color change.

[0300] In another example, combinatorial libraries were screened for compounds that had cytotoxic effects on cancer
cells growing in agar [Salmon et al., 1996, (168)].

[0301] Another high throughput screening method is described in Beutel et al., U.S. Patent 5,976,813. In this method,
30 test samples are placed in a porous matrix. One or more assay components are then placed within, on top of, or at the bottom of a matrix such as a gel, a plastic sheet, a filter, or other form of easily manipulated solid support. When samples are introduced to the porous matrix they diffuse sufficiently slowly, such that the assays can be performed without the test samples running together.

35 Binding Assays

[0302] For binding assays, the test compound is preferably a small molecule which binds to and occupies, for ex-
ample, the ATP/GTP binding site of the enzyme or the active site of a "BREAST CANCER GENE" polypeptide, such that normal biological activity is prevented. Examples of such small molecules include, but are not limited to, small peptides or peptide-like molecules.

[0303] In binding assays, either the test compound or a "BREAST CANCER GENE" polypeptide can comprise a
40 detectable label, such as a fluorescent, radioisotopic, chemiluminescent, or enzymatic label, such as horseradish peroxidase, alkaline phosphatase, or luciferase. Detection of a test compound which is bound to a "BREAST CANCER GENE" polypeptide can then be accomplished, for example, by direct counting of radioemission, by scintillation counting, or by determining conversion of an appropriate substrate to a detectable product.

[0304] Alternatively, binding of a test compound to a "BREAST CANCER GENE" polypeptide can be determined
45 without labeling either of the interactants. For example, a microphysiometer can be used to detect binding of a test compound with a "BREAST CANCER GENE" polypeptide. A microphysiometer (e.g., CytosensorJ) is an analytical instrument that measures the rate at which a cell acidifies its environment using a light-addressable potentiometric sensor (LAPS). Changes in this acidification rate can be used as an indicator of the interaction between a test compound
50 and a "BREAST CANCER GENE" polypeptide [McConnell et al., 1992, (169)].

[0305] Determining the ability of a test compound to bind to a "BREAST CANCER GENE" polypeptide also can be
accomplished using a technology such as real-time Bimolecular Interaction Analysis (BIA) [Sjolander & Urbaniczky, 1991, (170), and Szabo et al., 1995, (171)]. BIA is a technology for studying biospecific interactions in real time, without labeling any of the interactants (e.g., BIAcore™). Changes in the optical phenomenon surface plasmon resonance
55 (SPR) can be used as an indication of real-time reactions between biological molecules.

[0306] In yet another aspect of the invention, a "BREAST CANCER GENE" polypeptide can be used as a "bait
protein" in a two-hybrid assay or three-hybrid assay [see, e.g., U.S. Patent 5,283,317; Zervos et al., 1993, (172); Madura et al., 1993, (173); Bartel et al., 1993, (174); Iwabuchi et al., 1993, (175) and Brent WO 94/10300], to identify

other proteins which bind to or interact with the "BREAST CANCER GENE" polypeptide and modulate its activity.

[0307] The two-hybrid system is based on the modular nature of most transcription factors, which consist of separable DNA-binding and activation domains. Briefly, the assay utilizes two different DNA constructs. For example, in one construct, polynucleotide encoding a "BREAST CANCER GENE" polypeptide can be fused to a polynucleotide encoding the DNA binding domain of a known transcription factor (e.g., GAL4). In the other construct a DNA sequence that encodes an unidentified protein ("prey" or "sample") can be fused to a polynucleotide that codes for the activation domain of the known transcription factor. If the "bait" and the "prey" proteins are able to interact in vivo to form an protein- dependent complex, the DNA-binding and activation domains of the transcription factor are brought into close proximity. This proximity allows transcription of a reporter gene (e.g., LacZ), which is operably linked to a transcriptional regulatory site responsive to the transcription factor. Expression of the reporter gene can be detected, and cell colonies containing the functional transcription factor can be isolated and used to obtain the DNA sequence encoding the protein which interacts with the "BREAST CANCER GENE" polypeptide.

[0308] It may be desirable to immobilize either a "BREAST CANCER GENE" polypeptide (or polynucleotide) or the test compound to facilitate separation of bound from unbound forms of one or both of the interactants, as well as to accommodate automation of the assay. Thus, either a "BREAST CANCER GENE" polypeptide (or polynucleotide) or the test compound can be bound to a solid support. Suitable solid supports include, but are not limited to, glass or plastic slides, tissue culture plates, microtiter wells, tubes, silicon chips, or particles such as beads (including, but not limited to, latex, polystyrene, or glass beads). Any method known in the art can be used to attach a "BREAST CANCER GENE" polypeptide (or polynucleotide) or test compound to a solid support, including use of covalent and non-covalent linkages, passive absorption, or pairs of binding moieties attached respectively to the polypeptide (or polynucleotide) or test compound and the solid support. Test compounds are preferably bound to the solid support in an array, so that the location of individual test compounds can be tracked. Binding of a test compound to a "BREAST CANCER GENE" polypeptide (or polynucleotide) can be accomplished in any vessel suitable for containing the reactants. Examples of such vessels include microtiter plates, test tubes, and microcentrifuge tubes.

[0309] In one embodiment, a "BREAST CANCER GENE" polypeptide is a fusion protein comprising a domain that allows the "BREAST CANCER GENE" polypeptide to be bound to a solid support. For example, glutathione S-transferase fusion proteins can be adsorbed onto glutathione sepharose beads (Sigma Chemical, St. Louis, Mo.) or glutathione derivatized microtiter plates, which are then combined with the test compound or the test compound and the nonadsorbed "BREAST CANCER GENE" polypeptide; the mixture is then incubated under conditions conducive to complex formation (e.g., at physiological conditions for salt and pH). Following incubation, the beads or microtiter plate wells are washed to remove any unbound components. Binding of the interactants can be determined either directly or indirectly, as described above. Alternatively, the complexes can be dissociated from the solid support before binding is determined.

[0310] Other techniques for immobilising proteins or polynucleotides on a solid support also can be used in the screening assays of the invention. For example, either a "BREAST CANCER GENE" polypeptide (or polynucleotide) or a test compound can be immobilized utilizing conjugation of biotin and streptavidin. Biotinylated "BREAST CANCER GENE" polypeptides (or polynucleotides) or test compounds can be prepared from biotin NHS (N-hydroxysuccinimide) using techniques well known in the art (e.g., biotinylation kit, Pierce Chemicals, Rockford, Ill.) and immobilized in the wells of streptavidin-coated 96 well plates (Pierce Chemical). Alternatively, antibodies which specifically bind to a "BREAST CANCER GENE" polypeptide, polynucleotide, or a test compound, but which do not interfere with a desired binding site, such as the ATP/GTP binding site or the active site of the "BREAST CANCER GENE" polypeptide, can be derivatised to the wells of the plate. Unbound target or protein can be trapped in the wells by antibody conjugation.

[0311] Methods for detecting such complexes, in addition to those described above for the GST-immobilized complexes, include immunodetection of complexes using antibodies which specifically bind to a "BREAST CANCER GENE" polypeptide or test compound, enzyme-linked assays which rely on detecting an activity of a "BREAST CANCER GENE" polypeptide, and SDS gel electrophoresis under non-reducing conditions.

[0312] Screening for test compounds which bind to a "BREAST CANCER GENE" polypeptide or polynucleotide also can be carried out in an intact cell. Any cell which comprises a "BREAST CANCER GENE" polypeptide or polynucleotide can be used in a cell-based assay system. A "BREAST CANCER GENE" polynucleotide can be naturally occurring in the cell or can be introduced using techniques such as those described above. Binding of the test compound to a "BREAST CANCER GENE" polypeptide or polynucleotide is determined as described above.

Modulation of Gene Expression

[0313] In another embodiment, test compounds which increase or decrease "BREAST CANCER GENE" expression are identified. A "BREAST CANCER GENE" polynucleotide is contacted with a test compound, and the expression of an RNA or polypeptide product of the "BREAST CANCER GENE" polynucleotide is determined. The level of expression of appropriate mRNA or polypeptide in the presence of the test compound is compared to the level of expression of

mRNA or polypeptide in the absence of the test compound. The test compound can then be identified as a modulator of expression based on this comparison. For example, when expression of mRNA or polypeptide is greater in the presence of the test compound than in its absence, the test compound is identified as a stimulator or enhancer of the mRNA or polypeptide expression. Alternatively, when expression of the mRNA or polypeptide is less in the presence of the test compound than in its absence, the test compound is identified as an inhibitor of the mRNA or polypeptide expression.

[0314] The level of "BREAST CANCER GENE" mRNA or polypeptide expression in the cells can be determined by methods well known in the art for detecting mRNA or polypeptide. Either qualitative or quantitative methods can be used. The presence of polypeptide products of a "BREAST CANCER GENE" polynucleotide can be determined, for example, using a variety of techniques known in the art, including immunochemical methods such as radioimmunoassay, Western blotting, and immunohistochemistry. Alternatively, polypeptide synthesis can be determined in vivo, in a cell culture, or in an in vitro translation system by detecting incorporation of labeled amino acids into a "BREAST CANCER GENE" polypeptide.

[0315] Such screening can be carried out either in a cell-free assay system or in an intact cell. Any cell which expresses a "BREAST CANCER GENE" polynucleotide can be used in a cell-based assay system. A "BREAST CANCER GENE" polynucleotide can be naturally occurring in the cell or can be introduced using techniques such as those described above. Either a primary culture or an established cell line, such as CHO or human embryonic kidney 293 cells, can be used.

Therapeutic Indications and Methods

[0316] Therapies for treatment of breast cancer primarily relied upon effective chemotherapeutic drugs for intervention on the cell proliferation, cell growth or angiogenesis. The advent of genomics-driven molecular target identification has opened up the possibility of identifying new breast cancer-specific targets for therapeutic intervention that will provide safer, more effective treatments for malignant neoplasia patients and breast cancer patients in particular. Thus, newly discovered breast cancer-associated genes and their products can be used as tools to develop innovative therapies. The identification of the Her2/neu receptor kinase presents exciting new opportunities for treatment of a certain subset of tumor patients as described before. Genes playing important roles in any of the physiological processes outlined above can be characterized as breast cancer targets. Genes or gene fragments identified through genomics can readily be expressed in one or more heterologous expression systems to produce functional recombinant proteins. These proteins are characterized in vitro for their biochemical properties and then used as tools in high-throughput molecular screening programs to identify chemical modulators of their biochemical activities. Modulators of target gene expression or protein activity can be identified in this manner and subsequently tested in cellular and in vivo disease models for therapeutic activity. Optimization of lead compounds with iterative testing in biological models and detailed pharmacokinetic and toxicological analyses form the basis for drug development and subsequent testing in humans.

[0317] This invention further pertains to the use of novel agents identified by the screening assays described above. Accordingly, it is within the scope of this invention to use a test compound identified as described herein in an appropriate animal model. For example, an agent identified as described herein (e.g., a modulating agent, an antisense polynucleotide molecule, a specific antibody, ribozyme, or a human "BREAST CANCER GENE" polypeptide binding molecule) can be used in an animal model to determine the efficacy, toxicity, or side effects of treatment with such an agent. Alternatively, an agent identified as described herein can be used in an animal model to determine the mechanism of action of such an agent. Furthermore, this invention pertains to uses of novel agents identified by the above described screening assays for treatments as described herein.

[0318] A reagent which affects human "BREAST CANCER GENE" activity can be administered to a human cell, either in vitro or in vivo, to reduce or increase human "BREAST CANCER GENE" activity. The reagent preferably binds to an expression product of a human "BREAST CANCER GENE". If the expression product is a protein, the reagent is preferably an antibody. For treatment of human cells ex vivo, an antibody can be added to a preparation of stem cells which have been removed from the body. The cells can then be replaced in the same or another human body, with or without clonal propagation, as is known in the art.

[0319] In one embodiment, the reagent is delivered using a liposome. Preferably, the liposome is stable in the animal into which it has been administered for at least about 30 minutes, more preferably for at least about 1 hour, and even more preferably for at least about 24 hours. A liposome comprises a lipid composition that is capable of targeting a reagent, particularly a polynucleotide, to a particular site in an animal, such as a human. Preferably, the lipid composition of the liposome is capable of targeting to a specific organ of an animal, such as the lung, liver, spleen, heart brain, lymph nodes, and skin.

[0320] A liposome useful in the present invention comprises a lipid composition that is capable of fusing with the plasma membrane of the targeted cell to deliver its contents to the cell. Preferably, the transfection efficiency of a liposome is about 0.5 μ g of DNA per 16 nmol of liposome delivered to about 10^6 cells, more preferably about 1.0 μ g

of DNA per 16 nmol of liposome delivered to about 10^6 cells, and even more preferably about 2.0 μg of DNA per 16 nmol of liposome delivered to about 10^6 cells. Preferably, a liposome is between about 100 and 500 nm, more preferably between about 150 and 450 nm, and even more preferably between about 200 and 400 nm in diameter.

[0321] Suitable liposomes for use in the present invention include those liposomes usually used in, for example, gene delivery methods known to those of skill in the art. More preferred liposomes include liposomes having a polycationic lipid composition and/or liposomes having a cholesterol backbone conjugated to polyethylene glycol. Optionally, a liposome comprises a compound capable of targeting the liposome to a particular cell type, such as a cell-specific ligand exposed on the outer surface of the liposome.

[0322] Complexing a liposome with a reagent such as an antisense oligonucleotide or ribozyme can be achieved using methods which are standard in the art (see, for example, U.S. Patent 5,705,151). Preferably, from about 0.1 μg to about 10 μg of polynucleotide is combined with about 8 nmol of liposomes, more preferably from about 0.5 μg to about 5 μg of polynucleotides are combined with about 8 nmol liposomes, and even more preferably about 1.0 μg of polynucleotides is combined with about 8 nmol liposomes.

[0323] In another embodiment, antibodies can be delivered to specific tissues in vivo using receptor-mediated targeted delivery. Receptor-mediated DNA delivery techniques are taught in, for example, Findeis et al., 1993, (176); Chiou et al., 1994, (177); Wu & Wu, 1988, (178); Wu et al., 1994, (179); Zenke et al., 1990, (180); Wu et al., 1991, (181).

Determination of a Therapeutically Effective Dose

[0324] The determination of a therapeutically effective dose is well within the capability of those skilled in the art. A therapeutically effective dose refers to that amount of active ingredient which increases or decreases human "BREAST CANCER GENE" activity relative to the human "BREAST CANCER GENE" activity which occurs in the absence of the therapeutically effective dose.

[0325] For any compound, the therapeutically effective dose can be estimated initially either in cell culture assays or in animal models, usually mice, rabbits, dogs, or pigs. The animal model also can be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans.

[0326] Therapeutic efficacy and toxicity, e.g., ED_{50} (the dose therapeutically effective in 50% of the population) and LD_{50} (the dose lethal to 50% of the population), can be determined by standard pharmaceutical procedures in cell cultures or experimental animals. The dose ratio of toxic to therapeutic effects is the therapeutic index, and it can be expressed as the ratio, $\text{LD}_{50}/\text{ED}_{50}$.

[0327] Pharmaceutical compositions which exhibit large therapeutic indices are preferred. The data obtained from cell culture assays and animal studies is used in formulating a range of dosage for human use. The dosage contained in such compositions is preferably within a range of circulating concentrations that include the ED_{50} with little or no toxicity. The dosage varies within this range depending upon the dosage form employed, sensitivity of the patient, and the route of administration.

[0328] The exact dosage will be determined by the practitioner, in light of factors related to the subject that requires treatment. Dosage and administration are adjusted to provide sufficient levels of the active ingredient or to maintain the desired effect. Factors which can be taken into account include the severity of the disease state, general health of the subject, age, weight, and gender of the subject, diet, time and frequency of administration, drug combination(s), reaction sensitivities, and tolerance/response to therapy. Long-acting pharmaceutical compositions can be administered every 3 to 4 days, every week, or once every two weeks depending on the half-life and clearance rate of the particular formulation.

[0329] Normal dosage amounts can vary from 0.1 to 100,000 micrograms, up to a total dose of about 1 g, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature and generally available to practitioners in the art. Those skilled in the art will employ different formulations for nucleotides than for proteins or their inhibitors. Similarly, delivery of polynucleotides or polypeptides will be specific to particular cells, conditions, locations, etc.

[0330] If the reagent is a single-chain antibody, polynucleotides encoding the antibody can be constructed and introduced into a cell either ex vivo or in vivo using well-established techniques including, but not limited to, transferrin-polycation-mediated DNA transfer, transfection with naked or encapsulated nucleic acids, liposome-mediated cellular fusion, intracellular transportation of DNA-coated latex beads, protoplast fusion, viral infection, electroporation, a gene gun, and DEAE- or calcium phosphate-mediated transfection.

[0331] Effective in vivo dosages of an antibody are in the range of about 5 μg to about 50 $\mu\text{g}/\text{kg}$, about 50 μg to about 5 mg/kg, about 100 μg to about 500 $\mu\text{g}/\text{kg}$ of patient body weight, and about 200 to about 250 $\mu\text{g}/\text{kg}$ of patient body weight. For administration of polynucleotides encoding single-chain antibodies, effective in vivo dosages are in the range of about 100 ng to about 200 ng, 500 ng to about 50 mg, about 1 μg to about 2 mg, about 5 μg to about 500 μg , and about 20 μg to about 100 μg of DNA.

[0332] If the expression product is mRNA, the reagent is preferably an antisense oligonucleotide or a ribozyme. Polynucleotides which express antisense oligonucleotides or ribozymes can be introduced into cells by a variety of methods, as described above.

[0333] Preferably, a reagent reduces expression of a "BREAST CANCER GENE" gene or the activity of a "BREAST CANCER GENE" polypeptide by at least about 10, preferably about 50, more preferably about 75, 90, or 100% relative to the absence of the reagent. The effectiveness of the mechanism chosen to decrease the level of expression of a "BREAST CANCER GENE" gene or the activity of a "BREAST CANCER GENE" polypeptide can be assessed using methods well known in the art, such as hybridization of nucleotide probes to "BREAST CANCER GENE"-specific mRNA, quantitative RT-PCR, immunologic detection of a "BREAST CANCER GENE" polypeptide, or measurement of "BREAST CANCER GENE" activity.

[0334] In any of the embodiments described above, any of the pharmaceutical compositions of the invention can be administered in combination with other appropriate therapeutic agents. Selection of the appropriate agents for use in combination therapy can be made by one of ordinary skill in the art, according to conventional pharmaceutical principles. The combination of therapeutic agents can act synergistically to effect the treatment or prevention of the various disorders described above. Using this approach, one may be able to achieve therapeutic efficacy with lower dosages of each agent, thus reducing the potential for adverse side effects.

[0335] Any of the therapeutic methods described above can be applied to any subject in need of such therapy, including, for example, birds and mammals such as dogs, cats, cows, pigs, sheep, goats, horses, rabbits, monkeys, and most preferably, humans.

[0336] All patents and patent applications cited in this disclosure are expressly incorporated herein by reference. The above disclosure generally describes the present invention. A more complete understanding can be obtained by reference to the following specific examples which are provided for purposes of illustration only and are not intended to limit the scope of the invention.

Pharmaceutical Compositions

[0337] The invention also provides pharmaceutical compositions which can be administered to a patient to achieve a therapeutic effect. Pharmaceutical compositions of the invention can comprise, for example, a "BREAST CANCER GENE" polypeptide, "BREAST CANCER GENE" polynucleotide, ribozymes or antisense oligonucleotides, antibodies which specifically bind to a "BREAST CANCER GENE" polypeptide, or mimetics, agonists, antagonists, or inhibitors of a "BREAST CANCER GENE" polypeptide activity. The compositions can be administered alone or in combination with at least one other agent, such as stabilizing compound, which can be administered in any sterile, biocompatible pharmaceutical carrier, including, but not limited to, saline, buffered saline, dextrose, and water. The compositions can be administered to a patient alone, or in combination with other agents, drugs or hormones.

[0338] In addition to the active ingredients, these pharmaceutical compositions can contain suitable pharmaceutically acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically. Pharmaceutical compositions of the invention can be administered by any number of routes including, but not limited to, oral, intravenous, intramuscular, intraarterial, intramedullary, intrathecal, intraventricular, transdermal, subcutaneous, intraperitoneal, intranasal, parenteral, topical, sublingual, or rectal means. Pharmaceutical compositions for oral administration can be formulated using pharmaceutically acceptable carriers well known in the art in dosages suitable for oral administration. Such carriers enable the pharmaceutical compositions to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions, and the like, for ingestion by the patient.

[0339] Pharmaceutical preparations for oral use can be obtained through combination of active compounds with solid excipient, optionally grinding a resulting mixture, and processing the mixture of granules, after adding suitable auxiliaries, if desired, to obtain tablets or dragee cores. Suitable excipients are carbohydrate or protein fillers, such as sugars, including lactose, sucrose, mannitol, or sorbitol; starch from corn, wheat, rice, potato, or other plants; cellulose, such as methyl cellulose, hydroxypropylmethylcellulose, or sodium carboxymethylcellulose; gums including arabic and tragacanth; and proteins such as gelatin and collagen. If desired, disintegrating or solubilizing agents can be added, such as the cross-linked polyvinyl pyrrolidone, agar, alginic acid, or a salt thereof, such as sodium alginate.

[0340] Dragee cores can be used in conjunction with suitable coatings, such as concentrated sugar solutions, which also can contain gum arabic, talc, polyvinylpyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments can be added to the tablets or dragee coatings for product identification or to characterize the quantity of active compound, i.e., dosage.

[0341] Pharmaceutical preparations which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a coating, such as glycerol or sorbitol. Push-fit capsules can contain active ingredients mixed with a filler or binders, such as lactose or starches, lubricants, such as talc or magnesium stearate, and, optionally, stabilizers. In soft capsules, the active compounds can be dissolved or suspended in suitable liquids,

such as fatty oils, liquid, or liquid polyethylene glycol with or without stabilizers.

[0342] Pharmaceutical formulations suitable for parenteral administration can be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hanks' solution, Ringer's solution, or physiologically buffered saline. Aqueous injection suspensions can contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Additionally, suspensions of the active compounds can be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate or triglycerides, or liposomes. Non-lipid polycationic amino polymers also can be used for delivery. Optionally, the suspension also can contain suitable stabilizers or agents which increase the solubility of the compounds to allow for the preparation of highly concentrated solutions. For topical or nasal administration, penetrants appropriate to the particular barrier to be permeated are used in the formulation. Such penetrants are generally known in the art.

[0343] The pharmaceutical compositions of the present invention can be manufactured in a manner that is known in the art, e.g., by means of conventional mixing, dissolving, granulating, dragee making, levigating, emulsifying, encapsulating, entrapping, or lyophilizing processes. The pharmaceutical composition can be provided as a salt and can be formed with many acids, including but not limited to, hydrochloric, sulfuric, acetic, lactic, tartaric, malic, succinic, etc. Salts tend to be more soluble in aqueous or other protonic solvents than are the corresponding free base forms. In other cases, the preferred preparation can be a lyophilized powder which can contain any or all of the following: 150 mM histidine, 0.1 %2% sucrose, and 27% mannitol, at a pH range of 4.5 to 5.5, that is combined with buffer prior to use.

[0344] Further details on techniques for formulation and administration can be found in the latest edition of REMINGTON'S PHARMACEUTICAL SCIENCES (182). After pharmaceutical compositions have been prepared, they can be placed in an appropriate container and labeled for treatment of an indicated condition. Such labeling would include amount, frequency, and method of administration.

Material and Methods

[0345] One strategy for identifying genes that are involved in breast cancer is to detect genes that are expressed differentially under conditions associated with the disease versus non-disease conditions. The sub-sections below describe a number of experimental systems which may be used to detect such differentially expressed genes. In general, these experimental systems include at least one experimental condition in which subjects or samples are treated in a manner associated with breast cancer, in addition to at least one experimental control condition lacking such disease associated treatment. Differentially expressed genes are detected, as described below, by comparing the pattern of gene expression between the experimental and control conditions.

[0346] Once a particular gene has been identified through the use of one such experiment, its expression pattern may be further characterized by studying its expression in a different experiment and the findings may be validated by an independent technique. Such use of multiple experiments may be useful in distinguishing the roles and relative importance of particular genes in breast cancer. A combined approach, comparing gene expression pattern in cells derived from breast cancer patients to those of *in vitro* cell culture models can give substantial hints on the pathways involved in development and/or progression of breast cancer.

[0347] Among the experiments which may be utilized for the identification of differentially expressed genes involved in malignant neoplasia and breast cancer, for example, are experiments designed to analyze those genes which are involved in signal transduction. Such experiments may serve to identify genes involved in the proliferation of cells.

[0348] Below are methods described for the identification of genes which are involved in breast cancer. Such represent genes which are differentially expressed in breast cancer conditions relative to their expression in normal, or non-breast cancer conditions or upon experimental manipulation based on clinical observations. Such differentially expressed genes represent "target" and/or "marker" genes. Methods for the further characterization of such differentially expressed genes, and for their identification as target and/or marker genes, are presented below.

[0349] Alternatively, a differentially expressed gene may have its expression modulated, i.e., quantitatively increased or decreased, in normal versus breast cancer states, or under control versus experimental conditions. The degree to which expression differs in normal versus breast cancer or control versus experimental states need only be large enough to be visualized via standard characterization techniques, such as, for example, the differential display technique described below. Other such standard characterization techniques by which expression differences may be visualized include but are not limited to quantitative RT-PCR and Northern analyses, which are well known to those of skill in the art.

EXAMPLE 1*Expression profiling*5 a) Expression profiling utilizing quantitative RT-PCR

[0350] For a detailed analysis of gene expression by quantitative PCR methods, one will utilize primers flanking the genomic region of interest and a fluorescent labeled probe hybridizing in-between. Using the PRISM 7700 Sequence Detection System of PE Applied Biosystems (Perkin Elmer, Foster City, CA, USA) with the technique of a fluorogenic probe, consisting of an oligonucleotide labeled with both a fluorescent reporter dye and a quencher dye, one can perform such a expression measurement. Amplification of the probe-specific product causes cleavage of the probe, generating an increase in reporter fluorescence. Primers and probes were selected using the Primer Express software and localized mostly in the 3' region of the coding sequence or in the 3' untranslated region (see Table 5 for primer- and probe- sequences) according to the relative positions of the probe sequence used for the construction of the Affymetrix HG_U95A-E or HG-U133A-B DNA-chips. All primer pairs were checked for specificity by conventional PCR reactions. To standardize the amount of sample RNA, GAPDH was selected as a reference, since it was not differentially regulated in the samples analyzed. TaqMan validation experiments were performed showing that the efficiencies of the target and the control amplifications are approximately equal which is a prerequisite for the relative quantification of gene expression by the comparative $\Delta\Delta C_T$ method, known to those with skills in the art.

[0351] As well as the technology provided by Perkin Elmer one may use other technique implementations like Lightcycler™ from Roche Inc. or iCycler from Stratagene Inc..

b) Expression profiling utilizing DNA microarrays

[0352] Expression profiling can be carried out using the Affymetrix Array Technology. By hybridization of mRNA to such a DNA-array or DNA-Chip, it is possible to identify the expression value of each transcripts due to signal intensity at certain position of the array. Usually these DNA-arrays are produced by spotting of cDNA, oligonucleotides or sub-cloned DNA fragments. In case of Affymetrix technology app. 400.000 individual oligonucleotide sequences were synthesized on the surface of a silicon wafer at distinct positions. The minimal length of oligomers is 12 nucleotides, preferable 25 nucleotides or full length of the questioned transcript. Expression profiling may also be carried out by hybridization to nylon or nitrocellulose membrane bound DNA or oligonucleotides. Detection of signals derived from hybridization may be obtained by either colorimetric, fluorescent, electrochemical, electronic, optic or by radioactive readout. Detailed description of array construction have been mentioned above and in other patents cited. To determine the quantitative and qualitative changes in the chromosomal region to analyze, RNA from tumor tissue which is suspected to contain such genomic alterations has to be compared to RNA extracted from benign tissue (e.g. epithelial breast tissue, or micro dissected ductal tissue) on the basis of expression profiles for the whole transcriptome. With minor modifications, the sample preparation protocol followed the Affymetrix GeneChip Expression Analysis Manual (Santa Clara, CA). Total RNA extraction and isolation from tumor or benign tissues, biopsies, cell isolates or cell containing body fluids can be performed by using TRIzol (Life Technologies, Rockville, MD) and Oligotex mRNA Midi kit (Qiagen, Hilden, Germany), and an ethanol precipitation step should be carried out to bring the concentration to 1 mg/ml. Using 5-10 mg of mRNA to create double stranded cDNA by the Superscript system (Life Technologies). First strand cDNA synthesis was primed with a T7-(dT24) oligonucleotide. The cDNA can be extracted with phenol/chloroform and precipitated with ethanol to a final concentration of 1 mg /ml. From the generated cDNA, cRNA can be synthesized using Enzo's (Enzo Diagnostics Inc., Farmingdale, NY) *in vitro* Transcription Kit. Within the same step the cRNA can be labeled with biotin nucleotides Bio-11-CTP and Bio-16-UTP (Enzo Diagnostics Inc., Farmingdale, NY) . After labeling and cleanup (Qiagen, Hilden (Germany) the cRNA then should be fragmented in an appropriated fragmentation buffer (e.g., 40 mM Tris-Acetate, pH 8.1, 100 mM KOAc, 30 mM MgOAc, for 35 minutes at 94°C). As per the Affymetrix protocol, fragmented cRNA should be hybridized on the HG_U133 arrays A and B, comprising app. 40.000 probed transcripts each, for 24 hours at 60 rpm in a 45°C hybridization oven. After Hybridization step the chip surfaces have to be washed and stained with streptavidin phycoerythrin (SAPE; Molecular Probes, Eugene, OR) in Affymetrix fluidics stations. To amplify staining, a second labeling step can be introduced, which is recommended but not compulsive. Here one should add SAPE solution twice with an antistreptavidin biotinylated antibody. Hybridization to the probe arrays may be detected by fluorometric scanning (Hewlett Packard Gene Array Scanner; Hewlett Packard Corporation, Palo Alto, CA).

[0353] After hybridization and scanning, the microarray images can be analyzed for quality control, looking for major chip defects or abnormalities in hybridization signal. Therefore either Affymetrix GeneChip MAS 5.0 Software or other microarray image analysis software can be utilized. Primary data analysis should be carried out by software provided by the manufacturer..

[0354] In case of the genes analyses in one embodiment of this invention the primary data have been analyzed by further bioinformatic tools and additional filter criteria. The bioinformatic analysis is described in detail below.

c) Data analysis

[0355] According to Affymetrix measurement technique (Affymetrix GeneChip Expression Analysis Manual, Santa Clara, CA) a single gene expression measurement on one chip yields the average difference value and the absolute call. Each chip contains 16-20 oligonucleotide probe pairs per gene or cDNA clone. These probe pairs include perfectly matched sets and mismatched sets, both of which are necessary for the calculation of the average difference, or expression value, a measure of the intensity difference for each probe pair, calculated by subtracting the intensity of the mismatch from the intensity of the perfect match. This takes into consideration variability in hybridization among probe pairs and other hybridization artifacts that could affect the fluorescence intensities. The average difference is a numeric value supposed to represent the expression value of that gene. The absolute call can take the values 'A' (absent), 'M' (marginal), or 'P' (present) and denotes the quality of a single hybridization. We used both the quantitative information given by the average difference and the qualitative information given by the absolute call to identify the genes which are differentially expressed in biological samples from individuals with breast cancer versus biological samples from the normal population. With other algorithms than the Affymetrix one we have obtained different numerical values representing the same expression values and expression differences upon comparison.

[0356] The differential expression E in one of the breast cancer groups compared to the normal population is calculated as follows. Given n average difference values d_1, d_2, \dots, d_n in the breast cancer population and m average difference values c_1, c_2, \dots, c_m in the population of normal individuals, it is computed by the equation:

$$E \equiv \exp \left(\frac{1}{m} \sum_{i=1}^m \ln(c_i) - \frac{1}{n} \sum_{i=1}^n \ln(d_i) \right)$$

If $d_j < 50$ or $c_i < 50$ for one or more values of i and j, these particular values c_i and/or d_j are set to an "artificial" expression value of 50. These particular computation of E allows for a correct comparison to TaqMan results.

[0357] A gene is called up-regulated in breast cancer versus normal if $E \geq 1.5$ and if the number of absolute calls equal to 'P' in the breast cancer population is greater than $n/2$.

[0358] A gene is called down-regulated in breast cancer versus normal if $E \leq 1.5$ and if the number of absolute calls equal to 'P' in the normal population is greater than $m/2$.

[0359] The final list of differentially regulated genes consists of all up-regulated and all down-regulated genes in biological samples from individuals with breast cancer versus biological samples from the normal population. Those genes on this list which are interesting for a pharmaceutical application were finally validated by TaqMan. If a good correlation between the expression values/behavior of a transcript could be observed with both techniques, such a gene is listed in Tables 1 to 3.

[0360] Since not only the information on differential expression of a single gene within an identified ARCHEON, but also the information on the co-regulation of several members is important for predictive, diagnostic, preventive and therapeutic purposes we have combined expression data with information on the chromosomal position (e.g. golden path) taken from public available databases to develop a picture of the overall transcriptom of a given tumor sample. By this technique not only known or suspected regions of genomes can be inspected but even more valuable, new regions of dysregulation with chromosomal linkage can be identified. This is of value in other types of neoplasia or viral integration and chromosomal rearrangements. By SQL based database searches one can retrieve information on expression, qualitative value of a measurement (denoted by Affymetrix MAS 5.0 Software), expression values derived from other techniques than DNA-chip hybridization and chromosomal linkage.

EXAMPLE 2

Identification of the ARCHEON

a) Identification and localization of genes or gene probes (represented by the so called probe sets on Affymetrix arrays HG-U95A-E or HG-U133A-B) in their chromosomal context and order on the human genome.

[0361] For identification of larger chromosomal changes or aberrations, as they have been described in detail above,

a sufficient number of genes, transcripts or DNA-fragments is needed. The density of probes covering a chromosomal region is not necessarily limited to the transcribed genes, in case of the use of array based CGH but by utilizing RNA as probe material the density is given by the distance of genes on a chromosome. The DNA-microarrays provided by Affymetrix Inc. Do contain hitherto all transcripts from the known humane genome, which are be represented by 40.000 - 60.000 probe sets. By BLAST mapping and sorting the sequences of these short DNA-oligomers to the public available sequence of the human genome represented by the so called "golden path", available at the university of California in Santa Cruz or from the NCBI, a chromosomal display of the whole Transcriptome of a tissue specimen evolves. By graphical display of the individual chromosomal regions and color coding of over or under represented transcripts, compared to a reference transcriptome regions with DNA gains and losses can be identified.

b) Quantification of gene copy numbers by combined IHC and quantitative PCR (PCR karyotyping) or directly by quantitative PCR

[0362] Usually one to three paraffin-embedded tissue sections that are 5 µm thick are used to obtain genomic DNA from the samples. Tissue section are stained by colorimetric IHC after deparaffinization to identify regions containing disease associated cells. Stained regions are macrodissected with a scalpel and transferred into a microcentrifuge tube. The genomic DNA of these isolated tissue sections is extracted using appropriate buffers. The isolated DNA is then used for quantitative PCR with appropriate primers and probes. Optionally the IHC staining can be omitted and the genomic DNA can be directly isolated with or without prior deparaffinization with appropriate buffers. Those who are skilled in the art may vary the conditions and buffers described below to obtain equivalent results.

[0363] Reagents from DAKO (HercepTest Code No. K 5204) and TaKaRa were used (Biomedicals Cat.: 9091) according to the manufactures protocol.

[0364] It is convenient to prepare the following reagents prior to staining:

Solution No. 7

[0365] Epitope Retrieval Solution (Citrate buffer + antimicrobial agent) (10xconc.) 20 ml ad 200 ml aqua dest. (stable for 1month at 2-8°C)

Solution No. 8

[0366] Washing-buffer (Tris-HCl + antimicrobial agent) (10 x conc.) 30 ml ad 300 ml distilled water (stable for 1month at 2-8°C)

Staining solution: DAB

[0367] 1 ml solution is sufficient for 10 slides. The solution were prepared immediately before usage.:

[0368] 1 ml DAB buffer (Substrate Buffer solution, pH 7.5, containing H₂O₂, stabilizer, enhancers and an antimicrobial agent) + 1 drop (25-3 µl) DAB-Chromogen (3,3'-diaminobenzidine chromogen solution). This solution is stable for up to 5 days at 2-8°C. Precipitated substances do not influence the staining result. Additionally required are: 2x approx. 100 ml Xylol, 2x approx. 100 ml Ethanol 100%, 2x Ethanol 95%, aqua dest. These solution can be used for up to 40 stainings. A water bath is required for the epitope retrieval step.

Staining procedure:

[0369] All reagents are pre-warmed to room temperature (20-25°C) prior to immuno-staining. Likewise all incubations were performed at room temperature. Except the epitope retrieval which is performed in at 95°C water bath. Between the steps excess of liquid is tapped off from the slides with lintless tissue (Kim Wipe).

Deparaffinization

[0370] Slides are placed in a xylene bath and incubated for 5 minutes. The bath is changed and the step repeated once. Excess of liquid is tapped off and the slides are placed in absolute ethanol for 3 minutes. The bath is changed and the step repeated once. Excess of liquid is tapped off and the slides are placed in 95% ethanol for 3 minutes. The bath is changed and the step repeated once. Excess of liquid is tapped off and the slides are placed in distilled water for a minimum of 30 seconds.

Epitope Retrieval

[0371] Staining jars are filled with diluted epitope retrieval solution and preheated in a water bath at 95°C. The deparaffinized sections are immersed into the preheated solution in the staining jars and incubated for 40 minutes at 95°C. The entire jar is removed from the water bath and allowed to cool down at room temperature for 20 minutes. The epitope retrieval solution is decanted, the sections are rinsed in distilled water and finally soaked in wash buffer for 5 minutes.

Peroxidase Blocking:

[0372] Excess of buffer is tapped off and the tissue section encircled with a DAKO pen. The specimen is covered with 3 drops (100 µl) Peroxidase-Blocking solution and incubated for 5 minutes. The slides are rinsed in distilled water and placed into a fresh washing buffer bath.

Antibody Incubation

[0373] Excess of liquid is tapped off and the specimen are covered with 3 drops (100 µl) of Anti-Her-2/neu reagent (Rabbit Anti-Human Her2 Protein in 0.05 mol/L Tris/HCl, 0.1 mol/L NaCl, 15 mmol/L pH7.2 NaN₃ containing stabilizing protein) or negative control reagent (= IGG fraction of normal rabbit serum at an equivalent protein concentration as the Her2 Ab). After 30 minutes of incubation the slide is rinsed in water and placed into a fresh water bath.

Visualization

[0374] Excess of liquid is tapped off and the specimen are covered with 3 drops (100 µl) of visualization reagent. After 30 minutes of incubation the slide is rinsed in water and placed into a fresh water bath. Excess of liquid is tapped off and the specimen are covered with 3 drops (100 µl) of Substrate-Chromogen solution (DAB) for 10 minutes. After rinsing the specimen with distilled water, photographs are taken with a conventional Olympus microscope to document the staining intensity and tumor regions within the specimen. Optionally a counterstain with hematoxylin was performed.

DNA extraction

[0375] The whole specimens or dissected subregions are transferred into a microcentrifuge tubes. Optionally a small amount (10µl) of preheated TaKaRa solution (DEXPAT™) is preheated and placed onto the specimen to facilitate sample transfer with a scalpel. 50 to 150 µl of TaKaRa solution were added to the samples depending on the size of the tissue sample selected. The sample are incubated at 100°C for 10 minutes in a block heater, followed by centrifugation at 12.000 rpm in a microcentrifuge. The supernatant is collected using a micropipet and placed in a separate microcentrifuge tube. If no deparaffinization step has been undertaken one has to be sure not to withdraw tissue debris and resin. Genomic DNA left in the pellet can be collected by adding resin-free TaKaRa buffer and an additional heating and centrifugation step. Samples are stored at -20°C.

[0376] Genomic DNA from different tumor cell lines (MCF-7, BT-20, BT-474, SKBR-3, AU-565, UACC-812, UACC-893, HCC-1008, HCC-2157, HCC-1954, HCC-2218, HCC-1937, HCC1599, SW480), or from lymphocytes is prepared with the QIAamp® DNA Mini Kits or the QIAamp® DNA Blood Mini Kits according to the manufacturers protocol. Usually between 1ng up to 1µg DNA is used per reaction.

Quantitative PCR

[0377] To measure the gene copy number of the genes within the patient samples the respective primer/probes (see table below) are prepared by mixing 25 µl of the 100 µM stock solution "Upper Primer", 25 µl of the 100 µM stock solution "Lower Primer" with 12,5 µl of the 100 µM stock solution Taq Man Probe (Quencher Tamra) and adjusted to 500 µl with aqua dest. For each reaction 1,25 µl DNA-Extract of the patient samples or 1,25 µl DNA from the cell lines were mixed with 8,75 µl nuclease-free water and added to one well of a 96 Well-Optical Reaction Plate (Applied Biosystems Part No. 4306737). 1,5 µl Primer/Probe mix, 12, µl Taq Man Universal-PCR Mix (2x) (Applied Biosystems Part No. 4318157) and 1 µl Water are then added. The 96 well plates are closed with 8 Caps/Strips (Applied Biosystems Part Number 4323032) and centrifuged for 3 minutes. Measurements of the PCR reaction are done according to the instructions of the manufacturer with a TaqMan 7900 HT from Applied Biosystems (No. 20114) under appropriate conditions (2 min. 50°C, 10 min. 95°C, 0.15min. 95°C, 1 min. 60°C; 40 cycles). Software SDS 2.0 from Applied Biosystems is used according to the respective instructions. CT-values are then further analyzed with appropriate software (Microsoft Excel™).

REFERENCES**[0378]**

5	<i>Patents cited</i>
	U.S. Pat. No. 4,843,155
	U.S. Pat. No. 5,262,31
	U.S. Pat. No. 4,683,202
10	U.S. Pat. No. 5,593,839
	U.S. Pat. No. 5,578,832
	U.S. Pat. No. 5,556,752
	U.S. Pat. No. 5,631,734
	U.S. Pat. No. 5,599,695
15	U.S. Pat. No. 4,683,195
	U.S. Pat. No. 5,498,531
	U.S. Pat. No. 5,714,331
	U.S. Pat. No. 5,641,673
20	U.S. Pat. No. 5,223,409
	U.S. Pat. No. 5,976,813
	U.S. Pat. No. 5,283,317
	U.S. Pat. No. 6,203,987
25	U.S. Pat. No. 6,379,895
	WO 97/29212
	WO 97/27317
	WO 95/22058
	WO 99/12826
30	WO 97/02357
	WO 94/13804
	WO 94/10300
	WO 97/14028
	WO 99/52708
35	EP 0 785 280
	EP 0 799 897
	EP 0 728 520
	EP 0 721 016
40	EP 0 321 201

GB2188638B

Publications cited

45

[0379]

- (1) Gusterson et al., Journal of Clinical Oncology 10, 1049-1056, 1992
- (2) Achuthan et al., Cancer Genet Cytogenet. 130:166-72, 2001
- 50 (3) Tomasetto et al., FEBS Lett. 373: 245-249, 1995
- (4) Pragnell et al., FEBS Lett. 291: 253-258, 1991
- (5) Nakamichi et al. 1986
- (6) Feo et al., Proc. Nat. Acad. Sci. 86: 6691-6695, 1989
- (7) Davies et al., Proc. Nat. Acad. Sci. 86: 6691-6695, 1989
- 55 (8) Lee et al., Molec. Endocr. 9: 243-254, 1995
- (9) Drane et al., Oncogene 15: 3013-3024, 1997
- (10) Zhu et al., J. Biol. Chem. 272: 25500-25506, 1997
- (11) Yuan et al., Proc. Nat. Acad. Sci. 95: 7939-7944, 1998

- (12) Zhu et al., Proc. Nat. Acad. Sci. 96: 10848-10853, 1999
- (13) Lee et al., Science 268: 836-844, 1995
- (14) McCormick et al., Molec. Cell. Biol. 16: 5792-5800, 1996
- (15) Tamimi et al., Genomics 40: 355-357, 1997
- 5 (16) Valle et al., *FEBS Lett.* 415: 163-168, 1997
- (17) Kaneda et al., J. Biol. Chem. 263: 7672-7677, 1988
- (18) Hoehe et al., Hum. Molec. Genet. 1:175-178, 1992
- (19) Yang-Feng et al., Abstract Cytogenet. Cell Genet. 40: 784, 1985
- (20) Coussens et al., Science 230: 1132-1139, 1985
- 10 (21) van de Vijver et al., New Eng. J. Med. 319: 1239-1245, 1988
- (22) Slamon et al., Science 244: 707-712, 1989
- (23) Fukushima et al., Res. Commun. 134: 477-483, 1986
- (24) Kaneko et al., Jpn. J. Cancer Res. 78: 16-19, 1987
- (25) Di Fiore et al., Science 237: 178-182, 1987
- 15 (26) Popescu et al., Genomics 4: 362-366, 1989
- (27) Qiu et al., Nature 393: 83-85, 1998
- (28) Yu et al., Molec. Cell 2: 581-591, 1998
- (29) Doherty et al., Proc. Nat. Acad. Sci. 96: 10869-10874, 1999
- (30) Slamon et al., New Eng. J. Med. 344: 783-792, 2001
- 20 (31) Margolis et al., J. Clin. Invest. 102: 821-827, 1998
- (32) Tanaka et al., J. Clin. Invest. 102: 821-827, 1998
- (33) Dong et al., J. Biol. Chem. 272: 29104-29112, 1997
- (34) Stein et al., EMBO J.13:1331-40, 1994
- (35) Nagata et al., Nature 319: 415-418, 1986
- 25 (36) Le Beau et al., Leukemia 1: 795-799, 1987,
- (37) Jansson et al., EMBO J. 2: 561-565, 1983
- (38) Thompson et al., Science 237:1610-1614, 1987
- (39) Nakai et al., Proc. Nat. Acad. Sci. 85: 2781-2785, 1988
- (40) Miyajima et al., Cell 57: 31-39, 1989
- 30 (41) Debuire et al., Science 224: 1456-1459, 1984
- (42) Petkovich et al., Nature 330: 444-450, 1987
- (43) Mattei et al., Hum. Genet. 80: 186-188, 1988.
- (44) Williams et al., Molec. Cell. Biol. 18: 2758-2767, 1998
- (45) Saha et al, Molec. Cell. Biol. 18: 2758-2767, 1998
- 35 (46) Yan et al., Proc. Nat. Acad. Sci. 95: 3603-3608, 1998
- (47) Singh et al., Nucleic Acids Res. 16: 3919-3929, 1988
- (48) Tsai-Pflugfelder et al., Proc. Nat. Acad. Sci. 85: 7177-7181, 1988
- (49) Chung et al., Proc. Nat. Acad. Sci. 86: 9431-9435, 1989
- (50) Lang et al., Gene 221: 255-266, 1998
- 40 (51) Keith et al., Genes Chromosomes Cancer 4: 169-175, 1992
- (52) Kingsmore et al., Mammalian Genome 4: 288-289, 1993
- (53) Watt et al., Biochem. J. 303: 681-695, 1994
- (54) Kiefer et al., J. Biol. Chem. 267: 12692-12699, 1992
- (55) Shimasaki et al., Molec. Endocr. 4: 1451-1458, 1990
- 45 (56) Zazzi et al., Genomics 49: 401-410, 1998
- (57) Bajalica et al., Hum. Genet. 89: 234-236, 1992
- (58) Tonin et al., Genomics 18: 414-417, 1993
- (59) Birkenbach et al., J. Virol. 67: 2209-2220, 1993
- (60) Wang et al., Proc. Nat. Acad. Sci. 95: 492-498, 1998
- 50 (61) Klochendler-Yeivin et al., Curr Opin Genet Dev 12:73-9,2002
- (62) Ring et al., Genomics 51:140-3, 1998
- (63) Darmon et al., Molec. Biol. Rep. 12: 277-283, 1987
- (64) Zhou et al., J. Biol. Chem. 263: 15584-15589, 1988
- (65) Korge et al., Proc. Nat. Acad. Sci. 89: 910-914, 1992
- 55 (66) Lessin et al., J. Invest. Derm. 91: 572-578, 1988
- (67) Romano et al., Cytogenet. Cell Genet. 58: 2009-2010, 1991.
- (68) Fuchs et al., Proc. Nat. Acad. Sci. 89: 6906-6910, 1992
- (69) Rogaev et al., Nature Genet. 5: 158-162, 1993

- (70) Liu et al., Curr. Eye Res. 12: 963-974, 1993
 (71) Nishida et al., Invest. Ophthalm. Vis. Sci. 37: 1800-1809, 1996
 (72) Nishida et al., Am. J. Hum. Genet. 61: 1268-1275, 1997
 (73) Meesmann and Wilke, 1939
 5 (74) Corden et al., Exp. Eye Res. 70: 41-49, 2000
 (75) van de Vijver et al., Mol Cell Biol 7, 2091-23, 1987
 (76) Offlerdinger et al., Biochem Biophys Res Comm 251, 907-13, 1988
 (77) Sambrook et al., MOLECULAR CLONING: A LABORATORY MANUAL, 2d ed., 1989
 (78) Ausubel et al., CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, John Wiley & Sons, New York, N.Y.,
 10 1989.
 (79) Tedder, T. F. et al., Proc. Natl. Acad. Sci. U.S.A. 85:208-212, 1988
 (80) Hedrick, S. M. et al., Nature 308:149-153, 1984
 (81) Lee, S. W. et al., Proc. Natl. Acad. Sci. U.S.A. 88:4225, 1984
 (82) Sarkar, PCR Methods Applic. 2, 318-322, 1993
 15 (83) Triglia et al., Nucleic Acids Res. 16, 81-86, 1988
 (84) Lagerstrom et al., PCR Methods Applic. 1, 111-119, 1991
 (85) Copeland & Jenkins, Trends in Genetics 7: 113-118, 1991
 (86) Cohen, et al., Nature 366: 698-701, 1993
 (87) Bonner et al., J. Mol. Biol. 81, 123 1973
 20 (88) Bolton and McCarthy, Proc. Natl. Acad. Sci. U.S.A. 48, 1390 1962
 (89) Plump et al., Cell 71: 343-353, 1992
 (90) Altschul et al., Bull. Math. Bio. 48:603, 1986,
 (91) Henikoff & Henikoff, Proc. Natl. Acad. Sci. USA 89:10915, 1992
 (92) Pearson & Lipman, Proc. Nat'l Acad. Sci. USA 85:2444, 1988
 25 (93) Pearson et al., Meth. Enzymol. 183:63, 1990
 (94) Needleman & Wunsch, J. Mol. Biol.48:444, 1970
 (95) Sellers, SIAM J. Appl. Math.26:787, 1974
 (96) Takamatsu, EMBO J. 6, 307-311, 1987
 (97) Coruzzi et al., EMBO J. 3, 1671-1680, 1984
 30 (98) Broglie et al., Science 224, 838-843, 1984
 (99) Winter et al., Results Probl. Cell Differ. 17, 85-105, 1991
 (100) Engelhard et al., Proc. Nat. Acad. Sci. 91, 3224-3227, 1994
 (101) Logan & Shenk, Proc. Natl. Acad. Sci. 81, 3655-3659, 1984
 (102) Scharf et al., Results Probl. Cell Differ. 20, 125-162, 1994
 35 (103) Freshney R.I., ed., ANIMAL CELL CULTURE , 1986
 (104) Wigler et al., Cell 11, 223-232, 1977
 (105) Lowy et al., Cell 22, 817-823, 1980
 (106) Wigler et al., Proc. Natl. Acad. Sci. 77, 3567-3570, 1980
 (107) Colbere-Garapin et al., J. Mol. Biol. 150, 114, 1981
 40 (108) Hartman & Mulligan, Proc. Natl. Acad. Sci. 85, 8047-8051, 1988
 (109) Rhodes et al., Methods Mol. Biol. 55, 121-131, 1995
 (110) Hampton et al., SEROLOGICAL METHODS: A LABORATORY MANUAL, APS Press, St. Paul, Minn., 1990
 (111) Maddox et al., J. Exp. Med. 158, 1211-1216, 1983
 (112) Porath et al., Prot. Exp. Purif. 3, 263-281, 1992
 45 (113) Kroll et al., DNA Cell Biol. 12, 441-453, 1993
 (114) Caruthers et al., Nucl. Acids Res. Symp. Ser. 215-223, 1980
 (115) Horn et al. Nucl. Acids Res. Symp. Ser. 225-232, 1980
 (116) Merrifield, J. Am. Chem. Soc. 85, 2149-2154, 1963
 (117) Roberge et al., Science 269, 202-204, 1995
 50 (118) Creighton, PROTEINS: STRUCTURES AND MOLECULAR PRINCIPLES, WH and Co., New York, N.Y., 1983
 (119) Cronin et al., Human Mutation 7:244, 1996
 (120) Landegran et al., Science 241:1077-1080, 1988
 (121) Nakazawa et al., PNAS 91:360-364, 1994
 (122) Abravaya et al., Nuc Acid Res 23:675-682, 1995
 55 (123) Guatelli, J.C. et al., Proc. Natl. Acad. Sci. USA 87:1874-1878, 1990
 (124) Kwoh, D.Y. et al., Proc. Natl. Acad. Sci. USA 86:1173-1177, 1989
 (125) Lizardi, P.M. et al., Bio/Technology 6:1197, 1988
 (126) Brown, Meth. Mol. Biol. 20, 18, 1994

- (127) Sonveaux, Meth. Mol. Biol. 26, 1-72, 1994
- (128) Uhlmann et al., Chem. Rev. 90, 543-583, 1990
- (129) Gee et al., in Huber & Carr, MOLECULAR AND IMMUNOLOGIC APPROACHES, Publishing Co., Mt. Kisco, N.Y., 1994
- 5 (130) Agrawal et al., Trends Biotechnol. 10, 152-158, 1992
- (131) Uhlmann et al., Tetrahedron. Lett. 215, 3539-3542, 1987
- (132) Cech, Science 236, 1532-1539, 1987
- (133) Cech, Ann. Rev. Biochem. 59, 543-568, 1990
- (134) Couture & Stinchcomb, Trends Genet. 12, 510-515, 1996
- 10 (135) Haseloff et al. Nature 334, 585-591, 1988
- (136) Kohler et al., Nature 256, 495-497, 1985
- (137) Kozbor et al., J. Immunol. Methods 81, 3142, 1985
- (138) Cote et al., Proc. Natl. Acad. Sci. 80, 2026-2030, 1983
- (139) Cole et al., Mol. Cell Biol. 62, 109-120, 1984
- 15 (140) Morrison et al., Proc. Natl. Acad. Sci. 81, 6851-6855, 1984
- (141) Neuberger et al., Nature 312, 604-608, 1984
- (142) Takeda et al., Nature 314, 452-454, 1985
- (143) Burton, Proc. Natl. Acad. Sci. 88, 11120-11123, 1991
- (144) Thirion et al., Eur. J. Cancer Prev. 5, 507-11, 1996
- 20 (145) Coloma & Morrison, Nat. Biotechnol. 15, 159-63, 1997
- (146) Mallender & Voss, J. Biol. Chem. 269, 199-206, 1994
- (147) Verhaar et al., Int. J. Cancer 61, 497-501, 1995
- (148) Nicholls et al., J. Immunol. Meth. 165, 81-91, 1993
- (149) Orlandi et al., Proc. Natl. Acad. Sci. 86, 3833-3837, 1989
- 25 (150) Winter et al., Nature 349, 293-299, 1991
- (151) Lam, Anticancer Drug Des. 12, 145, 1997
- (152) DeWitt et al., Proc. Natl. Acad. Sci. U.S.A. 90, 6909, 1993
- (153) Erb et al. Proc. Natl. Acad. Sci. U.S.A. 91, 11422, 1994
- (154) Zuckermann et al., J. Med. Chem. 37, 2678, 1994
- 30 (155) Cho et al., Science 261, 1303, 1993
- (156) Carell et al., Angew. Chem. Int. Ed. Engl. 33, 2059 & 2061, 1994
- (157) Gallop et al., J. Med. Chem. 37, 1233, 1994
- (158) Houghten, BioTechniques 13, 412-421, 1992
- (159) Lam, Nature 354, 8284, 1991
- 35 (160) Fodor, Nature 364, 555-556, 1993
- (161) Cull et al., Proc. Natl. Acad. Sci. U.S.A. 89, 1865-1869, 1992
- (162) Scott & Smith, Science 249, 386-390, 1990
- (163) Devlin, Science 249, 404-406, 1990
- (164) Cwirla et al., Proc. Natl. Acad. Sci. 97, 6378-6382, 1990
- 40 (165) Felici, J. Mol. Biol. 222, 301-310, 1991
- (166) Jayawickreme et al., Proc. Natl. Acad. Sci. U.S.A. 19, 1614-1618, 1994
- (167) Chelsky, Strategies for Screening Combinatorial Libraries 1995
- (168) Salmon et al., Molecular Diversity 2, 57-63, 1996
- (169) McConnell et al., Science 257, 1906-1912, 1992
- 45 (170) Sjolander & Urbaniczky, Anal. Chem. 63, 2338-2345, 1991
- (171) Szabo et al., Curr. Opin. Struct. Biol. 5, 699-705, 1995
- (172) Zervos et al., Cell 72, 223-232, 1993
- (173) Madura et al., J. Biol. Chem. 268, 12046-12054, 1993
- (174) Bartel et al., BioTechniques 14, 920-924, 1993
- 50 (175) Iwabuchi et al., Oncogene 8, 1693-1696, 1993
- (176) Findeis et al. Trends in Biotechnol. 11, 202-205, 1993
- (177) Chiou et al., GENE THERAPEUTICS: METHODS AND APPLICATIONS OF DIRECT GENE TRANSFER J. A. Wolf ed., 1994
- (178) Wu & Wu, J. Biol. Chem. 263, 621-24, 1988
- 55 (179) Wu et al., J. Biol. Chem. 269, 542-46, 1994
- (180) Zenke et al., Proc. Natl. Acad. Sci. U.S.A. 87, 3655-59, 1990
- (181) Wu et al., J. Biol. Chem. 266, 338-42, 1991
- (182) REMINGTON'S PHARMACEUTICAL SCIENCES Maack Publishing Co., Easton, Pa.

EP 1 365 034 A2

(183) Hille, Excitable Membranes, Sunderland, MA, Sinauer Associates, Inc.

Table 1

5	DNA SEQ ID NO:	Protein SEQ ID NO:	Genbank ID	Unigene_v133_ID	Locus Link ID	Gene Name
	1	27	NM_006148.1	75080	3927	LASP1
	2	28	NM_000723.1	635	782	CACNB1
10	3	29	NM_000981.1	252723	6143	RPL19RPL19
	4	30	Y13467	15589	5469	PPARGBP
	5	31	NM_016507.1	123073		CrkRS
15	6	32	AB021742.1	322431	4761	NEUROD2
	7	33	NM_006804.1	77628	10948	MLN64
	8	34	NM_003673.1	111110	8557	TELETHONIN
	9	35	NM_002686.1 I	1892	5409	PNMT
20	10	36	X03363.1	323910	2064	ERBB2
	11	37	AB008790.1	86859	2886	GRB7
	12	38	NM_002809.1	9736	5709	PSMD3
25	13	39	NM_000759.1	2233	1440	GCSFG
	14	40	AI023317	23106	9862	KIAA0130
	15	41	X55005		7067	c-erbA-1
	16	42	X72631	211606	9572	NRID1
30	17	43	NM_007359.1	83422	22794	MLN51
	18	44	U77949.1	69563	990	CDC6
	19	45	U41742.1		5914	RARA
35	20	46	NM_001067.1	156346	7153	TOP2A
	21	47	NM_001552.1	1516		IGFBP4
	22	48	NM_001838.1	1652		CCR7 EBI1
40	23	49	NM_003079.1	332848	6605	SMARCE1 BAF57
	24	50	X14487	99936	3858	KRT10
	25	51	NM_000223.1	66739		KRT12
45	26	52	NM_002279.2	32950	3884	hHKa3-II
	53	76	NM_005937	349196	4302	MLLT6
	54	77	XM_008147	184669	7703	ZNF144
	55	78	NM_138687	432736	8396	PIP5K2B
50	56	79	NM_020405	125036	57125	TEM7
	57	80	XM_012694	258579	22806	ZNFN1A3
	58	81	XM_085731	13996	147179	WIRE
55	59	82	NM_002795	82793	5691	PSMB3
	60	83	NM_033419	91668	93210	MGC9753 Variant a

EP 1 365 034 A2

Table 1 (continued)

DNA SEQ ID NO:	Protein SEQ ID NO:	Genbank ID	Unigene_v133_ID	Locus Link ID	Gene Name
61	84				MGC9753 Variant c
62	85				MGC9753 Variant d
63	86				MGC9753 Variant e
64	87				MGC9753 Variant g
65	88				MGC9753 Variant h
66	89				MGC9753 Variant i
67	90	AF395708	374824	94103	ORMDL3
68	91	NM_032875	194498	84961	MGC15482
69	92	NM_032192	286192	84152	PPP1R1B
70	93	NM_032339	333526	84299	MGC14832
71	94	NM_057555	12101	51242	LOC51242
72	95	NM_017748	8928	54883	FLJ20291
73	96	NM_018530	19054	55876	Pro2521
74	97	NM_016339	118562	51195	Link-GEFII
75	98	NM_032865	294022	84951	CTEN

Table 2

DNA SEQ ID NO:	Gene description
1	Member of a subfamily of LIM proteins that contains a LIM domain and an SH3 (Src homology region 3) domain
2	Beta 1 subunit of a voltage-dependent calcium channel (dihydropyridine receptor), involved in coupling of excitation and contraction in muscle, also acts as a calcium channel in various other tissues
3	Ribosomal protein L19, component of the large 60S ribosomal subunit
4	Protein with similarity to nuclear receptor-interacting proteins; binds and co-activates the nuclear receptors PPARalpha (PPARA), RARalpha (RARA), RXR, TRbeta1, and VDR
5	we26e0 CDC2-related protein kinase 7
6	Neurogenic differentiation, a basic-helix-loop-helix transcription factor that mediates neuronal differentiation
7	Protein that is overexpressed in malignant tissues, contains a putative trans-membrane region and a StAR Homology Domain (SHD), may function in steroidogenesis and contribute to tumor progression
8	Telethonin, a sarcomeric protein specifically expressed in skeletal and heart muscle, caps titin (TTN) and is important for structural integrity of the sarcomere

Table 2 (continued)

DNA SEQ ID NO:	Gene description
5 9	Phenylethanolamine N-methyltransferase, acts in catecholamine biosynthesis to convert norepinephrine to epinephrine
10 10	Tyrosine kinase receptor that has similarity to the EGF receptor, a critical component of IL-6 signaling through the MAP kinase pathway, overexpression associated with prostate, ovary and breast cancer
10 11	Growth factor receptor-bound protein, an SH2 domain-containing protein that has isoforms which may have a role in cell invasion and metastatic progression of esophageal carcinomas
12	Non-ATPase subunit of the 26S proteasome (prosome, macropain)
15 13	Granulocyte colony stimulating factor, a glycoprotein that regulates growth, differentiation, and survival of neutrophilic granulocytes
14	Member of the Vitamin D Receptor Interacting Protein co-activator complex, has strong similarity to thyroid hormone receptor-associated protein (murine Trap 100) which function as a transcriptional coregulator
20 15	Thyroid hormone receptor alpha, a high affinity receptor for thyroid hormone that activates transcription; homologous to avian erythroblastic leukemia virus oncogene
16	encoding Rev-ErbAalp nuclear receptor subfamily 1, group D, member 1
17	Protein that is overexpressed in breast carcinomas
25 18	Protein which interacts with the DNA replication proteins PCNA and Orc1, translocates from the nucleus following onset of S phase; <i>S. cerevisiae</i> homolog Cdc6p is required for initiation of S phase
30 19	Retinoic acid receptor alpha, binds retinoic acid and stimulates transcription in a ligand-dependent manner
20	DNA topoisomerase II alpha, member of a family of proteins that relieves torsional stress created by DNA replication, transcription, and cell division;
35 21	Insulin-like growth factor binding protein, the major IGFBP of osteoblast-like cells, binds IGF1 and IGF2 and inhibits their effects on promoting DNA and glycogen synthesis in osteoblastic cells
22	HUMEBI103 G protein-coupled receptor (EBI 1) gene exon 3 chemokine (C-C motif) receptor 7 G protein-coupled receptor
40 23	Protein with an HMG 1/2 DNA-binding domain that is subunit of the SNF/SWI complex associated with the nuclear matrix and implicated in regulation of transcription by affecting chromatin structure
45 24	Keratin 10, a type I keratin that is a component of intermediate filaments and is expressed in terminally differentiated epidermal cells; mutation of the corresponding gene causes epidermolytic hyperkeratosis
25	Keratin 12, a component of intermediate filaments in corneal epithelial cells; mutation of the corresponding gene causes Meesmann corneal dystrophy
50 26	Hair keratin 3B, a type I keratin that is a member of a family of structural proteins that form intermediate filaments
53	MLLT6 Myeloid/lymphoid or mixed-lineage leukemia (trithorax homolog, <i>Drosophila</i>); translocated to, 6
54	zinc finger protein 144 (Mel-18)
55 55	phosphatidylinositol-4-phosphate 5-kinase type II beta isoform a
56	tumor endothelial marker 7 precursor

Table 2 (continued)

DNA SEQ ID NO:	Gene description
57	zinc finger protein, subfamily 1A, 3
58	WASP-binding protein putative cr16 and wip like protein similar to Wiskott-Aldrich syndrome protein
59	proteasome (prosome, macropain) subunit, beta type, 3
60	Predicted
67	ORM1-like 3 (<i>S. cerevisiae</i>)
68	F-box domain A Receptor for Ubiquitination Targets
69	protein phosphatase 1, regulatory (inhibitor) subunit 1B (dopamine and cAMP regulated phosphoprotein, DARPP-32)
70	Predicted Protein
71	Predicted Protein
72	Predicted Protein
73	Predicted Protein
74	Link-GEFII: Link guanine nucleotide exchange factor II
75	C-terminal tensin-like

Table 3

DNA SEQ ID NO:	Gene function	Subcellular localization
1	SH3/SH2 adapter protein	-
3	voltage-gated calcium channel membrane fraction Channel [passive transporter]	Plasma membrane
4	RNA binding structural protein of ribosome protein biosynthesis	Cytoplasm
5	transcription co-activator nucleus Pol II transcription	Nucleus
6	-	-
7	transcription factor transcription regulation from Pol II promoter neurogenesis	-
8	mitochondrial transport steroid and lipid metabolism	Cytoplasm
9	structural protein of muscle sarcomere alignment	Cytoplasm
10	phenylethanolamine N-methyltransferase Transferase	-
11	Neu/ErbB-2 receptor receptor signaling protein tyrosine kinase	Plasma membrane
12	SH3/SH2 adapter protein EGF receptor signaling pathway	Cytoplasm
13	26S proteasome Protein degradation Proteasome subunit	Cytoplasm
14	developmental processes positive control of cell proliferation	Extracellular space
15	fatty acid omega-hydroxylase fatty acid omega-hydroxylase	-
16	DNA-binding protein Transcription factor	Nucleus
17	steroid hormone receptor transcription co-repressor	Nucleus
18	-	-
19	nucleotide binding cell cycle regulator DNA replication checkpoint regulation of CDK activity	nucleus
20	retinoic acid receptor transcription co-activator transcription factor	nucleus
21	DNA binding DNA topoisomerase (ATP-hydrolyzing)	nucleus
22	skeletal development DNA metabolism signal transduction cell proliferation	plasma membrane
23	chromatin binding transcription co-activator nucleosome disassembly transcription	nucleus nuclear chromosome
24	Cell structure Cytoskeletal Epidermal Development and Maintenance	cytoplasm
25	structural protein vision cell shape and cell size control intermediate filament	cytoplasm
26	cell shape and cell size control Cell structure	cytoplasm
53	leucine-zipper containing fusion	-

Table 3 (continued)

DNA SEQ ID NO:	Gene function	Subcellular localization
54		-
55	Tumor endothelial marker 7 precursor; may be involved in angiogenesis	-
56	Aiolos; DNA binding protein that may be a transcription factor; has strong similarity to murine Znf1a3, contains zinc finger domain	-
57	The WASP-binding protein WIRE has a role in the regulation of the actin filament system downstream of the platelet-derived growth factor receptor	-
58		-
59		-
60		-
67		-
68		-
69	Midbrain dopaminergic neurons play a critical role in multiple brain functions, and abnormal signaling through dopaminergic pathways has been implicated in several major neurologic and psychiatric disorders. One well-studied target for the actions of dopamine is DARPP32.	-
70		-
71		-
72		-
73		-
74	Brain-specific guanine nucleotide exchange factor; activates the ERK/MAP kinase cascade plus R-Ras and H-ras; activates targets through a Ca2+- and diacylglycerol-sensitive mechanism; active protein associates with membranes	-
75	C-terminal tensin-like <u>Phosphotyrosine-binding domain, phosphotyrosine-interaction (PTI) domain</u>	-

EP 1 365 034 A2

Table 4

DNA SEQ ID NO:	Protein SEQ ID NO:	Gene Name	DBSNPID	Type	Codon	AA-Seq
9	34	ERBB2	rs2230698	coding-synon	TCA TCG	S S
9	34	ERBB2	rs2230700	noncoding		
9	34	ERBB2	rs1058808	coding-nonsynon	CCC GCC	P A
9	34	ERBB2	rs1801200	noncoding		
9	34	ERBB2	rs903506	noncoding		
9	34	ERBB2	rs2313170	noncoding		
9	34	ERBB2	rs1136201	coding-nonsynon	ATC GTC	I V
9	34	ERBB2	rs2934968	noncoding		
9	34	ERBB2	rs2172826	noncoding		
9	34	ERBB2	rs1810132	coding-nonsynon	ATC GTC	I V
9	34	ERBB2	rs1801201	noncoding		
14	39	c-erbA-1	rs2230702	coding-synon	TCC TCT	S S
14	39	c-erbA-1	rs2230701	coding-synon	GCC GCT	A A
14	39	c-erbA-1	rs1126503	coding-nonsynon	ACC AGC	T S
14	39	c-erbA-1	rs3471	noncoding		
19	44	TOP2A	rs13695	noncoding		
19	44	TOP2A	rs471692	noncoding		
19	44	TOP2A	rs558068	noncoding		
19	44	TOP2A	rs1064288	noncoding		
19	44	TOP2A	rs1061692	coding-synon	GGA GGG	G G
19	44	TOP2A	rs520630	noncoding		
19	44	TOP2A	rs782774	coding-nonsynon	AAT ATT ATT TTT	N I I F
19	44	TOP2A	rs565121	noncoding		
19	44	TOP2A	rs2586112	noncoding		
19	44	TOP2A	rs532299	coding-nonsynon	TTT GTT	F V
19	44	TOP2A	rs2732786	noncoding		
19	44	TOP2A	rs1804539	noncoding		
19	44	TOP2A	rs1804538	noncoding		
19	44	TOP2A	rs1804537	noncoding		
19	44	TOP2A	rs1141364	coding-synon	AAA AAG	K K
23	48	KRT10	rs12231	noncoding		
23	48	KRT10	rs1132259	coding-nonsynon	CAT CGT	H R
23	48	KRT10	rs1132257	coding-synon	CTG TTG	L L
23	48	KRT10	rs1132256	coding-synon	GCC GCT	A A
23	48	KRT10	rs1132255	coding-synon	CTG TTG	L L
23	48	KRT10	rs1132254	coding-synon	GGC GGT	G G
23	48	KRT10	rs1132252	coding-synon	TTC TTT	F F

Table 4 (continued)

DNA SEQ ID NO:	Protein SEQ ID NO:	Gene Name	DBSN ^{PID}	Type	Codon	AA-Seq
23	48	KRT10	rs1132268	coding-nonsynon	CAG GAG	Q E
23	48	KRT10	rs1132258	coding-nonsynon	CGG TGG	R W

Table 5

PRIMER	SEQUENCE
CACNB1	FAM 5' CCATATATAAAACCACTGCTCCTGTCCTTTGTGGCT 3'TAMRA
CACNB1FOR	5' CCCCCATCTGCTGCTATATTGTC 3'
CACNB1REV	5' TGCCTACGCTGACGACTATGTG 3'
CDC6	FAM 5' TTTGGTTTCTACAACTGTTGCTAT 3'TAMRA
CDC6 FOR	5' GGGCTCCACACACCAAGATG 3'
CDC6 REV	5' ACGTCTGAGCACCCCTCTACA 3'
EBI1-1	FAM 5' TGTACACAGGACTGAAAACCTCTCCTCATGT 3'TAMRA
EBI1-1 FOR	5' CCCAAGGCCACGAGCTT 3'
EBI1-1 REV	5' TGTGCTCTCTTAACGAATCGAA 3'
EBI1-2	FAM 5' CTGGTCAACAACTCTCTGAACCCCTCC 3'TAMRA
EBI1-2 FOR	5' TGGTGAGGAAAAGCGGACAT 3'
EBI1-2 REV	5' CTGGCTTGAGGACAGTGAAG 3'
GCSF	FAM 5' CCAAGCCCTCCCCATCCCATGTAT 3'TAMRA
GCSF FOR	5' GAGGTGTCGTACCGCGTTCTA 3'
GCSF REV	5' CCGTCTGCTCTTCCCTGTCT 3'
GRB7	FAM 5' CCAGACCCGCTTCACTGACCTGC 3'TAMRA
GRB7 FOR	5' CGCCTGTACTTCAGCATGGA 3'
GRB7 REV	5' GCGGTTCAAGTGGTGAA 3'
HKA3	FAM 5' ACCCCGAGGCATCACCACAAATCAT 3'TAMRA
HKA3 FOR	5' AGTCTGCCCTCTCTGACAAACCAT 3'
HKA3 REV	5' TAGGCTCAGAGTCAGACCCCAAC 3'
MLN50	FAM 5' CCCCTCGTGGCTTGTGCTCGG 3'TAMRA
MLN50 FOR	5' AAGCCGCCAGTTCATCTTTT 3'
MLN50 REV	5' CTTGTGGTTCAAGTCAAAATGTTTTCAG 3'
MLN64-1	FAM 5' TCTGCTGCGCTCTCGTCGGT 3'TAMRA
MLN64-1 FOR	5' GGGCTGGGCACCTGACTT 3'

Table 5 (continued)

PRIMER	SEQUENCE
MLN64-1REV	5' CCCAACAAAGGTCCCAGACT 3'
MLN64-2	FAM 5' CGGCGCATTTAGCGGCG 3'TAMRA
MLN64-2 FOR	5' CCCAAGGACTTCGTGAATG 3'
MLN64-2REV	5' GGCATCCCTGATGACAAAGTA 3'
PPARBP	FAM 5' AGCACCAACTGTGAACCAGGTACAATGGC 3'TAMRA
PPARBP FOR	5' GAGGGAGGCTCTGCTTGG 3'
PPARBP REV	5' TCACAACTAGCGGGTGAGGAG 3'
PSMD3	FAM 5' TGCAGAGGAACGGCGTGAGCG 3'TAMRA
PSMD3 FOR	5' TGAGGTTTCCTCCCAAATCGTA 3'
PSMD3 REV	5' CAGCTCAAGGGAAGCTGTCATC 3'
RAR	FAM 5' CCCCCACATGTTCCCCAAGATGCT 3'TAMRA
RAR FOR	5' GGAGGCGCTAAAGGTCTACGT 3'
RAR REV	5' TGATGCTTCGCAGGTCAGTAA 3'
RPL23A	FAM 5' CTCCTGCCCTCCTAAAGCTGAAGCC 3'TAMRA
RPL23A FOR	5' GGACGCGTGGGCTTTTC 3'
RPL23A REV	5' TGTGGCTGTGGACACCTTTC 3'
RPL19	FAM 5' CCACAAGCTGAAGGCAGACAAGGCC 3'TAMRA
RPL19 FOR	5' GCGGATTCATGGAACACA 3'
RPL19 REV	5' GGTACGCCAGGAGCTTCTTG 3'
NEUROD2	FAM 5' ACCACCTTGCGCAGGTTGTCCAG 3'TAMRA
NEUROD2 FOR	5' CGCATGCACGACCTGAAC 3'
NEUROD2 REV	5' GTCTCGATCTTGACAGCTTCTG 3'
TELE TELETHONIN	FAM 5' AACTGTCCACACGGCCCGGAGG 3'TAMRA
TELE TELETHONIN FOR	5' CTGGGCAGAAATGGAAGGATCT 3'
TELE TELETHONIN REV	5' GGGACTCTAGCAGACCCACACT 3'
PENT PNMT	FAM 5' CACCCACCTGGATTCCCTGTTT 3'TAMRA
PENT PNMT FOR	5' CCTTCAGACAGGCGTAGATGATG 3'
PENT PNMT REV	5' GGGTATTATTCTTTATTAGTGCCACTT 3'

Table 5 (continued)

PRIMER	SEQUENCE
HER2/NEU;ERBB2	FAM 5' TTCCCTAAGGCTTTTCAGTACCCAGGATCTG 3'TAMRA
HER2/NEU;ERBB FOR	5' CCAGCTTGGCCCTTTCCT 3'
HER2/NEU;ERBB REV	5' GAATGGGTCGCTTTTGTCTTAG 3'
KIA0130	FAM 5' TCACGGACCTCAGCCTGCCCT 3'TAMRA
KIA0130 FOR	5' TGGTGAAGGTGTCAGCCATGT 3'
KIA0130 REV	5' TCAGAGTGCAGCAATGGCTTT 3'
THRA	FAM 5' ACCTCCTTCCCCAGCTCCCC 3'TAMRA
THRA FOR	5' GGCAACATCTTACTTGTCTTTGA 3'
THRA REV	5' CCAAGGAAGCAGACAACTATTC 3'
MLN51	FAM 5' TCCTCCCTATCCATGGCACTAAACCCTTC 3'TAMRA
MLN51 FOR	5' TGGGCAAGGGCTCCTATCT 3'
MLN51 REV	5' GTTACCCCTGGCAGACGTATG 3'
TOP2A	FAM 5' TGCCTCTGAGTCTGAATCTCCCAAAGAGAGA 3'TAMRA
TOP2A FOR	5' GAGTAGTTATGTGATTATTTTCAGCTCTTGAC 3'
TOP2A REV	5' TCAAATGTTGTCCCGAGTCT 3'
KRT10	FAM 5' CAGAAATTCGGAAGACAGAACTATTGTCATGCCT 3'TAMRA
KRT10 FOR	5' GATTAGTAACCCATAGCAGTTGAAGGT 3'
KRT10 REV	5' ATTTACTGACGGTGGTCTGAACATAC 3'
K12 KRT12	FAM 5' TGACAGACTCCAAATCACAAAGCACAGTCAAC 3'TAMRA
K12 KRT12 FOR	5' TGATGGTTTGGAGGAAAGTTATTT 3'
K12 KRT12 REV	5' TTTGGTTGGTCTTTAGAGGAATC 3'
NR1D1	FAM 5' TGCCAAACCATGCATCAGGTAGCCC 3'TAMRA
NR1D1 FOR	5' CAGCTCACCTGGCAACTCA 3'
NR1D1 REV	5' CCTGATTTTCCCGAGGATGT 3'
HSEB1	FAM 5' CGCCGCTCCCGTTCTGCT 3'TAMRA
HSEB1 FOR	5' TGGCCAAGCGTAAGCTGAT 3'
HSEB1 REV	5' GCTGCAGTGATCGGATCATCT 3'
MLLT6	FAM 5' CACCATGGAGCCCATCGTGCTG 3'TAMRA
MLLT6 FOR	5' ATCCCCGAGGTGCAATTG 3'

Table 5 (continued)

PRIMER	SEQUENCE
MLT6 REV	5' AGCGATCATGAGGCACGTACT 3'
ZNF144	FAM 5' CCTGCCAGAGATAGGAGACCCAGACAGCT 3'TAMRA
ZNF144 FOR	5' ATCCCCCTGAGCCTTTTCA 3'
ZNF144 REV	5' CAGCCTCTGGTCCCACCAT 3'
PIP5K2B	FAM 5' TGATCATCAATTCCAAACCTCTCTCCCGAA 3'TAMRA
PIP5K2B FOR	5' CCCCATGGTGTCCGAAAC 3'
PIP5K2B REV	5' TGCCAGGAGCCTCCATACC 3'
TEM7	FAM 5' CAGCCTTCTAAAAACACAATGTATTTCATGT 3'TAMRA
TEM7 FOR	5' CCTGAACCTTAATGGTAGAATTCAAAGATC 3'
TEM7 REV	5' TATTAACACTGAGAATCCATGCAGAGA 3'
ZNFN1A3	FAM 5' TATCTGTCTCAGGGATTGCTCCTATGTATTTCAGC 3'TAMRA
ZNFN1A3 FOR	5' CACAGAGCCCTGCTGAAGTG 3'
ZNFN1A3 REV	5' GCGAGGTCATTGGTTTITAGAAA 3'
WIRE	FAM 5' CTGTGATCCGAAATGGTGCCAG 3'TAMRA
WIRE FOR	5' CCGTCTCCACATCCAAACCT 3'
WIRE REV	5' ACCCATGCATTCCGGTATGGT 3'
PSMB3	FAM 5' AGTGGCACCTGCGCCGAAACAA 3'TAMRA
PSMB3 FOR	5' CCCCATGGTGACTGATGACTT 3'
PSMB3 REV	5' CCAGAGGGACTCACACATTCC 3'
MGC9753	FAM 5' CCAGAAACTTTCCATCCCAAAGGCAGTCT 3'TAMRA
MGC9753 FOR	5' CTGCCCCACAGGAATAGAAATG 3'
MGC9753 REV	5' AAAAATCCAGTCTGCTTCAACCA 3'
ORMDL3	FAM 5' AGCTGCCCCAGCTCCACGGA 3'TAMRA
ORMDL3 FOR	5' TCCCTGATGAGCGTGCTTATC 3'
ORMDL3 REV	5' TCTCAGTACTTATTGATTCCTCAAAATCC 3'
MGC15482	FAM 5' TCCAGTGAAGCAACCCAGTGTTTC 3'TAMRA
MGC15482 FOR	5' CACTTCTAGAGCTACCGTGAGTCT 3'
MGC15482 REV	5' CCCTCACTTTGTAACCCCTTGCT 3'
PPP1R1B	FAM 5' CAGCGTGGCGCAACAACCCA 3'TAMRA

Table 5 (continued)

PRIMER	SEQUENCE
PPP1R1B FOR	5' GGGATTGTTTCGCCACACATA 3'
PPP1R1B REV	5' CCGATGTTAAGGCCCATAGC 3'
MGC14832	FAM 5' TAAATGTCCGGCCCAACATGAGTTCCC 3'TAMRA
MGC14832 FOR	5' CGCAGTGCCTGGCACAT 3'
MGC14832 REV	5' GACACCCCTGACCTATGGA 3'
LOC51242	FAM 5' CAGTGACCTCTCCCGTTCCCTTGG 3'TAMRA
LOC51242 FOR	5' TGGTCCCTGTGTCTCTTTC 3'
LOC51242 REV	5' AGGGTCAGGAGGGAGAAAAC 3'
FLJ20291	FAM 5' CCAGTGCCCAACCCGTTAAAGAGTCAA 3'TAMRA
FLJ20291 FOR	5' TTGTGGGACACTCAGTAACTTTGG 3'
FLJ20291 REV	5' ACAAGCACTCCCAACCGAGAT 3'
PRO2521	FAM 5' AGTCTGTCTCACTGCCATCGCCA 3'TAMRA
PRO2521 FOR	5' AAGCCTCTGGGTTTTCCTTT 3'
PRO2521 REV	5' CCCACTGGTGACAGGATGGT 3'
Link-GEFII	FAM 5' CATCTGACATCTTCCCGTGGAG 3'TAMRA
Link-GEFII FOR	5' CTTTGACGATGTCTCAACCA 3'
Link-GEFII REV	5' TTTCCCGTGGAGCAGGAA 3'
CTEN	FAM 5' CCGCCGCCCTAATATGCAACATTAGGG 3'TAMRA
CTEN FOR	5' CGAGTATTCCAAAGCTGGTATCG 3'
CTEN REV	5' ATCACAGAGAGATGGCCCTTATCT 3'

Table 6

No.	ID	forward	reverse	PCR size (bp)	GB ID
1	D17S946	ACAGTCTATCAAGCAGAAAAATCCT	TGCCGTGCCAGAGAGA	128-142	Z24029
2	D17S1181	GACAAACAGAGCGAGACTGCC	GCCAGCCTGTCACTTATTC	122	-
3	D17S2026	TGGTCATTGACACGAA	CAGCATTTGGATGCAATCC	171-318	G05498 X53777
4	D17S838	CTCCAGAAATCCAGACCATGA	AGGACAGTGTAGCCCTTC	71-103	Z51080
5	D17S250	GGAAGAAATCAAAATAGACAAAT	GCTGGCCATATATATATTTAAACC	151	-
6	D17S1818	CATAGGTATGTTTCAAGAAATGTGA	TGCCTACTGGAAACCAGA	119-151	Z52895
7	D17S614	AAGGGAAAGGGCTTTCAAAGCT	NGGAGGTTGCAGTGAGCCAAAGAT	136	L29873
8	D17S2019	CAAAAGCTTATGATGCTCAAACC	TTGTTTCCCTTTGACTTTCGA	151-152	G07286 Z39013
9	D17S608	TAGGTTACCTCTCATTTTCTTCAG	GTCGGGTCTTTATGGNGCTTGTG	136	L29870
10	D17S1655	CGGACCAGAGTGTCCATGG	GCATACAGCACCCCTCTACCT	240	-
11	D17S2147	AGGGAGAAATAAAATCTGTGG	CAGGAGTGAGACACTCTCCATG	138	G15195
12	D17S754	TGGATTCACTGACTCAGCCTGC	GCGTGTCTGTCTCCATGTGTGC	145	-
13	D17S1814	TCCCAATGACGGTGATG	CTGGAGGTTGGCTTGTGGAT	150-166	Z52854
14	D17S2007	GGTCCCACGAATTTGCTG	CCACCCAGAAAAACAGGAGA	102-103	G07073 X03438
15	D17S1246	TCGATCTCCTGACCTTGTA	TTGTCAACCCCATTGCCTTTC	115	-
16	D17S1979	CCTTGGATAGATTCAGCTCCC	CTTGTCCCTTCTCAATCCTCC	199	G11172 X55068
17	D17S1984	TTAAGCAAGGTTTAAATTAAGCTGC	GATTACAGTGCTCCCTCTCCC	134	G14779 T50487
18	D17S1984	GGTTTAAATTAAGCTGCATGGC	GATTACAGTGCTCCCTCTCCC	126	G11580 T50487
19	D17S1867	AGTTTGACACTGAGGCTTTG	TTTAGACTTGGTAACTGCCG	94	Z51301
20	D17S1788	TGCAGATGCCTAAGAACTTTTCAG	GCCATGATCTCCCAAAGCC	156-168	Z52160
21	D17S1836	TCGAGGTTATGGTGAGCC	AAACTGTGTGTCAAAAGGATACT	167-173	Z53182
22	D17S1787	GCTGATCTGAAGCCAAATGA	TACATGAAGGCATGGTCTG	239-251	Z52130
23	D17S1660	CTAATATAATCCTGGGCACATGG	GCTGCGGACCAGACAGAT	201	G06069
24	D17S2154	GATAAAACAAGCACTGGCTCC	CCCACGGCTTCTTGATCTA	137	G15440
25	D17S1955	TGTAATGTAAGCCCCCATGAGG	CACTCAACTCAACAGTCTAAAGGTG	180	G11900
26	D17S2098	GTGAGTTCAAGCATAGTAATTATCC	ATTCAGCCTCAGTTCACCTGCTTC	181	G13994
27	D17S518	GATCCAGTGGAGACTCAGAG	TAGTCTCTGGACACCCAGA	88 - 100	X60690
28	D17S1851	ATTCTGTAGTGTACCCCTGTTGAG	ACTGACTGGCCACTGC	237 - 253	Z53675
29	D11S4358	TCGAGAAAGGACAAAATCACC	GAACAGGGTTAGTCCATTCC	58	-

Table 6 (continued)

No.	ID	forward	reverse	PCR size (bp)	GB ID
30	D17S964	GTTCTTTCCCTCTTGTTGGG	AGTCAGCTGAGATTGTGCC	224	L36695
31	D19S1091	CAAGCCAAGACATCCCAGTT	CCCCACACACAGCTCATATG	238	G14589
32	D17S1179	TTTCTCTCTCATTCATTTGGG	GCAACAGAGGGAGACTCCAA	113 - 125	-
33	D10S2160	TCCCATCCCGTAAGACCTC	TATGGAGTACCTACTCTATGCCAGG	349	G06592
34	D17S1230	ATTCAAAGCTGGATCCCCTT	AGCTGTGACAAATGCCCTGTA	108	L32949
35	D17S1338	TCACCTGAGATTGGGAGACC	AAATGGGGCAGGAATGG	178 - 200	-
36	D17S2011	TCACCTGCTCCTCCAAGCCAG	AAACACCACACTCTCCCCTG	115	G07143
37	D17S1237	TTCTTGGGCTTCCCGTAGCC	GGGGCAGACGACTTCTCCTT	186	L32947
38	D17S2038	GGGATACAAACCTTTAAAGTTCC	ATTACACCTAATGAGGATTCTTCTTT	228	G6219
39	D17S2091	GCTGAAATAGCCATCTTGAGCTAC	TCCGCATCCTTTTAAAGAGGCAC	157	G13941
40	D17S649	CTTTCACCTCTTTCAGCTGAAGAGG	TGACGTGCTATTTCCTGTTTGTCT	146	L36685
41	D17S1190	GTTTGTGCTATGCCCTGC	CAACACACTACCCCAGGA	122	L18197
42	M87506	ACTCCTCATCTGTAGGGTCT	GAGTCCGCTACCTGAGTGCT	102 - 120	m87506

5 ggcacagctt gtgattccta cccctcttcc agcttctttt gccaaactgaa gccttcttct 1020
 gccacttctg cgggctccct cctctggcag gcttcccccg tgatcgactt cttgggttttc 1080
 tctctggatg gaacgggtat gggcctctct gggggaggca gggctggaat gggagacctg 1140
 ttggcctgtg ggccctacct gccctctctg tctctccctc cacatcctcc tgcccagcto 1200
 ctcacatacc cacacattcc agggctgggg tgagcctgac tgccaggacc ccaggtcagg 1260
 ggctccctac attccccaga gtgggatoca cttcttggtt cctgggatgg cgtgggggac 1320
 tctgccgctg tgtagggacc agtgggatgg gctctacctc tctttctcaa agagggggct 1380
 ctgcccacct ggggtctctc tccctaccto cctcctcagg ggcaacaaca ggagaatggg 1440
 10 gttcctgtct tggggcgaat tcatccctc cccgcgcgtt ccttcgcaca ctgtgatttt 1500
 gccctcctgc ccacgcagac ctgcagcggg caaagagctc ccgaggaagc acagcttggg 1560
 tcaggttctt gcccttctta attttaggga cagctaccgg aaggagggga acaaggagt 1620
 ctcttccgca gcccttttcc ccaagccccc cccagctctc cagggaccct tgccctgcctc 1680
 ctaggctgga agccatggtc ccgaagtgtg gggcaagggt gcctcaggac ctttttggtct 1740
 15 tcagcctccc tcagccccc ggatctgggt taggtggccg cctcctccctg ctccctcatgg 1800
 gaagatgtct cagagccttc catgacctcc cctccccagc ccaatgocaa gtggactgg 1860
 agctgcacaa agtcagcagg gacctataa tctccaagac ctggtgtgag gaggcaggag 1920
 cagtgtatgt tgacacgcaa gtgtgtgagt gtgagtgtga gagatggggc 1980
 gggggtgtgt ctgtaggtgt ctctgggcct gtgtgtgggt ggggttatgt gagggtatga 2040
 agagctgtct tcccttgaga gtttctcctc aaccacagt gagaggggag ggctcctggg 2100
 20 gcagagaagt tcttaggtt ttctttggaa tgaaattoct ccttcccccc atctctgagt 2160
 ggaggaagcc caccaatctg ccttttgagc tgtgtcaggg tggaaggtaa gaggttgggt 2220
 tggagtggg gctgccatag ggtctgcagc ctgctggggc taagcgggtg aggaaggctc 2280
 tgtcactoca ggcatatgtt tccccatctc tgtctggggc tacagaatag gttggcagaa 2340
 gtgtcaccct gtgggtgtct cctcggggg ctcttccctc agacctccc ctcacttaca 2400
 25 taaagctccc ttgaagcaag aaagaggggt ccagggtctc aaaaactggaa gcacagcctc 2460
 ggggatgggg agggaaagac ggtgctatat ccagttcctg ctctctgctc atgggtggct 2520
 gtgacaaccc tggcctcact tgattcatct ctggttttct tgccaccctc tgggagtccc 2580
 oatcccatct tcatcctgag cccaaccagg ccttgccatt ggccctctgt cccttggcac 2640
 acttgtaccc acaggtgagg ggcaggacct gaaggtattg gcctgttcaa caatcagtca 2700
 30 tcatgggtgt ttttgtcaac tgettgttaa ttgatttggg gatgtttgce ccgaatgaga 2760
 ggttgaggaa aagactgttg gtggggaggc cctgcctgac ccatcccttt tctttctgg 2820
 cccagccta ggtggaggca agtggaaat cttatatggg gcgatttggg ggctcgggga 2880
 ggcagagaat ctcttgggag tcttgggtgg cgctgggtga ttctgtttcc tcttgatctc 2940
 aaagcacaat gtggatttgg ggaccaaagg tcagggacac atccccttag aggacctgag 3000
 tttgggagag tgggtgagtgg aaggaggag cagcaagaag cagcctgttt tcaactcagct 3060
 35 taattctcct tcccagataa ggcaagccag tcatggaatc ttgtctgagg ccctccctct 3120
 actcttctct tctaaaaaat aggggcogtt ttcttacaca ccccagaga gaggagggac 3180
 tgtcacactg gtgctgagt accggggggt gctgggcgtc tgttctttac caaaaccatc 3240
 catccctaga agacacaga gcctgaggg gctgggctgg gctgggctga gccctgggtc 3300
 ttctctacag ttcacagagg tcttctcagc catttaatcc caggaaagag gcatcaaagc 3360
 40 tagaatgtga atataaactt tgtgggcaa tactaagaat aacaagaagc ccagtgtgga 3420
 ggaaagtgg ttctccagc actgcctcct gttttctccc tctcatgtcc ctccagggaa 3480
 aatgacttta ttgcttaatt tctgccttcc cccctcaca catgcacttt tgggcctttt 3540
 tttatagctg gaaaaaacia aataccaccc tacaacactg tatttaaaaa gaaacagaaa 3600
 tgaccacgtg aaatttgcct ctgtccaaac atttcatccg tgtgtatgtg tatgtgtgtg 3660
 45 agtgtgtgaa gccgcagtt catcttttta tatggggttg ttgtctcatt ttggtctgtt 3720
 ttggctccct cctctgtgg cttgtgctcg ggatcaaacc tttctggcct gttatgatcc 3780
 tgaacatttg acttgaacca caagtgaatc tttctcctgg tgactcaaat aaaagtataa 3840
 ttttta 3846

<210> 2

<211> 1711

<212> DNA

<213> Homo sapiens

<400> 2							
5	gaggggaaggc	aggaaggagg	cagccgaagg	ccgagctggg	tggctggacc	gggtgctggc	60
	tgcgcgcgct	gctttcggct	cccacggcct	ctcccatgog	ctgagggagc	ccggctgcgg	120
	gccggcggcg	ggaggggagg	ctcctctcca	tggctccagaa	gaccagcatg	tcccggggcc	180
	cttaccacc	ctcccaggag	atccccatgg	aggtcttcga	ccccagccc	cagggcacaat	240
10	acagcaagag	gaaagggcga	ttcaaacggg	cagatgggag	caogtctctg	gataccacat	300
	ccaacagctt	tgtccgccag	ggctcagogg	agtcctacac	cagcogtcca	tcagactctg	360
	atgtatctct	ggaggaggac	cgggaagcct	taaggaagga	agcagagcgc	caggcattag	420
	ogcagctcga	gaaggccaag	accaagccag	tggcatttgc	tgtgaggaca	aatgttggt	480
	acaatccgtc	tccaggggat	gaggtgcctg	tgcagggagt	ggccatcacc	ttcgagccca	540
	aagacttctt	gcacatcaag	gagaaataca	ataatgactg	gtggatcggg	cggctggtga	600
15	aggagggctg	tgaggttggc	ttcattccca	gccccgtcaa	actggacagc	cttcgcctgc	660
	tgcaggaaca	gaagctgcgc	cagaacggcc	tcggctccag	caaatcaggc	gataactcca	720
	gttccagctt	gggagatgtg	gtgactggca	ccgcgcggcc	cacacccctt	gccagtcca	780
	aacagaagca	gaagtgcaca	gagcatgtgc	ccccctatga	cgtggtgcct	tccatgaggc	840
	ccatcatctt	ggtgggaccg	tcgctcaagg	gctacgaggt	tacagacatg	atgcagaaag	900
20	ctttatttga	cttcttgaag	catcggtttg	atggcaggat	ctccatcact	cgtgtgacgg	960
	cagatatttc	cctggctaag	cgctcagttc	tcaacaaccc	cagcaaacac	atcatcattg	1020
	agcgctccaa	cacacgctcc	agcctggctg	aggtgcagag	tgaaatcgag	cgaatcttcg	1080
	agctggcccg	gacccttcag	ttggctcgct	tggatgctga	caccatcaat	cacccagccc	1140
	agctgtccaa	gacctcgctg	gccccatca	ttgtttacat	caagatcacc	tctcccaagg	1200
25	tacttcaaag	gctcatcaag	tcccaggagaa	agtctcagtc	caaacacctc	aatgtccaaa	1260
	tagcggcctc	ggaaaagctg	gcacagtgcc	cccctgaaat	gtttgacatc	atcctggatg	1320
	agaaccaatt	ggaggatgcc	tgcagagcat	tggcggagta	cttggaagcc	tattggaagg	1380
	ccacacaccc	gcccagcagc	acgccacca	atccgctgct	gaaccgcacc	atggctaccg	1440
	cagccctgcg	ccgtagccct	gcccctgtct	ccaacctcca	ggtacagggt	ctcacctcgc	1500
	tcaggagaaa	cctcggtctt	tggggcgggc	tggagtcttc	acagcggggc	agtgtggtgc	1560
30	cccaggagca	ggaacatgcc	atgtagtggg	cgccctgccc	gtcttccctc	ctgctctggg	1620
	gtcggaaact	gagtgcaggg	aacatggagg	aggaagggaa	gagctttatt	ttgtaaaaaa	1680
	ataagatgag	cggcaaaaaa	aaaaaaaaaa	a			1711
<210> 3							
35	<211> 698						
<212> DNA							
40	<213> Homo sapiens						
<400> 3							
45	ttttcctttc	gctgctgcgg	ccgcagccat	gagtatgctc	aggcttcaga	agaggctcgc	60
	ctctagtgtc	ctccgctgtg	gcaagaagaa	ggctctggta	gaccccaatg	agaccaatga	120
	aatcgccaat	gccaactccc	gtcagcagat	ccggaagctc	atcaaagatg	ggctgatcat	180
	ccgcaagcct	gtgaagggtc	attcccgggc	tcgatgcggg	aaaaacacct	tggcccgcgg	240
	gaagggcagg	cacatgggca	taggtaagcg	gaagggatca	gccaatgccc	gaatgccaga	300
	gaaggtcaca	tggatgagga	gaatgaggat	tttgccggcg	ctgctcagaa	gatacgtga	360
50	atctaagaag	atcgatcgcc	acatgtatca	cagcctgtac	ctgaagggtga	aggggaatgt	420
	gttcaaaaac	aagcggattc	tcattggaaca	catccacaag	ctgaaggcag	acaaggcccg	480
	caagaagctc	ctggctgacc	aggctgaggc	ccgcaggtct	aagaccaagg	aagcacgcaa	540
	gcgcctgtaa	gagcgcctcc	aggccaagaa	ggaggagatc	atcaagactt	tatccaagga	600
	ggaagagacc	aagaaataaa	acctcccact	ttgtctgtac	atactggcct	ctgtgattac	660
55	atagatcagc	cattaaaata	aaacaagcct	taactctgc			698

<210> 4

<211> 5810

<212> DNA

<213> Homo sapiens

<400> 4

gggaagatgg	cgggggcctc	gagcaccctc	ctctttcttg	cgccggggac	ttcagattga	60
tccttcccgg	gaagagtagg	gactgctggt	gccttgctgc	ccgggatccc	gagccaactt	120
gtttcctcgg	ttagtgggtg	ggaagggtt	atccttttgt	ggcggatcta	gcttctcctc	180
gccttcagga	tgaagctca	gggggaaac	cgaggagtoa	gaaaagctga	gtaagatgag	240
ttctctcctg	gaacggctcc	atgcaaaatt	taaccaaaat	agaccctgga	gtgaaacat	300
taagcttggtg	cgtaagctca	tggagaagag	ggttggtgatg	agttctggag	ggcatcaaca	360
tttggtcagc	tgtttggaga	cattgcagaa	ggctctcaaa	gtaacatctt	taccagcaat	420
gactgatcgt	ttggagtcca	tagcaggaca	gaatggactg	ggctctcatc	tcagtgccag	480
tggaactgaa	tgtttacatca	cgtaagatat	gttctatgtg	gaagtgcagt	tagatcctgc	540
aggcagcctt	tgtgatgtaa	aagtggctca	ccatggggag	aatcctgtga	gctgtccgga	600
gcttgtagag	cagctaaggg	aaaaaaattc	tgtatgaattt	tctaagcacc	ttaaggccct	660
tgttaatctg	tataaccttc	caggggacaa	caaaactgaag	actaaaatgt	acttggtctc	720
ccaatcctta	gaacaagatc	ttcttaaaat	ggcaattatg	tactggaaag	caactaatgc	780
tgggtcccttg	gataagattc	ttcatggaag	tgttggtctat	ctcacaccaa	ggagtggggg	840
tcatttaaatg	aacctgaagt	actatgtctc	tcctcttgac	ctactggatg	acaagactgc	900
atctcccatc	attttgcattg	agaataatgt	ttctcgatct	ttgggcatga	atgcatcagt	960
gacaattgaa	ggaacatctg	ctgtgtacaa	actcccaatt	gcaccattaa	ttatgggggtc	1020
acatocagtt	gacaataaat	ggaccccttc	cttctcctca	atcaccagtg	ccaacagttg	1080
tgatcttccct	gcctgtttct	tcttgaaatt	tcaccagcca	atccagctat	ctagagcatt	1140
tgttcagaaa	ctgcagaact	gcacaggaat	tcattgtttt	gaaactcaac	caacttatgc	1200
aacctgtgat	gaactgatca	ctcagtttga	gctatcaaag	gacctgacc	ccataccttt	1260
gaatocacaac	atgagatttt	atgctgtctc	tcctggctcag	cagcactgct	atttctcctca	1320
caaggatgct	cctcttccag	atggccgaag	tctacagga	accttgttta	gcaaaatcac	1380
cttctagcac	cctggccgag	ttctctttat	cctaaatctg	atcagacacc	aagtggccta	1440
taacaccctc	attggaagct	gtgtcaaaag	aactattctg	aaagaagatt	ctcctgggct	1500
tctccaatttt	gaagtgtgtc	ctctctcaga	gtctcgtttc	agcgtatctt	ttcagcacc	1560
tgtgaatgac	tcctgtgtgt	gtgtggtaat	ggatgtgcag	ggcttaacac	atgtgagctg	1620
taaaactctac	aaagggtgtg	cggatgcact	gatctgcaca	gatgacttca	ttgccaaagt	1680
tgttcaaaaga	tgtatgtcca	tcctgtgtac	gatgagggct	attcggagga	aagctgaaac	1740
caatcaagcc	gacacccag	cactgtccct	cattgcagag	acagttgaag	acatgggtgaa	1800
aaagaacctg	cccccgcta	gcagcccagg	gtatggcatg	accacaggca	acaacccaat	1860
gagtgggtacc	actacatcaa	ccaacacctt	tcgggggggt	ccatttgcca	ccttggtttaa	1920
tatgagcatg	agcatcaaag	atcggcatga	gtcgggtggc	catggggagg	acttcagcaa	1980
ggtgtctcag	aacccaattc	ttaccagttt	gttgcaaatc	acagggaacg	gggggtctac	2040
cattggctcg	agtcgcagcc	ctcctcatca	cacgcgcgca	cctgtctctt	cgatggccgg	2100
caacaccaag	aaccacccga	tgctcatgaa	ccttctcaaa	gataatcctg	cccaggattt	2160
ctcaaccttt	tatggaagca	gccttttaga	aaggcagaac	tcctcttccg	gctcaccctc	2220
catggaataa	tgctcgggga	gcaacaagac	caagaaaaag	aagtcatcaa	gattaccacc	2280
tgagaaacca	aagcaccaga	ctgaagatga	ctttcagagg	gagctatctt	caatggatgt	2340
tgactcacag	aacctatctc	ttgatgtcaa	catgacagct	gacacgctgg	atacgccaca	2400
catcactcca	gctccaagcc	agtgtagcac	tcctcccaaca	acttaccac	aaccagtacc	2460
tcacccccaa	cccagttatc	aaaggatggt	ccgactatcc	agttcagaca	gcattggccc	2520
agatgtaact	gacatccttt	cagacattgc	agaagaagct	tctaaacttc	ccagcactag	2580
tgatgattgc	ccagccattg	gcacccctct	tcgagattct	tcaagctctg	ggcattctca	2640
gagtaccctg	tttgactctg	atgtctttca	aactaacaat	aatgaaaatc	catacaactga	2700
tcagctgat	cttattgcag	atgctgctgg	aagccccagt	agtgaactctc	ctaccaatca	2760
tttttttcat	gatggagtag	atttcaatcc	tgattttattg	aacagccaga	gccaaagtgg	2820
ttttggagaa	gaatattttg	atgaaagcag	ccaaagtggg	gataatgatg	atttcaaagg	2880

atttgcattct caggcactaa atactttggg ggtgccaatg cttggagggtg ataattgggga 2940
 gaccaagttt aagggcaata accaagccga cacagttgat ttcagtatta tttcagtagc 3000
 cggcaaaagct ttagctcctg cagatcttat ggagcatcac agtggtagtc agggtcottt 3060
 actgaccact ggggacttag ggaaagaaaa gactcaaaag agggtaaagg aaggcaatgg 3120
 caccagtaat agtactctct cggggcccgg attagacagc aaaccaggga agcgcatcg 3180
 gaccccttct aatgatggga aaagcaaaaga taagcctcca aagcggaaga aggcagacac 3240
 tgagggaag tctccatctc atagtctctc taacagacct tttaccccac ctaccagtac 3300
 aggtggatct aaatcgccag gcagtgcagg aagatctcag actccccag gtgttgccac 3360
 accacccatt cccaaaatca ctattcagat tccaaaggga acagtgatgg tgggcaagcc 3420
 ttcctctcac agtcagtata ccagcagtggt ttctgtgtct tcctcaggca gcaaaagcca 3480
 ccatagccat tcttctctct ctctctctc tgcttcacc tcagggaaga tgaaaagcag 3540
 taaatcagaa gggtcatcaa gttccaagtt aagtcagat atgtattcta gccagggtc 3600
 ttctggatct agccagttca aaaattcatc ccagtctggg gggaagccag gctcctctcc 3660
 cataaccaag catggactga gcagtggctc tagcagcacc aagatgaaac ctcaaggaaa 3720
 gccatcatca ottatgaatc cttctttaag taaaccaaac atatccctt ctcatccaag 3780
 gccacctgga ggctctgaca agcttgccct tccaatgaag cctgttctct gaactcctcc 3840
 atcctctaaa gccaaagtccc ctatcagttc aggttctggt ggttctcata tgtctggaac 3900
 tagttcaagc totggcatga agtcatcttc aggtgttagga tcctcaggct cgttgtccca 3960
 gaaaactccc ccatcatcta attcctgtac ggcatcttcc tctcctttt cctcaagtgg 4020
 ctcttccatg tcatcctctc agaaccagca tgggagttct aaaggaaaat ctcccagcag 4080
 aaacaagaag cgtctcttga cagctgtcat agataaaact aagcatgggg ttgtcaccag 4140
 tggccctggg ggtgaagacc cactggacgg ccagatgggg gtgagcaca atctctccag 4200
 ccatcctatg tcctccaaac ataacatgtc aggaggagag tttcagggca agcgtgagaa 4260
 aagtataaaa gacaaatcaa aggtttccac ctccgggagt tcagtggatt cttctaagaa 4320
 gacctcagag tcaaaaaatg tggggagcac aggtgtggca aaaattatca tcagtaagca 4380
 tgatggaggc tcccctagca ttaaagccaa agtgactttg cagaaacctg gggaaagtag 4440
 tggagaaggg cttaggcctc aaatggcttc ttctaaaaac tatggctctc cactcatcag 4500
 tggttccact ccaaagcatg agcgtggctc toccagccat agtaagtcac cagcatatac 4560
 cccccagaat ctggacagtg aaagtgaatc aggtcctcc atagcagaga aatcttatca 4620
 gaatagtccc agctcagacg atggtatccg accaottcca gaatacagca cagagaacaa 4680
 taagaagcac aaaaaggaaa agaagaaagt aaaagacaaa gatagggacc gagaccggga 4740
 caaagaccga aagttaaaaa aatctcatag catcaagcca gagagttggt ccaaatcacc 4800
 catctcttca gaccagtcct tgtctatgac aagtaacaca atottatctg cagacagacc 4860
 ctcaaggctc agcccagact ttatgattgg ggaggaagat gatgatctta tggatgtggc 4920
 cctgattggg aattaggaac cttatttctc aaaagaaaca gggccagagg aaaaaaaact 4980
 attgataagt ttataggcaa accaccataa ggggtgagtc agacaggctc gatttggtta 5040
 agaatcctaa atggcatggc tttgacatca agctgggtga attagaaagg catatccaga 5100
 ccctattaaa gaaaccacag ggtttgattc tggttaccag gaagtcttct ttgttctgt 5160
 gccagaaaga aagttaaaaa acttgcttaa gaaagggagg ggggtgggag ggggttaggg 5220
 agagggagg gagggaaaca gttttgtggg aaatattcat atatattttc ttctcccttt 5280
 ttccattttt aggccatgtt ttaaactcat tttagtgcag gtatatgaag ggctgggcag 5340
 aaaatgaaaa agcaatacat tocttgatgc atttgcattg aggttgttca actttgtttg 5400
 aggtagtgtt ccgtttgagt catgggcaaa tgaaggactt tggctatttt ggacacttaa 5460
 gtaatgtttg gtgtctgttt cttaggagtg actgggggag ggaagattat tttagctatt 5520
 tatttgtaat attttaacc tttatctgtt tgtttttata cagtgtttcg ttctaaatct 5580
 atgaggttta gggttcaaaa tgatggaagg ccgaagagca aggttatat ggtggtaggg 5640
 agcttatagc ttgtgcta atctgtagcat caagcccaag caaattagtc agagcccgcc 5700
 ttttagagta aatataatag aaaaaccaa atgatatttt tatttttagga gggtttaa at 5760
 agggttcaga gatcatagga atattaggag ttaoctctct gtggagggtat 5810

<210> 5

<211> 5515

<212> DNA

<213> Homo sapiens

[illegible]

	cagctgaatc	aaagtgaatt	ggcagtggtta	ttaaacctgc	tgcagagcca	aaccgacctg	3420
	agcatccctc	aaatggcaca	gctgcttaac	atccactcca	acccagagat	gcagcagcag	3480
	ctggaagccc	tgaaccaatc	catcagtgcc	ctgaoggaag	ctacttocca	gcagcaggac	3540
5	tcagagacca	tggccccaga	ggagtgctttg	aaggaagcac	cctctgcccc	agtgtacctg	3600
	ccttcagcag	aacagatgac	ccttgaagct	tcaagcacac	cagctgacat	gcagaaatata	3660
	ttggcagttc	tcttgagtca	gctgatgaaa	acccaagagc	cagcaggoag	tctggaggaa	3720
	aacaacagtg	acaagaacag	tgggccacag	gggccccgaa	gaactcccac	aatgccacag	3780
	gaggaggcag	cagcatgtcc	tcctcacatt	cttccaccag	agaagaggcc	ccctgagccc	3840
	cccggaacctc	caccgcccgc	acctccaccc	cctctggttg	aaggcgatct	ttccagcgcc	3900
10	ccccaggagt	tgaaccacag	cgtgacagcc	gccttgctgc	aacttttato	ccagcctgaa	3960
	gcagagcctc	ctggccacct	gccacatgag	caccaggcct	tgagaccaat	ggagtactcc	4020
	acccgacccc	gtccaaaacag	gacttatgga	aacactgatg	ggcctgaaac	agggttcagt	4080
	gccattgaca	ctgatgaacg	aaaactctggt	ccagccttga	cagaatcctt	ggtccagacc	4140
	ctggtgaaga	acaggacctt	ctcaggctct	ctgagccacc	ttggggagtc	cagcagttac	4200
15	cagggcacag	ggtcagtgca	gtttccaggg	gaccaggacc	tcogttttgc	caggggtcccc	4260
	ttagcgttac	acccggtggt	ogggcaacca	ttcctgaagg	ctgaggggaag	cagcaattct	4320
	gtggtacatg	cagagaccaa	attgcaaaac	tatggggagc	tggggccagg	aaccactggg	4380
	gccagcagct	caggagcagg	ccttcactgg	gggggcccac	ctcagttctc	tgcttatgga	4440
	aaactctatc	gggggcctac	aagagtccca	ccaagagggg	gaagagggag	aggagtctct	4500
	tactaaccac	gagacttcag	tgtcctgaaa	gattcctttc	ctatocatcc	ttccatccag	4560
20	ttctctgaat	ctttaatgaa	atcatttgcc	agagcgaggt	aatcatctgc	atttggtctac	4620
	tgcaaaagctg	tcogttgtat	tccttgctca	cttgctacta	gcaggcgact	taggaaataa	4680
	tgatgttggc	accagttccc	cotggatggg	ctatagccag	aacatttact	tcaactotac	4740
	cttagtagat	acaagtagag	aatatggaga	ggatcattac	attgaaaagt	aaatgtttta	4800
	ttagttcatt	gcctgcactt	actggctcga	agagagaaa	aacagtttca	gtattgagat	4860
25	ggctcaggag	aggtctcttg	atttttaaag	ttttgggggtg	gggggttgtg	tgtggtttot	4920
	ttcttttgaa	ttttaattta	gggtgttttg	gtttttttcc	tttaaagaga	atagtgttca	4980
	caaaaattga	gotgctcttt	ggcttttgct	ataagggaaa	cagagtggcc	tggtctgattt	5040
	gaataaatgt	ttctttctct	tcacccatct	cacattttgc	ttttaagtga	acactttttc	5100
	cccattgagc	atcttgaaac	tacttttttt	ccaaaataat	tactcatcct	taaaagtttac	5160
30	tcacttttga	caaaagatac	gcccttctcc	ctgcacataa	agcaggttgt	agaaagttggc	5220
	attctttggc	aagtagtag	actttaccca	gtctctttcc	ttttttgctg	atgtgtgtct	5280
	tctctctctc	tttctctctc	tctctctctc	tctctctctc	tctgtctgtc	tcgttctgtc	5340
	gctctogotg	tttctctctc	tttgaggcat	ttgtttggaa	aaaatcgttg	agatgcccac	5400
	gaacctggga	taattcttta	ctttttttga	aataaaggaa	aggaaattoa	aaaaaaaaaa	5460
	aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	aaaaa	5515
35	<210> 6						
	<211> 6131						
40	<212> DNA						
	<213> Homo sapiens						
45	<400> 6						
	gaattctagg	cccagttctg	tgtttccctc	gtgtgttcc	aggcaggtca	gtttccctcc	60
	atgggcctct	gtaagatgag	gagttggaga	ggtacattct	caggctactt	tcaactocca	120
	gccaaagtac	tcaagagtcc	caggcagcac	cagcaccctc	atctccaagg	cctcctgatg	180
	tgtgtctcta	tttagaactt	aatccaacct	acccaacatc	agatcagtg	cttaccagcc	240
50	caaggctcc	ggggagcctc	ctagagggag	agagccctgc	ccaccagat	tgagggtaaa	300
	ggcctccccc	tgctcatttt	tgtaccacca	cagtgccttg	cacatggtag	acatcaaaat	360
	gtgtgtgctg	aaagtataat	tgaagttgtg	tatatatgtc	agctagagtg	tctggagggg	420
	cagaaatgtg	ggtctaaaac	atacaaatgc	tccaaatggg	gtgtgggcaa	gggtctgtct	480
	acaccaggct	gtgattacct	gctcacatac	atgtgtctat	ctgagtaggg	gtatgttata	540
	tatttttcta	caccacaggg	tgaggaacag	gtatatgtgt	gcatgtgtat	gcatccgtgt	600
55	gtgtgtgtat	gtgtgtgtgc	atgagtggtg	gtgtgtgtgt	ccaaagccac	ctcttcaacc	660

	tgtgccattt	gtatctgtgt	ctggcccaat	gagagtgttg	aaaggtgagc	cacaagataa	720
	aacagcaact	toctacctcc	cttatcaaga	cagctgtctg	acctacctcc	ccttggccac	780
	tcttgggatt	actgggggtt	gcttcagtat	tttcagattt	ttcagaaggg	gaggagaatg	840
5	cttgagtctc	atccaggaac	ttaggcagtt	ctcagcactg	cctgcttoct	ctccctcaaa	900
	taaccaagtc	tgaagaccag	gagagaaagc	cgctggtgga	ctggtcacct	gtctggcagt	960
	gggaggagga	gagtgcagag	tttctaggta	ggaatccaga	cttagacct	ccctccacc	1020
	cccagatggg	tggtgcacag	gctcatctcg	cgccccctcc	ccactccacc	ctaactgga	1080
	tacgccccca	acaaccaagg	aaagatctcc	catcggtcga	ctccacagat	acacacatgt	1140
10	ccccacagac	acacacacgc	ccatgcagag	gcacagacat	ccaggcacat	ctttcccttt	1200
	ctctgtcttt	cccttggttt	gaatttctgt	tagccacata	tgttgttgt	gcgtgagggt	1260
	gggtggggga	ggggcagaca	gggatgaggg	atggcatggt	gccaacatct	acctatgggg	1320
	ctcggggccag	ggacgcccc	tacagccatc	ctgggagggg	gtctcagctg	tccttttgtg	1380
	gccaaggggg	ccctcctggg	gagtgggggc	aagcacagag	gtcctttctc	cccaaccggg	1440
	ggctctggct	ctgaccacc	ttgggggct	gcaggggagg	aaatggacag	agcgggaccc	1500
15	tgaggggagca	tagaattggc	caccacgagc	ccccagtgtc	cagccttgcc	accccatgt	1560
	tcccgtagag	gggtctctat	atacaggggg	caactcctcc	caccttctc	tcaatccctg	1620
	ctttccctgc	gttgggcggg	gaggggaggg	cggcagaaat	atttatttat	ttcctttatt	1680
	tatttaattt	tttttttttt	tttttgaggt	agagagtgc	agatggcggc	gggtcccggg	1740
	ggagccggct	ctccccaggt	gcagacgcac	gccaatccac	gtctctcatg	tgatagctgc	1800
20	tgccctgagc	gtgccaaagg	catatggcct	ggcatagagg	ctggtacccc	gcctggtaga	1860
	gatgccacac	tcgtctcccg	gttcgcatgg	cgctotgaag	acgcggcgcc	ccgcgcctt	1920
	gaggagccgc	tgcccccgct	ccctgaagat	gggggaacaa	tgaaataagc	gagaagatcc	1980
	ctctctctcc	ccctctctct	cttgccccct	ccccccctcc	cctccccctc	cccttgact	2040
	cctctccgag	gtaagttgtc	cgaaagggag	cgagatctga	ccgcgcgggt	gggaggaggg	2100
	gcgccagctt	ogggcgacag	gagggctctc	aaatacctcc	ttcctgggat	gatgcccccc	2160
25	tcattgggtg	ggcatcgag	ggggcccgag	ttctctctcc	cttaggggct	gcagccacag	2220
	gggctgcaga	ggaggtgtct	ctgcctgcga	tgggtctggg	ggggggggaa	ggcaggatca	2280
	cgagggggga	tatgcgaaga	ggccgagacg	gaggaccctc	ccatggttgt	cccaaaaagc	2340
	ctgccacctt	tccccaccac	cgaaaaaagg	gaagcaaaac	aacaaatttg	gatttttccc	2400
	ccatcaatcc	caaaaatacaa	cgagatctga	agagccttgt	gggaggggag	cagcttgaag	2460
30	ggggaagggg	gtccctgacc	gcagagggga	cgactgggc	tcgtctctct	cagtctctcc	2520
	cccacgcccc	gctgcttcag	tcctgcgcgc	ccagagccgg	ctccgggagc	tggggaccca	2580
	tcggctagag	ggacgcgac	tcgccctctc	ggatattggg	gtgcgggggt	gggggcccag	2640
	caaggggcgg	cgccgagcca	agttgcaaat	tggattaggg	agcgtggggg	tgagagccac	2700
	gggaggggtg	agggagctgg	gccccggggc	ccgggcccgc	agagcgcgga	goggggcagc	2760
	tgtccccacc	ggcgcccgac	cagcctctct	ccaccgccag	gagagaacgg	gcttcagggg	2820
35	cgagcgcgcc	gcctccccct	gcaaagatat	ctggtcccta	aaacccccac	ccggtccctg	2880
	ccctgacctt	gagaagaagc	aggcgcgggg	agcagccccc	cattcaagcg	aggggcccag	2940
	ccggggccca	gcgcggggga	gagggcctgg	gcagagatcc	caggccggca	gcccggtagg	3000
	gctgggcggg	ctctgggcgg	ggcagggcgg	ggaggtgggc	atccagggta	gcctaggcag	3060
	gagcccgcac	gagactcggg	ggtggaggag	ggttgtgggg	gggogtcggg	accccgccgc	3120
40	gccccctcact	ttgtgtgtgc	tgtctccctc	tcgcccccgc	ggggcgccct	caggcaccat	3180
	gctgaccogc	ctgttcagcg	agcccgccct	tctctcgga	gtgcccaggt	tcgccagctg	3240
	gggcgacggc	gaagacgacg	agccgaggag	cgacaagggc	gacgogccgc	caccgccacc	3300
	gcctgcgccc	gggcccaggg	ctccggggcc	agcccgggcg	gccaagccag	tcctctcccg	3360
	tggagaagag	gggacggagg	ccacgttggc	cgaggtcaag	gaggaaggcg	agctgggggg	3420
45	agaggaggag	gaggaagagg	aggaggaaga	aggactggac	gaggcggagg	gcgagcgggc	3480
	caagaagcgc	gggcccaga	agcgcaagat	gaocaaaggc	cgcttgagc	gctccaagct	3540
	tcggcggcag	aaggcgaaag	cgccgggagc	caaccgcag	cacgacctga	acgcagccct	3600
	ggacaacctg	cgcaaggtgg	tgccctgcta	ctccaagacg	cagaagctgt	ccaagatcga	3660
	gacgtgcgc	ctagccaaga	actatatctg	ggcgtctctg	gagatctctg	gctccggcaa	3720
	gcgccagac	ctagtgtcct	aogtgcagac	tctgtgcaag	ggtctgtctg	agcccaccac	3780
50	caatctgggtg	gcccgtgtgc	tgagctcaa	ctctcgcaac	ttcctcacgg	agcaaggcgc	3840
	cgacggtgcc	ggccgcttcc	acggctcggg	cgcccggttc	gccaatgcacc	cctaccgta	3900
	cccggtgctg	cgccgtggcg	ggcacagtg	ccaggcgggc	ggcgccctgg	gcggcgggcg	3960
	ggcgcaogcc	ctgcggaccc	acggctactg	cgccgcctac	gagacgctgt	atgcggcggc	4020
	aggcggtggc	ggcgcgagcc	cggactacaa	cagctccgag	tacgagggcc	cgctcagccc	4080
55	cccgctctgt	ctcaatggca	acttctcact	caagcaggac	tcctcgcccg	accacgagaa	4140
	aagctaccac	tactctatgc	actactcggc	gctgcccggg	tcgcgcacag	gccacggggt	4200

agtcttcggc togctggctg tgcgcggggg cgtccactcg gagaatctct tgtcttaoga 4260
 tatgcacctt caccacgacc ggggccccat gtacgaggag ctcaatgcgt tttttcataa 4320
 ctgagacttc ggcgcggctc ccttcttttt cttttgcctt tgcccgcccc cctgtcccca 4380
 5 gccccagca ggcgcagggt cacccccac ctaccccggc gccgggcggc gggagcgggc 4440
 caccggctct gccgctctcc tggggcagcg cagtctctgt acctgtgggt ggctgtctcc 4500
 aggggcctcg cttccccag gggactcgcc ttctctctcc ccaagggtt cctcctct 4560
 ctctcccaag gagtgtctt ccaggacct ctctccggg gctccctgga ggcacccctc 4620
 cccattcccc aatatctctg ctgaggtttc ctctccccc tctccctgc aggcccaagg 4680
 cgttggtaag ggggcagctg agcaatggaa cgcgtttccc cctctcatta ttattttaaa 4740
 10 aacagacacc cagctgccga ggcaaaaagg agccaggcgc tccctctttc ttgaagaggg 4800
 tagtattttg gggcgcggag cccgggcctg gaacgcctc acccgcaacc tccagtctcc 4860
 gogttttgcg attttaattt tggcgggagg ggaagtggat tgagaggaaa gagagaggcc 4920
 aagacaattt gtaactagaa tccgtttttc ctttttcoct tttttaaaca aacaacata 4980
 caaaaaaaa aaaaaaaa aaaaaaaa aagctaagag gcgacggaag ccgaacgcag 5040
 15 agtcggatc ggagagaaaa cgcagtaagg acttttagaa gcaataaaa gcaaaaaaaa 5100
 caaaaaacaa aaaaacaaac aaaaaaaa cactactacc aataatcaaa gacacaaata 5160
 tctatgcaag gaggctccac tgagcctcgc gcccgggccc gcccccgga tgccccgcc 5220
 ggctgcggg cgcgcccgcc cgagcgggga tctgtgcaact ttggtgaagt gggggcccg 5280
 gccccccct cccctccccc aggttcttac aatcagtgac tcggagattt ggggccccag 5340
 20 tgccactgoc ctccccgcc cctccccgt tgtgcgtcat gctgtttttt aaaaacctgt 5400
 ttccaaattt gtatggaatg gcaaaactgtt ggggggtcgg tttggggagg gagggtttgc 5460
 atgaaagaca cagcacacc acacgcacg cacaagcagg cccggcgccg gcgtccgggg 5520
 ggcagaagga ggtgagctcg ccggtctctc ctccccggc ccattctgtc cctcctggg 5580
 gtgaggggtg gggatggaga cctgggggca gccccacccc tgcccggaact gtgcctcgg 5640
 ggggtgccacc tggcgatttc cgggtgtctg agagagtatt ttttggtcca aggagtctc 5700
 25 ttggctttag cttgtgggtg ggcggggaga ggtctgagg gctctactgg aggttcccc 5760
 aaaaaggggc aaagagagac cctctgccc cgggaggcag gggatcagg atccaaatac 5820
 acgatgcaaa aatgcaatcc cacaggogac acaccacac actcaccac acacacgcaa 5880
 ttttaccttc ctcttgtagc gaagatgaaa ctcccgctg acaccogaag tgcattgcgt 5940
 gtttctgttc agtttaatga cgattaataa atatttatgt aaatgagatg caaagccgga 6000
 30 ccggtttctc acggtggcct catttcattg aggggggaga gaaggtttga gctggggctg 6060
 ggggtgatgaa ggcagagtgt caagtgactg tgcagaggcc aaacagaggg acttccagc 6120
 aaaaagcact g 6131

<210> 7

35 <211> 2020

<212> DNA

<213> Homo sapiens

40

<400> 7

gctactgagg ccgcggagcc ggactgcggt tggggcggga agagccgggg ccgtggctga 60
 catggagcag ccctgtctgt gaggcgcgc cctccccgcc ctgaggtggg ggcccaccag 120
 45 gatgagcaag ctgccaggg agctgacccg agacttggag cgcagcctgc ctgccgtggc 180
 ctccctgggc tctcactgt cccacagcca gagcctctcc tcgcacctcc ttccgcccgc 240
 tgagaagoga agggccatct ctgatgtccg ccgcaacctc tgtctctctg tcaccttcga 300
 cctgtctctc atctccctgc tctggatcat cgaactgaat accaacacag gcatccgtaa 360
 gaacttggag caggagatca tccagtacaa ctttaaaact tccttctctg acatctttgt 420
 50 cctggccttc ttccgcttct ctggactgct cctaggtctat gccgtgctgc agctccggca 480
 ctggtgggtg attcgggtca cgaagctggg gtccagtgca ttctcattg tcaaggtcat 540
 cctctctgag ctgctcagca aaggggcatt tggctaacct ctccccatog tctcttttgt 600
 cctcgcctgg ttggagacct ggttccttga cttcaaagtc ctacccagg aagctgaaga 660
 ggagcgatgg tatcttgccg ccaggttgc tgttgcccg ggacccctgc tgttctccgg 720
 55 tgcctgtctc gagggacagt totattcacc ccagaaatcc tttgcagggt ctgacaatga 780
 atcagatgaa gaagtgtctg ggaagaaaag tttctctgct caggagcggg agtacatccg 840

	ccaggggaag	gaggccacgg	cagtgggtgga	ccagatcttg	gcccaggaag	agaactggaa	900
	gtttgagaag	aataatgaat	atggggacac	cggttacacc	attgaagttc	cotttcacgg	960
	caagacgttt	atcctgaaga	ccttcctgcc	ctgtcctgog	gagctcgtgt	accaggaggt	1020
5	gatcctgcag	cccgagagga	tgggtcgtgtg	gaacaagaca	gtgactgcct	gccagatcct	1080
	gcagcgagtg	gaagacaaca	ccctcatctc	ctatgacgtg	tctgcagggg	ctgoggggcgg	1140
	cgtgggtctcc	ccaagggact	tcgtgaatgt	ccggcgcat	gagcggcgca	gggaccgata	1200
	cttggtcatca	gggatcgcca	cctcacacag	tgccaagccc	ccgacgcaca	aatatgtccg	1260
	gggagagaat	ggccctgggg	gcttcatogt	gctcaagtog	gccagtaacc	cccggtgttg	1320
	cacctttgtc	tggattctta	atacagatct	caagggccgc	ctgccccggg	acctcatcca	1380
10	ccagagcctc	gcggccacca	tgtttgaatt	tgcctttcac	ctgcgacagc	gcacagcgga	1440
	gctggggggc	cgggcgtgac	tgtgccccct	cccaccctgc	ggggccagggt	cctgtcgcca	1500
	ccacttccag	agccagaaaag	ggtgccagtt	gggctcgcac	tgcccacatg	ggacctggcc	1560
	ccaggctgtc	accctccacc	gagccacgca	gtgcctggag	ttgactgact	gagcaggctg	1620
	tgggggtggag	cactggactc	cggggcccca	ctggctggag	gaagtggggg	ctggcctgtt	1680
15	gatgtttaca	tggcgccctg	cctcctggag	gaccagattg	ctctgccccca	ccttgccagg	1740
	gcaggggtctg	ggctggggcac	ctgacttggc	tggggaggag	cagggccctg	ggcagggcag	1800
	ggcagccctgt	cacccgtgtg	aagatgaagg	ggctotctcat	ctgcctgcgc	tctcgtcgggt	1860
	tttttttagga	ttattgaaag	agtctgggac	ccttgttggg	gagtgggtgg	caggtggggg	1920
20	tgggctgctg	gccatgaatc	tctgcctctc	ccaggctgtc	cccctcctcc	cagggcctcc	1980
	tgggggacct	ttgtattaa	ccaattaaaa	acatgaattt			2020
	<210> 8						
	<211> 1730						
25	<212> DNA						
	<213> Homo sapiens						
30	<400> 8						
	gtggtgaggg	tgaactggga	ctaggcaacta	ggcctttgggt	gcaggcgcct	gaggacktgg	60
	ttgcactctc	ccttotgggg	atatgcoctt	gagcccaggc	agaggagagc	acagcccagg	120
	gcaggacctg	gcagccctgg	tacagagccc	agagggggca	tcagttcctg	ctggtcctgc	180
35	tctgtttaca	gacaasctgc	tgtcctccct	gcaaagggga	gtgggtgggg	cagaggggcaa	240
	ktgccagggg	ggcacaaggc	tgggcatgtg	gctggcatga	gacggtgtct	gagtaatgtc	300
	aggcacctgg	aggcattgac	cccaggacct	tggaccccag	acctctgacc	gtggggcagc	360
	cagcgtccag	gtaccccac	ccctgcccctg	ggtccggcgt	ccccccatta	gtgagtcttg	420
	gctctactta	tagcatctga	caccagaggg	gcccgaataa	gcccctggag	aagggggagg	480
40	agggggctat	ttaaagggcc	tgggagggga	gagagaatga	ggagtgatca	tggctacctc	540
	agagctgagc	tgcgaggtgt	cggaggagaa	ctgtgagcgc	cgggaggcct	tctgggcaga	600
	atggaaggat	ctgacactgt	ccacacggcc	cgaggagggg	tgagtgtggg	tctgctagag	660
	tccttgccctc	tgtcccccca	gagcacctc	actgagccat	gaggccagag	catgaagccc	720
	tggagaaatt	tctgggggtg	ggggcaggaa	gaatgccccca	tggggagagc	aaaggggaac	780
	caccttccct	gccccccagg	cccagcagcc	caggggagcc	ccccccccag	cctgtgcccc	840
45	gagagcaaca	gctcccagga	gctcaactgc	cctccccctct	ccccagctgc	tccttgcatg	900
	aggaggacac	ccagagacat	gagacctacc	accagcaggg	gcagtgccag	gtgctggtgc	960
	agcgcctcgc	ctggctgatg	atgcggatgg	gcacctctcg	ccgtgggctg	caggagtacc	1020
	agctgcccta	ccagcgggta	ctgcctgctc	ccatcttcac	ccttgccaag	atgggcgcca	1080
	ccaaggagga	gcgtgaggac	acccccatcc	agcttcaggga	gctgctggcg	ctggagacag	1140
50	ccctgggtgg	ccagtgtgtg	gaccgccagg	aggtggctga	gatcacaaag	cagctgcccc	1200
	ctgtggtgcc	tgtcagcaag	cccggtgcmc	ttcgtcgtct	cctgtcccgc	tccatgtccc	1260
	aggaagcaca	gagaggctga	gagggactgt	gacttgggct	ccgctgtgcc	cgccccctgg	1320
	gctggggcct	tcctggctag	gacctgtgga	ggggcagctc	gctggcccat	ggctgctttg	1380
	tagtttgccc	agagttgggg	gctagggggg	gggggagcca	gaggccagga	tgcttgagcc	1440
55	ccctgagttc	ccaaagggag	ggtggcagag	acagtgggca	ctaagggtgg	agagttgggg	1500
	gocagcacag	ctgaggaccc	tcagccccag	gagaagggac	aaaagggtact	ggtgagggga	1560

agaggtgcct	gggaggagt	gccctgatcc	aggaaaatgt	gaggggaatc	tggaacgctc	1620
taggcagaag	aagctgggag	ggagggggag	gtgaaaaggg	cagaggcaag	gatggtgggg	1680
ccccagcac	cctotgttag	tgccgcaata	aatgctcaat	catgtgccag		1730

<210> 9

<211> 3799

<212> DNA

<213> Homo sapiens

<400> 9

ctggcactgg	gtggtaacca	gcaagccagc	tgccatccgc	atccagggtt	tgtttcaatg	60
atgtctcgtg	gagaatatgg	aggggctggg	gccaggactg	tccttggctt	tgccctgggg	120
tgtgaacggg	gtcagtgaac	tctaaaacta	acctgcctct	cagttctgaa	tccagacaga	180
atcaatcctc	agctgtgtct	cgctccacac	ccctgcccct	ggaagccagg	gaaggttgga	240
ggtgctaggg	ggtcaggctc	ccctctgtga	ccctgcagc	tggtgtgggtg	actcatgtcc	300
caacctagct	gcctctccca	aggagacttt	ccctggggac	aagggggagg	gaatggcatg	360
gaggaggccc	acatcaagcg	gggccaggaa	cccacggtgg	caggagctgg	gctggtgacc	420
tacccagggc	agaaggggcc	gggactcatt	cagaggggaa	ggaaggggtc	ttcagggaaga	480
ccacggagat	gccacaggca	gaattggctt	cccatctggg	agatagggtg	ggagaccctg	540
gcattttgac	agccagaacc	tggggtgctg	agcagaatct	tcatgcctgg	cctggccggc	600
ttcggaggga	agotggaggg	ttgggtgcga	gaggagtggg	gtcagagccc	ctacatccgc	660
aggaccccaa	atcggtggg	ccccaggcc	cgactgcgc	tcccgggtgg	ccccggcggc	720
cctccgcgaa	tgctgcctgc	ccctcccttg	cccaagccct	ctgccctcac	ccgggtccgg	780
cgccgcccc	gaagtggcgg	gaacaaccgg	aaccggaacc	ttctgtcctc	gggagccccc	840
agataagcgg	ctgggaaccc	gcggggcccg	caggggaggg	ccggctgttc	cgcccgcctaa	900
gtgcattagc	acagctcacc	tcccttatcg	cgctgccat	cggacgggca	gtgccgcgcc	960
ctgctctggg	gccccgggag	cgaccacagc	ggaggccgga	acggactgtc	ctttctgggg	1020
cggggtgggg	agggggtgtc	gctggagggc	ccggtggcat	agcaacggac	gagagagccc	1080
tggaggaggg	gcggggaggg	ggagttgtgt	ggcagttota	agggaaagggt	gggtgctggg	1140
acgggtgtcc	gggaggggag	ggagcctggc	gggtctggg	gcctctggc	ggaggggcgt	1200
gcgaggggga	aactggggaa	agggcctaata	tcccagttot	ccacctcgaa	tcaggaaaaga	1260
gaaggggagg	gctgctgggc	aaaagagggtg	aatggctgcg	gggggctgga	gaagagagat	1320
gggagggggc	ggccggcggg	ggtgaggggg	tctaaagatt	gtgggggtga	ggaactgagg	1380
gtggggggcg	cccagaggcg	ggactcgggg	cggggcaggc	gaggcggagg	gcgagggctg	1440
cgggagcaag	tacggagccg	gggtgtggg	ggcagattgc	cgctgcagcc	gccgccccac	1500
tcacctccgg	tgtgtctgca	gcccggacac	taaggagat	ggatgaatgg	gtggggagga	1560
tgccggcgac	atggccccgg	gcggctcggc	ggtcagctgc	cgccccaca	gcggaccggt	1620
cggggccccg	gtcggggcgt	agaaaaaagg	gcgcgaggc	gagcggggca	ctgggcggac	1680
cgccggcgga	gcatgagcgg	cgagaccgt	agccccaatg	cgggcgcagc	ccctgactcg	1740
gccccggggc	aggcggcggt	ggcttcggcc	taccagcgct	tcgagccggc	cgccctacctc	1800
cgcaacaact	acgcgcccc	tcgcggggac	ctgtgcaacc	cgaacggcgt	cgggccgtgg	1860
aagctgcgct	gcttggcgca	gaccttcgac	accggtgagc	gggggaaact	gaggcacgag	1920
ggacaagagg	tcgtcgggga	gtgaaagcag	gcgcagggaa	ataaaaagaa	ggaaaggagg	1980
acagaccagg	cgccctaacag	atggggacca	agaaacaaga	gatagctgag	aggtgcaaac	2040
agaagagaaa	aaggagcaac	atcccttagg	agaggggcag	aggagagaga	ggtggagaga	2100
gggggcccgg	agtgtctaga	attgagagct	aaggtggggg	atgcaggaca	gactgagggtg	2160
gagatgcata	ggaggaaatg	gaggcagatg	tgggacaggg	gtgagaaact	ccaggatttc	2220
ctogctgagc	ctggctggta	ggtatagttg	ttttctttct	ttttctttat	tttattttca	2280
tttatttact	tattttttatt	ttttatttgt	tttgagacgg	agtttcgctc	ttgttgccca	2340
ggctggagta	caatggcgcc	atctcggctc	actgcaacct	ccgcctcccc	gggttcaagc	2400
gattctcttg	cctcagcttc	cctagtagct	gggattacag	gcatgcgccc	ccatgcctgg	2460
ctaattttatt	tgtattttta	gtagagacgg	gacttctcca	tggttggtcag	gctggtctcg	2520
aactcccaac	cttaggateg	accacccccg	gcctcccaaa	gtgctgggat	tacagggtgtg	2580
agccactgcg	cccggccaggt	aggtatagtc	ttctagatgt	gaaacctgag	tctcagagcg	2640

	gtgaagttcc	cttcogaagg	gcagcccatg	ttggagctgg	gttcagtcta	actctggggc	2700
	caatgotttt	tccagatgga	gacacatttg	cagaggagaa	ggaagaacta	gagagaggca	2760
	gggagatgca	ggggagggaa	gggtaaggag	gcaggggctg	cctgggctgg	ctggcaccag	2820
5	gacctctctc	ctctgccctg	cccaggtgaa	gtgtccggac	gcacctcat	cgacattggt	2880
	tcaggcccca	ccgtgtacca	gctgctcagt	gcctgcagcc	actttgagga	catcaccatg	2940
	acagatttcc	tggaggtcaa	ccgccaggag	ctggggcgct	ggctgcagga	ggagccgggg	3000
	gccttcaact	ggagcatgta	cagccaacat	gcctgcctca	ttgagggcaa	ggggtaagga	3060
	ctgggggggtg	agggttgggg	aggaggcttc	ccatagagtg	gctggttggg	gcaacagagg	3120
10	cctgagcgta	gaacagcctt	gagccctgcc	ttgtgcctoc	tgcacaggga	atgctggcag	3180
	gataaggagc	gccagctgcg	agccagggtg	aaacgggtcc	tgcccatcga	cgtgcaccag	3240
	ccccagcccc	tgggtgctgg	gagcccagct	ccccctgctg	ctgacgccct	ggtctctgcc	3300
	ttctgcttgg	aggctgtgag	cccagatctt	gccagcttcc	agcggggcct	ggaccacatc	3360
	accacgctgc	tgaggcctgg	ggggcacctc	ctcctcatcg	gggcccctga	ggagtcgtgg	3420
	tacctggctg	gggagggcag	gctgacggtg	gtgccagtgt	ctgaggagga	ggtgagggag	3480
15	gccctgggtgc	gtagtggcta	caaggctcgg	gacctccgca	cctatatcat	gcctgcccac	3540
	cttcagacag	gcgtagatga	tgtcaagggc	gtcttcttog	cctgggctca	gaaggttggg	3600
	ctgtgagggc	tgtacctggt	gcctgtggc	ccccaccac	ctggattccc	tgttctttga	3660
	agtggcacct	aataaagaaa	taataccctg	ccgtgcgggt	cagtgtctgt	tgtggctctc	3720
	ctgggaagca	gcaagggccc	agagatctga	gtgtccgggt	aggggagaca	ttcacccatg	3780
20	gctttttttc	cagaagctt					3799
	<210> 10						
	<211> 4530						
25	<212> DNA						
	<213> Homo sapiens						
30	<400> 10						
	aattctcgag	ctcgtcgacc	ggtcgaagag	ctcaggggtc	gacgagctcg	agggcgcgcg	60
	cccgcccccc	acccctcgca	gcaccccgcg	cccccgcccc	tcccagccgg	gtccagccgg	120
	agccatgggg	ccggagccgc	agtgagcacc	atggagctgg	cgcccttgtg	ccgctggggg	180
	ctcctcctcg	ccctcttgcc	ccccggagcc	gcgagcacc	aagtgtgcac	cggcacagac	240
35	atgaagctgc	ggctccctgc	cagtcccgag	acccacctgg	acatgctccg	ccacctctac	300
	cagggtcgcc	aggtggtgca	gggaaacctg	gaactcaoot	acctgcccac	caatgcccagc	360
	ctgtccttcc	tgcaggatat	ccaggagggtg	cagggctacg	tgctcatcgc	tcacaaccac	420
	gtgagcgagg	tcccactgca	gaggtctcgg	attgtgcgag	gcacccagct	ctttgaggac	480
	aactatgccc	tggccgtgct	agacaatgga	gacccgctga	acaataccac	ccctgtcaca	540
40	ggggccctcc	caggaggcct	gogggagotg	cagcttcgaa	gcctcacaga	gatcttgaaa	600
	ggaggggtct	tgatccagcg	gaacccccag	ctctgctacc	aggacacgat	tttgtggaag	660
	gacatcttcc	acaagaacaa	ccagctggct	ctcacactga	tagacaccaa	ccgctctcgg	720
	gcctgccacc	cctgtttctcc	gatgtgtaag	ggctcccgct	gctggggaga	gagttctgag	780
	gattgtcaga	gcctgacgcg	cactgtctgt	gcoggtggct	gtgcccgtg	caaggggcca	840
	ctgccactcg	actgctgcca	tgagcagtg	gctgcgggt	gcacggggcc	caagcactct	900
45	gactgcctgg	cctgcctcca	cttcaaccac	agtggcatct	gtgagctgca	ctgccagacc	960
	ctggtcacct	acaacacaga	cacgtttgag	tccatgcccc	atcccagagg	ccggtatata	1020
	ttcggcgcca	gctgtgtgac	tgctgtcccc	tacaactacc	tttctacgga	cgtgggatcc	1080
	tgcacccctcg	tctgccccct	gcacaaccac	gaggtgacag	cagaggatgg	aacacagcgg	1140
	tgtgagaagt	gcagcaagcc	ctgtgcccga	gtgtgctatg	gtctgggcat	ggagcacttg	1200
50	cgagaggtga	gggcagttac	cagtgtccaat	atccaggagt	ttgctggctg	caagaagatc	1260
	tttgggagcc	tggcatttct	gccggagagc	tttgatgggg	acccagcctc	caacactgcc	1320
	ccgctccagc	cagagcagct	ccaagtgttt	gagactctgg	aagagatcac	aggttacctc	1380
	tacatctcag	catggccgga	cagcctgcct	gacctcagcg	tcttcacaga	cctgcaagta	1440
	atccggggac	gaattctgca	caatggcgcc	tactogctga	ccctgcaagg	gctgggcac	1500
55	agctggctgg	ggctgcgctc	actgagggaa	ctgggcagtg	gactggccct	catccaccat	1560
	aacacccacc	tctgcttctg	gcacacgggtg	ccctgggacc	agctctttcg	gaacccgcac	1620

caagctctgc tccacactgc caaccggcca gaggacgagt gtgtgggcga gggcctggcc 1680
 tgccaccagc tgtgogcccg agggcactgc tggggtccag ggcccaccca gtgtgtcaac 1740
 tgcagccagt tccttogggg ccaggagtg cgtggaggaat gccgagtact gcaggggctc 1800
 5 cccagggagt atgtgaatgc caggcactgt ttgcctgccc accctgagtg tcagccccag 1860
 aatggctcag tgacctgttt tggaccggag gctgaccagt gtgtggcctg tgcccactat 1920
 aaggaccctc ccttctgcgt ggcccgtgc cccagcgggtg tgaacactga cctctcctac 1980
 atgccatctt ggaagtttcc agatgaggag ggcccatgcc agccttgccc catcaactgc 2040
 aoccatcctt gtgtggacct ggtgacaag ggctgccccg ccgagcagag agccagccct 2100
 10 ctgaogtcca togtctctgc ggtggttggc attctgctgg tctgtgtctt gggggtggct 2160
 tttgggatcc tcatcaagcg acggcagcag aagatccgga agtacacgat gcggagactg 2220
 ctgcaggaaa cggagctggt ggagccgctg acacctagcg gaggcatgcc caaccaggcg 2280
 cagatgcgga tcctgaaaga gacggagctg aggaaggtga aggtgcttgg atctggcgct 2340
 tttggcacag agacagagg catctggatc ccttagtggg agaattgtga aattccagt 2400
 15 gccatcaaaag tgttagggga aaacacatcc cccaaagcca acaaagaaat cttagacgaa 2460
 gcatacgtga tggctggtgt gggctcccca tatgtctccc gccttctggg catctgcctg 2520
 acatccacgg tgcagctggt gacacagott atgcccattg gctgcctctt agaccatgtc 2580
 cgggaaaacc gcggacgcct gggctcccag gacctgctga actggtgtat gcagattgcc 2640
 aaggggatga gctacctgga ggtgtgctgg ctgtacaca gggacttggc cgtctgggaa 2700
 gtgctggtca agagtcccaa ccatgtcaaa attacagact tcgggctggc toggctgctg 2760
 20 gacattgacg agacagagta ccattgcagat gggggcaagg tggccatcaa gtggatggcg 2820
 ctggagtcca ttctccgccc gcggttcacc caccagagtg atgtgtggag ttatggtgtg 2880
 actgtgtggg agctgatgac ttttggggcc aaaccttaag atgggatccc agcccgggag 2940
 atccctgacc tgcctgaaaa gggggagcgg ctgcccagc ccccatctg caccattgat 3000
 gtctacatga tcatggtcaa atgttggatg attgactctg aatgtoggcc aagattccgg 3060
 gagttggtgt ctgaattctc ccgcatggcc agggaccccc agcgccttct ggtcatccag 3120
 25 aatgaggact tgggcccagc cagtccttg gacagacct tctaccgctc actgctggag 3180
 gacgatgaca gggggacct ggtggatgct gaggagtatc tggtaaccca gcagggttc 3240
 ttctgtccag accctgcccc gggcgttggg ggcatggtcc accacaggca ccgcagctca 3300
 totaccagga gtggcggtgg ggacctgaca ctagggtctg agccctctga agaggaggcc 3360
 cccaggtctc cactggcacc ctccgaagg gctgcaaagc ctccccacac atgaccccag 3420
 30 ggaatggggg cagccaaggg agtaaccctg ccctctgaga ctgatggcta cgttgcccc 3480
 oggtacagtg aggaccccac tgaatatgtg aaccagccag atgttcggcc ccagccccct 3540
 ctgacctgca gccccagcc gcctgctgcc cgacctgctg gtgccactct ggaaagggcc 3600
 togcocogag agggccctct gaatggggtc gtcaaagacg tttttgcctt tgggggtgoc 3660
 aagactctct cccaggggaa gacaccccag ggaggagctg cccctcagcc ccacctcct 3720
 35 gtggagaacc ccgagtactt gacaccccag ggtgagctg accaggaccc accagagcgg 3780
 ootgccttca gccagcctt cgacaacctc cctacggcag agaaccaga gtacctgggt 3840
 ggggctccac ccagcacctt caaagggaca agtcocgaga agccctgatg tgcctcagg 3900
 ctggacgtgc cagtgtgaac cagaaggcca caagaggtgg gaggggccctc cgaccacttc 3960
 gagcagggaa ggctgactt ctgctggcat ctaagggaacc ttcttctctg cttgagttcc 4020
 caggggaacc tgccatgcca ggaacctgtc ctaagggaacc agcactgggg agtctttgtg 4080
 40 cagatggctg gaaggggtcc agcctcgttg gaagagggaac agcactgggg cctagcttgg 4140
 gattctgagg ccctgcccac tgagactcta ggttccagtg gatgccacag cccagcttgg 4200
 ccctttcctt ccagatcctg ggtactgaaa gccttaggga agctggcctg agaggggaa 4260
 cggccctaag ggagtgtcta agaacaaaag cgaccatttc agagactgtc cctgaaacct 4320
 agtactgccc ccctgagga aggaacagca atggtgtcag tatccaggct ttgtacagag 4380
 45 tgctttctct tttagttttt actttttttt ttttgttttt ttaaagacga aataaagacc 4440
 caggggagaa tgggtgttgt atggggaggc aagtgtgggg ggtccttctc cacaccaact 4500
 ttgtccattt gcaaatatat tttggaaaa 4530

<210> 11

50 <211> 2205

<212> DNA

<213> Homo sapiens

55

	<400> 11						
5	cacagggctc	ccccccgcct	ctgactttctc	tgtccgaagt	cgggacaccc	tcctaccacc	60
	tgtagagaag	cgggagtggg	tctgaaataa	aatccaggaa	tctgggggtt	cctagacgga	120
	gccagacttc	ggaacgggtg	tctgtctact	cctgtctggg	ctcctccagg	acaagggcac	180
	acaactgggt	cogttaagcc	cctctctcgc	tcagacgcca	tggagctgga	tctgtctcca	240
10	cctcatctta	gcagctctcc	ggaagacott	tggccagccc	ctgggacccc	tcctgggact	300
	ccccggcccc	ctgatacccc	tctgcctgag	gaggtaaaga	ggtcccagcc	tctcctcatc	360
	ccaaccaccg	gcaggaaact	tcgagaggag	gagaggcgtg	ccacctccct	ccccctcttc	420
	cccaacccct	tccttgagct	ctgcagtcct	ccctcacaga	gccaattct	cgggggcccc	480
	tccagtgcaa	gggggctgct	cccccgcat	gccagccgcc	cccatgtagt	aaaggtgtac	540
	agtgaggatg	gggcctgcag	gtctgtggag	gtggcagcag	gtgccacagc	tcgccacgtg	600
15	tgtgaaatgc	tgggtgcagc	agctcacgcc	ttgagcgacg	agacctgggg	gctgggtggg	660
	tggcaccccc	acctagcact	ggagcggggt	ttggaggacc	acgagtcogt	ggtgggaagt	720
	caggctgcct	ggcccgtggg	cggagatagc	cgcttcgtct	tcoggaaaaa	cttcgccaaag	780
	tacgaactgt	tcaagagctc	cccacaactcc	ctgttcccag	aaaaaatggt	ctccagctgt	840
	ctcgatgcac	acactgggtat	atcccatgaa	gacctcatcc	agaacttct	gaatgctggc	900
	agctttcctg	agatccaggg	ctttctgcag	ctgcggggtt	caggacggaa	gctttggaaa	960
20	cgctttttct	gtttcttgcg	cogattotggc	ctctattact	ccaccaaggg	cacctctaag	1020
	gatoogaggc	acctgcagta	cgtggcagat	gtgaacgagt	ccaactgtga	cgtgggtgacg	1080
	cagggccgca	agctctacgg	gatgccact	gacttcgggt	tctgtgtcaa	gccaacaacg	1140
	cttcgaaatg	gacacaaggg	gotttcggatc	ttctgcagtg	aagatgagca	gagccgcacc	1200
	tgtctggctgg	ctgccttccg	cctcttcaag	tacggggtgc	agctgtacaa	gaattaccag	1260
25	caggcacagt	ctcgccatct	gcatccatct	tgtttgggt	ccccaccctt	gagaagtgcc	1320
	tcagataata	ccctgggtggc	catggaacttc	tctggccatg	ctgggcgtgt	cattgagaac	1380
	ccccgggagg	ctctgagtg	ggccctggag	gaggcccagg	cctggaggaa	gaagacaaac	1440
	caccgcctca	gcctgcccct	gccagcctcc	ggcacgagcc	tcagtgcagc	catccaccgc	1500
	acccaactct	ggttccacgg	gcgcatttcc	cgtgaggaga	gccagcgggt	tattggacag	1560
30	cagggccttg	tagacggcct	gttctctggt	cgggagagtc	agcggaaacc	ccagggtctt	1620
	gtcctctctt	tgtgccacct	gcagaaagtg	aagcattatc	tcctcctgcc	gagcgaggag	1680
	gagggtcgcc	tgtacttcag	catggatgat	ggccagaccc	gcttcaactga	cctgctgcag	1740
	ctcgtggagt	tcaccagct	gaaccggggc	atcctgcctg	gcttgcctgc	ccattgctgc	1800
	acgcgggtgg	ccctctgacc	aggccgtgga	ctggctcatg	cctcagcccg	ccttcaggct	1860
	gcccgcggcc	cctccaccca	tcagtggtac	tctggggcgc	ggccacaggg	gacgggatga	1920
35	ggagcgggag	ggttccgcca	ctccagtttt	ctcctctgct	tctttgcctc	cctcagatag	1980
	aaaacagccc	ccactccagt	ccactcctga	cccctctcct	caaggggaag	ccttgggtgg	2040
	ccccctctcc	ttctcctagc	tctggaggtg	ctgctctagg	gcaggggaatt	atgggagaag	2100
	tgggggcagc	ccaggcggtt	tcacgcccc	cactttgtac	agaccgagag	gccagttgat	2160
	ctgctctggt	ttatactagt	gacaataaag	attatttttt	gatac		2205
40	<210> 12						
	<211> 2177						
	<212> DNA						
45	<213> Homo sapiens						
50	<400> 12						
	gaattcgagg	ccgttggttt	gcagctgctc	cgctcatcgt	cggcccgaag	ctatctcgcg	60
	ctcgtgtgca	ggccgggtc	ggctcctggt	ccccggtgcg	agggttaacg	cgaggccccg	120
	gcctcggtcc	ccggactagg	ccgtgacccc	gggtgccatg	aagcaggagg	gctcggcgcg	180
	gcgcgcgggc	gcggacaagg	cgaaaaccgc	gcccggcgga	ggagaacaag	aacccccacc	240
	gccgcgggcc	ccccaggatg	tggagatgaa	agaggaggca	gcgacgggtg	gcgggtcaac	300
55	gggggaggca	gacggcaaga	cggcggcggc	agcgggttag	cactcccagc	gagagctgga	360

	cacagtcacc	ttggaggaca	tcaaggagca	cgtgaaacag	ctagagaaag	cggtttcagg	420
	caaggagccg	agattcgtgc	tgcggggcct	goggatgctg	ccttccacat	cacggccgct	480
	caaccactat	gttctgtata	aggctgtgca	gggcttcttc	acttcaaata	atgccactcg	540
5	agactttttg	ctcccccttc	tggaaagagcc	catggacaca	gaggctgatt	tacagttccg	600
	tccccgcacg	ggaaaagctg	cgtcgacacc	cctcctgcct	gaagtggag	cctatctcca	660
	actcctcgtg	gtcatcttca	tgatgaacag	caagcgctac	aaagaggcac	agaagatctc	720
	tgatgatctg	atgcagaaga	tcagtactca	gaaccgccgg	gccctagacc	ttgtagccgc	780
	aaagtgttac	tattatcacg	cccgggtcta	tgagttcctg	gacaagctgg	atgtggtgcg	840
	cagcttcttg	catgctoggc	tcoggacagc	taogcttogg	catgacgcag	acgggcaggc	900
10	caocctgttg	aacctcctgc	tgcgggaatta	cctacactac	agcttgtacg	accaggctga	960
	gaagctggtg	tccaagtctg	tgttcccaga	gcaggccaac	aacaatgagt	gggccaggta	1020
	cctctactac	acaggggcgaa	tcaaagccat	gcagctggag	tactcagagg	cccggaagac	1080
	gatgaccaac	gcccttcgca	aggccccca	gcacacagct	gtcggcttca	aacagacggt	1140
	gcacaagctt	gtcatcgtgg	tggagctggt	gctgggggag	atccctgacc	ggctgcagtt	1200
15	ccgccagccc	tccctcaagc	gctcaactcat	gccctatttc	cttctgactc	aagctgtcag	1260
	gacaggaaac	ctagccaagt	tcaaccaggt	cctggatcag	tttggggaga	agtttcaagc	1320
	agatgggacc	tacaccctaa	ttatccggct	gcggcacaac	gtgattaaga	cagggtgtacg	1380
	catgatcagc	ctctcctatt	cccgaatctc	cttggtgac	atcgcccaga	agctgcagtt	1440
	ggatagcccc	gaagatgcag	agttcattgt	tgccaaggcc	atccgggatg	gtgtcattga	1500
	ggccagcatc	aaccacgaga	agggctatgt	ccaatccaag	gagatgattg	acatctattc	1560
20	cacccgagag	ccccagctag	ccttccacca	gcgcctctcc	ttctgcctag	atatccacaa	1620
	catgtctgtc	aaggccatga	ggtttcctcc	caaatcgtac	aacaaggact	tggagtctgc	1680
	agaggaacgg	cgtgagcgag	aacagcagga	cttggagttt	gccaaaggaga	tggcagaaga	1740
	tgatgatgac	agcttccctt	gagctggggg	gctggggagg	ggtaggggga	atggggacag	1800
	gctctttccc	ccttgggggt	cccctgccca	gggcactgtc	cccattttcc	cacacacagc	1860
25	tcatatgctg	cattcgtgca	gggggtgggg	gtgctgggag	ccagccaccc	tgacctcccc	1920
	cagggtcctt	ccccagccgg	tgacttactg	tacagcaggc	aggaggggtg	gcaggcaacc	1980
	tccccgggca	gggtcctggc	cagcagtggt	ggagcaggag	gggaaggata	gttctgtgta	2040
	ctccttttag	gagtggggga	ctagaactgg	gatgtcttgg	cttgtatgtt	ttttgaagct	2100
	togattatga	tttttaacaa	ataaaaagtt	ctcccaaaaa	aaaaaaaaaa	aaaaaaaaaa	2160
30	aaagcggccg	cgaattc					2177

<210> 13

<211> 2960

35 <212> DNA

<213> Homo sapiens

40 <400> 13

	ctgccgcttc	caggcgtcta	tcagoggctc	agcctttggt	cagctgttct	gttcaaacac	60
	tctggggcca	ttcaggcctg	ggtggggcag	cgggaggaag	ggagtttgag	gggggcaagg	120
	cgacgtcaaa	ggaggtacag	agattccaca	atttcacaaa	actttcgcaa	acagcttttt	180
	gttccaaccc	ccctgcattg	tottggacac	caaatttgca	taaatcctgg	gaagttatta	240
45	ctaagcctta	gtcgtggccc	caggtaattt	cctcccaggc	ctccatgggg	ttatgtataa	300
	agggccccc	agagctgggc	cccaaaacag	cccggaagct	gcagcccagc	cccacccaga	360
	cccatggctg	gacctgccac	ccagagcccc	atgaagctga	tgggtgagtg	tcttggccca	420
	ggatgggaga	gccgcctgcc	ctggcatggg	agggaggctg	gtgtgacaga	ggggctgggg	480
	atccccgttc	tgggaatggg	gattaaaggc	accagtgctc	cccgaagagg	cctcaggtgg	540
50	tagggaacag	catgtctcct	gagcccgctc	tgccccagc	cctgcagctg	ctgctgtggc	600
	acagtgcact	ctggacagtg	caggaagcca	ccccctggg	ccctgccagc	tccttgcccc	660
	agagcttctt	gctcaagtgc	ttagagcaag	tgaggaagat	ccagggcgat	ggcgagcgcc	720
	tccaggagaa	gctggtgagt	gaggtgggtg	agagggctgt	ggaggggaag	ccgggtgggga	780
	gagctaaggg	ggatggaact	gcagggccaa	catcctctgg	aagggacatg	ggagaatatt	840
	aggagcagtg	gagctgggga	aggctgggga	gggacttggg	gaggaggacc	ttggtgggga	900
55	cagtgcctcg	gagggctggc	tgggatggga	gtggaggcat	cacattcagg	agaaagggca	960

agggccctg tgagatcaga gagtgggggt gcagggcaga gaggaactga acagcctggc 1020
 aggacatgga gggaggggaa agaccagaga gtcggggagg acccgggaag gagcggcgac 1080
 ccggccacgg cgagtctcac tcagcatcct tccatcccca gtgtgccacc tacaagctgt 1140
 5 gccacccoga ggagctggtg ctgctcggac actctctggg catcccctgg gctcccctga 1200
 gcagctgccc cagccaggcc ctgcagctgg tgagtgtcag gaaaggataa ggctaattgag 1260
 gagggggaag gagaggagga acaccatgg gctcccccatt gtctccagggt tccaagctgg 1320
 gggcctgacg tatctcaggc agcaccacct aactottccg ctctgtctca caggcaggct 1380
 gcttgagcca actccatago ggccttttcc totaccaggg gctcctgcag gccctggaag 1440
 10 ggatctcccc cgagttgggt cccaccttgg acacactgca gctggacgtc gccgaacttg 1500
 ccaccacat ctggcagcag gtgagccttg ttgggcaggg tggccaagggt cgtgctggca 1560
 ttctgggcac cacagccggg cctgtgtatg ggccctgtcc atgtgtcag cccccagcat 1620
 ttctcattt gtaataacgc cactcagaa gggcccaacc actgatcaca gctttccccc 1680
 acagatggaa gaactgggaa tggcccctgc cctgcagccc acccaggggtg ccatgccggc 1740
 cttoecctct gcttccagc gccgggcagg aggggtcctg gttgcctccc atctgcagag 1800
 15 cttoctggag gtgtcgtacc gcgttctacg ccaocttgcc cagccctgag ccaagccctc 1860
 cccatcccat gtatttatot ctatttaata tttatgtcta ttaagcctc atatttaaag 1920
 acaggggaaga gcagaacgga gcccaggcc tctgtgtcct tccctgcatt tctgagtttc 1980
 attctcctgc ctgtagcagt gagaaaaagc tcctgtcctc ccatcccctg gactgggagg 2040
 tagataggt aataccaagt atttattact atgactgtc cccagccctg gctctgcaat 2100
 20 gggcactggg atgagccgct gtgagccctt ggtcctgagg gtccccacct gggacccttg 2160
 agagtatcag gtctcccaag tgggagacaa gaaatccctg ttaataattt aaacagcagt 2220
 gttocccac tgggtccttg caccctcac tctggcctca gccgactgca cagcggcccc 2280
 tgcatccctt tggctgtgag gcccctggac aagcagaggt gccagagct gggaggcatg 2340
 gccctggggt cccacgaatt tgctggggaa tctogttttt cttcttaaga cttttgggac 2400
 25 atggtttgac tcccgaaat caccgacgtg tctcctgttt ttctgggtgg cctcgggaca 2460
 cctgcctgc cccacgagg gtcaggactg tgactctttt tagggccagg cagggtgcctg 2520
 gacatttgcc ttgctggatg gggactgggg atgtgggagg gagcagacag gaggaatcat 2580
 gtcaggcctg tgtgtgaaag gaagctccac tgtoacccct caoctottca cccccactc 2640
 accagtgtcc cctccactgt cacttgtaa ctgaacttca ggataataaa gtgtttgcct 2700
 30 ccagtcacgt ccttccctct tcttgagtc agctggtgcc tggccagggg ctggggagggt 2760
 ggctgaaggg tgggagaggc cagaggaggg tggggaggga ggtctgggga ggaggtccag 2820
 ggaggaggag gaaagttctc aagttcgtct gacattcatt ccgttagcac atatttatct 2880
 gagcacctac tctgtgcaga cgctgggcta agtgotgggg acacagcagg gaacaaggca 2940
 gacatggaat ctgcactoga 2960

35 <210> 14
 <211> 850
 <212> DNA
 40 <213> Homo sapiens

45 <220>
 <221> misc_feature
 <222> (3) .. (4)
 50 <223> n=a, c, g or t

55

<220>

<221> misc_feature

<222> (9) .. (9)

<223> n=a, c, g or t

<220>

<221> misc_feature

<222> (11) .. (11)

<223> n=a, c, g or t

<220>

<221> misc_feature

<222> (18) .. (18)

<223> n=a, c, g or t

<220>

<221> misc_feature

<222> (202) .. (202)

<223> n=a, c, g or t

<220>

<221> misc_feature

<222> (205) .. (205)

<223> n=a, c, g or t

<220>

<221> misc_feature

<222> (273) .. (273)

<223> n=a, c, g or t

5
 10
 15
 20
 25
 30
 35
 40
 45
 50
 55

<220>
 <221> misc_feature
 <222> (327) .. (327)
 <223> n=a, c, g or t

<220>
 <221> misc_feature
 <222> (367) .. (367)
 <223> n=a, c, g or t

<220>
 <221> misc_feature
 <222> (581) .. (581)
 <223> n=a, c, g or t

<220>
 <221> misc_feature
 <222> (599) .. (599)
 <223> n=a, c, g or t

<220>
 <221> misc_feature
 <222> (628) .. (628)
 <223> n=a, c, g or t

<220>
 <221> misc_feature
 <222> (673) .. (673)
 <223> n=a, c, g or t

<220>

<221> misc_feature

<222> (675) .. (675)

<223> n=a, c, g or t

<220>

<221> misc_feature

<222> (682) .. (682)

<223> n=a, c, g or t

<220>

<221> misc_feature

<222> (693) .. (693)

<223> n=a, c, g or t

<220>

<221> misc_feature

<222> (698) .. (698)

<223> n=a, c, g or t

<220>

<221> misc_feature

<222> (700) .. (700)

<223> n=a, c, g or t

<220>

<221> misc_feature

<222> (720) .. (720)

<223> n=a, c, g or t

<220>

<221> misc_feature

<222> (730)..(730)

<223> n=a, c, g or t

<220>

<221> misc_feature

<222> (734)..(734)

<223> n=a, c, g or t

<220>

<221> misc_feature

<222> (742)..(743)

<223> n=a, c, g or t

<220>

<221> misc_feature

<222> (746)..(746)

<223> n=a, c, g or t

<220>

<221> misc_feature

<222> (748)..(748)

<223> n=a, c, g or t

<220>

<221> misc_feature

<222> (752)..(752)

<223> n=a, c, g or t

5
 <220>
 <221> misc_feature
 <222> (762) .. (762)
 <223> n=a, c, g or t

10
 <220>
 <221> misc_feature
 <222> (767) .. (767)
 <223> n=a, c, g or t

20
 <220>
 <221> misc_feature
 <222> (777) .. (777)
 <223> n=a, c, g or t

30
 <220>
 <221> misc_feature
 <222> (783) .. (784)
 <223> n=a, c, g or t

35
 <220>
 <221> misc_feature
 <222> (789) .. (789)
 <223> n=a, c, g or t

45
 <220>
 <221> misc_feature
 <222> (794) .. (794)
 <223> n=a, c, g or t

55

<220>

<221> misc_feature

<222> (797) .. (798)

<223> n=a, c, g or t

<220>

<221> misc_feature

<222> (803) .. (805)

<223> n=a, c, g or t

<220>

<221> misc_feature

<222> (810) .. (810)

<223> n=a, c, g or t

<220>

<221> misc_feature

<222> (817) .. (817)

<223> n=a, c, g or t

<220>

<221> misc_feature

<222> (826) .. (827)

<223> n=a, c, g or t

<220>

<221> misc_feature

<222> (831) .. (832)

<223> n=a, c, g or t

<220>

<221> misc_feature

<222> (834)..(834)

<223> n=a, c, g or t

<220>

<221> misc_feature

<222> (837)..(838)

<223> n=a, c, g or t

<220>

<221> misc_feature

<222> (840)..(840)

<223> n=a, c, g or t

<220>

<221> misc_feature

<222> (844)..(844)

<223> n=a, c, g or t

<220>

<221> misc_feature

<222> (846)..(848)

<223> n=a, c, g or t

<400> 14

ttnnctttnt	ngccatgncc	agttcaactc	agcctctcag	ttccacacgg	acaacatgog	60
ggaccctctg	aaccgagtc	tggccaacot	gttctctctc	atctcctcca	tcctggggtc	120
tcgcacogct	ggccccaca	cccagttcgt	gcagtgggtc	atggaggagt	gtgtggactg	180
cctggagcag	ggtggccgtg	gnagngtcct	gcagttcatg	cccttcacca	ccgtgtcgga	240
actggtgaag	gtgtcagcca	tgtctagccc	canggtggtt	ctggccatca	cggacotcag	300
cctgccccctg	ggccgccagg	tggtcgtntaa	agccattgct	gcactctgag	gggcttggca	360
tggccgnagt	gggggctggg	gactggcgca	gccccaggcg	cctccaaggg	aagcagttag	420
gaaagatgag	gcctcgtgcc	tcacatccgt	tcacatgggt	gcaagagcct	ctagcggcctt	480
ccagttcccc	gctctgact	cctgactcca	ggatgtctcc	cggtttcttc	ttttcaaaat	540
tttctctctc	atcttgctgg	caactgagga	gagttagcag	nctggaccac	aagcccagng	600
ggtcacccct	gtgttgctgg	cgcccagncc	aggagtagtc	ttacctcttg	aggaactttc	660
ttggatggaa	agnnggtttt	tntgtgttgt	gtntgtgnan	gtgtttttcg	gggttttttn	720
gggcaatatn	ttangggaat	cnnccntncc	cncatttttt	cnttagagct	ccccggngga	780

aanntcttna tccnctnnct ttnnnctecn tcacctnctt totttnttct nntnttnnen 840
tccnccnncc 850

<210> 15

<211> 2309

<212> DNA

<213> Homo sapiens

<400> 15

ccccgggcgc	aggaggcggg	cgccccggcc	ccaccggccc	cccatggaag	ccccagcac	60
ggggcgctga	gacccccgcg	tcgctgcccc	gcccggtccg	gogcgccacg	ccagggatct	120
ctggacagga	caagactccg	aagctactcc	cccagcacac	agcccgggac	ccacaaaccc	180
agcttgcccc	cagccctccc	acctgccact	ccctggcccc	tcccaccgcc	cgccccctt	240
ggggcgccgc	gcatggtgtg	aaaggccaag	tgctgaggcg	ggtatcatgg	gtgctgtgcc	300
ctaggggcctg	ggtggcaggg	ggtgggtggc	ctgtgggtgt	gcccgggggg	ccagtgtgcc	360
cacccocagtc	tcttggcgtg	ctggaggggca	tcctggatgg	aattgaagtg	aatggaacag	420
aagccaagca	aggtggagtg	tgggtcagac	ccagaggaga	acagtgccag	gtcaccagat	480
ggaaagcgaa	aaagaaagaa	cggccaatgt	tcctgaaaa	ccagcatgtc	aggttatatc	540
cctagttacc	tgacaaaaga	cgagcagtg	gtcgtgtgtg	gggacaaggc	aactggttat	600
cactaccgct	gtatcacttg	tgagggctgc	aagggtctct	ttcgccgcac	aatccagaag	660
aacctccatc	ccacctattc	ctgcaaatat	gacagctgct	gtgtcattga	caagatcacc	720
cgcaatcagt	gccagctgtg	ccgcttcaag	aagtgcacgt	ccgtgggcat	ggccatggac	780
ttggtttotag	atgactcgaa	gcgggtggcc	aagcgttaag	tgattgagca	gaaccgggag	840
cgggcgogga	aggaggagat	gatccgatca	ctgcagcagc	gaccagagcc	cactcctgaa	900
gagtgggatc	tgatccacat	tgccacagag	gccatcgcga	gcaccaatgc	ccagggcagc	960
cattggaaac	agaggcggaa	attcctgccc	gatgacattg	gccagtcacc	cattgtctcc	1020
atgccggacg	gagacaaggt	ggacctggaa	gccttcagcg	agtttaacca	gatcatcacc	1080
ccggccatca	cccggtgtgt	ggactttgct	aaaaaactgc	ccatgttctc	cgagctgcct	1140
tgccgaagac	agatcatctc	octgaagggg	tgctgcatgg	agatcatgtc	cctgcgggcg	1200
gctgtccgct	acgacctga	gagcgacacc	ctgacgctga	gtggggagat	ggctgtcaag	1260
cgggagcagc	tcaagaatgg	cgccctgggc	gtagtctccg	acgccatctt	tgaactgggc	1320
aagtcactct	ctgcctttaa	cctggatgac	acggaagtgg	ctctgctgca	ggctgtgctg	1380
ctaattgtcaa	cagaccgctc	gggcctgctg	tgtgtggaca	agatcgagaa	gagtcaggag	1440
gcgtacctgc	tgccgttcga	gcactacgtc	aaccaccgca	aacacaacat	tccgcacttc	1500
tgccccaagc	tgctgatgaa	ggagagagaa	gtgcagagtt	cgattctgta	caagggggca	1560
gcggcagaag	gccggccggg	cggttcactg	ggcgtccacc	cggaaggaca	gcagcttctc	1620
ggaatgcatg	ttgttcaggg	tcgcaggttc	cggcagcttg	agcagcagct	tggtgaagcg	1680
ggaagtctcc	aagggccggt	tcttcagcac	cagagcccga	agagcccgca	gcagcgtctc	1740
ctggagctgc	tccaccgaag	cggaattctc	catgcccag	cggtctgtgg	ggaagacgac	1800
agcagtgagg	cggactcccc	gagctcctct	gaggaggaac	cggaggtctg	cgaggacctg	1860
gcaggcaatg	cagcctctcc	ctgaagcccc	ccagaaggcc	gatggggaag	gagaaggagt	1920
gccatacctt	ctcccaggcc	tctgccccaa	gagcaggagg	tgccgtgaaag	ctgggagcgt	1980
gggctcagca	gggctggtca	cctcccatcc	cgtaagacca	cctccctctc	ctcagcaggc	2040
caaacatggc	cagactccct	tgctttttgc	tgtgtagtgc	cctctgcctg	ggatgccctt	2100
ccccctttct	ctgcctggca	acatcttact	tgtcctttga	ggccccaact	caagtgtcac	2160
ctccttcccc	agctccccca	ggcagaaata	gttgtctgtg	cttccttggt	tcatgcttct	2220
actgtgacac	ttatctcact	gttttataat	tagtcgggca	tgagtctgtt	tcccaagcta	2280
gactgtgtct	gaatcatgtc	tgtatcccc				2309

<210> 16

<211> 2355

<212> DNA

<213> Homo sapiens

<400> 16

ccgttgcc	aacgtccaac	ccttctgcag	ggctgcagtc	cgccacccc	aagaccttgc	60
tgcagggtgc	ttcggatcct	gatogtgagt	cgcggggtcc	actccccgcc	cttagccagt	120
gcccaggggg	caacagcggc	gatogcaacc	tctagtttga	gtcaagggtcc	agtttgaatg	180
accgctctca	gctgggtgaag	acatgaccac	cctggactcc	aacaacaaca	caggtggcgt	240
catcacctac	attggctcca	gtggctcttc	ccaagccgc	accagccctg	aatccctcta	300
tagtgacaac	tccaatggca	gcttccagtc	cctgacccaa	ggctgtccca	cctacttccc	360
accatccccc	actggctccc	tcacccaaga	ccggctcgc	tcctttggga	gcattccacc	420
cagcctgagt	gatgacggct	ccccttcttc	ctcatcttcc	togtgcgcat	cctcctcttc	480
cttctataat	gggagccccc	ctgggagttc	acaagtggcc	atggaggaca	gcagccgagt	540
gtcccccagc	aagagcacca	gcaacatcac	caagtgaat	ggcatgggtg	tactgtgtaa	600
agtgtgtggg	gacgttgcc	cggtcttcca	ctacgggtgtg	ctcgccctgcg	agggctgcaa	660
gggttttttc	cgtcggagca	tccagcagaa	tccacgtac	aaaaggtgtc	tgaagaatga	720
gaattgtctc	atcgtccgca	tcaatcgcaa	ccgctgccag	caatgtcgct	tcaagaagtg	780
tctctctgtg	ggcatgtctc	gagaogctgt	gcgttttggg	cgcatcccca	aacgagagaa	840
gcagcggatg	cttgctgaga	tgcagagtgc	catgaacctg	gccaacaacc	agttgagcag	900
ccagtgcocg	ctgggagactt	cacccaccca	gcaccccaac	ccaggcccca	tgggcccctc	960
gccacccccc	gctccgggtc	cctcaccctc	ggtgggtctc	tcccagtttc	cacaacagct	1020
gacgcctccc	agatccccc	gccctgagcc	cacagtggag	gatgtgatat	cccaggtggc	1080
cogggcccac	cgagagatct	tcacctacgc	ccatgacaag	ctgggcagct	cacctggcaa	1140
cttcaatgoc	aaccatgcat	caggtagccc	tccagccacc	acccacatc	gctgggaaaa	1200
tcagggtgtg	ccacctgccc	ccaatgacaa	caacaccttg	gctgcccagc	gtcataacga	1260
ggccctaaat	ggtctgcgcc	aggctccctc	ctcctaccct	cccacctggc	ctcctggccc	1320
tgcacaccac	agctgccacc	agtccaacag	caacgggcac	ogtctatgoc	ccacccacgt	1380
gtatgcagcc	ccagaaggca	aggcacctgc	caacagtccc	cggcagggca	actcaaagaa	1440
tgttctgctg	gcatgtccta	tgaacatgta	ccgcctatga	cgcagtgggc	gaacggtgca	1500
ggagatctgg	gaggatttct	ccatgagctt	cacgcccgtc	gtgcgggagg	tggttagagtt	1560
tgccaaacac	atcccgggct	tccgtgacct	ttctcagcat	gaccaagtca	ccctgcttaa	1620
ggctggcacc	tttgaggtgc	tgatggtgcg	ctttgcttgc	ttgttcaacg	tgaaggacca	1680
gacagtgatg	ttcctaagcc	ggaccaccta	cagcctgcag	gagcttggtg	ccatgggcat	1740
gggagacctg	ctcagtgcc	tgttgcactt	cagcgagaag	ctcaactccc	tggcgcttac	1800
cgaggaggag	ctgggcctct	tcacccgggt	ggtgcttgct	tctgcagacc	gctcgggcat	1860
ggagaattcc	gcttcggtgg	agcagctcca	ggagacgctg	ctgcgggctc	ttcgggctct	1920
ggtgctgaag	aaccggccct	tggagacttc	ccgcttcacc	aagctgctgc	tcaagctgcc	1980
ggacctggcg	accctgaaca	acatgcattc	cgagaagctg	ctgtccttcc	gggtggacgc	2040
ccagtgaacc	gcccggccgg	ccttctgccc	ctgccccctt	gtacagaatc	gaactctgca	2100
cttctctctc	ctttacgaga	cgaaaaggaa	aagcaaacca	gaatcttatt	tatatgttta	2160
taaaatattc	caagatgagc	ctctggcccc	ctgagccttc	ttgtaaatac	ctgcctccct	2220
cccccatcac	cgaacttccc	ctcctccctc	atttaaacca	ctctgtctcc	cccacaaccc	2280
tcccctggcc	ctctgatttg	ttctgttctc	gtctcaaatc	caatagttca	cagctaaaaa	2340
aaaaaaaaaa	aaaag					2355

<210> 17

<211> 4119

<212> DNA

<213> Homo sapiens

<400> 17

gaattccggtt	gctgtcgcac	acacacacac	acacacacac	acacacacac	acacacacac	60
acaccccaac	acacacacac	acacacacac	acacacacac	acacacacac	acacacacac	120
atggccgagc	gccgcacgcg	tagcacgcgc	ggactagcta	tccagcctcc	cagcagcctc	180
tgcgacgggc	gcggtgcgta	agtaacctgc	cggtggtggc	cgttctccgt	aagatggcgg	240
acogggcggc	gcagcgcgct	tcgcaagaca	ccgaggacga	ggaatctggt	gcttcgggct	300
ccgcacgcgg	cggctccccg	ttgcggggag	gcgggagctg	cagcggtagc	gccggaggcg	360
goggcagcgg	ctctctgcct	tcacagcgcg	gaggccgaac	cggggccctt	catctgcggc	420
gggtggagag	cgggggcggc	aagagtgcctg	aggagtccga	gtgtgagagt	gaagatggca	480
ttgaaggtga	tgctgtttct	tcggattatg	aaagtgcaga	agactcggaa	ggtgaagaag	540
gtgaatacac	tgaagaggaa	aactccaaag	tggagctgaa	atcagaagct	aatgatgctg	600
ttaattcttc	aacaaaagaa	gagaaggagg	tgaaaaagcc	tgacacccaa	agcaotgtga	660
ctggagagag	gcaaagtggg	gacggacagg	agagcacaga	gcctgtggag	aacaaagtgg	720
gtaaaaaggg	ccctaagcat	ttggatgatg	atgaagatcg	gaagaatcca	gcatacatat	780
ctcggaaagg	gctcttcttt	gagcatgatc	ttcggaggga	aactcaggag	gaggaagtca	840
gacccaaggg	gcgtcagcga	aagctatgga	aggatgaggg	tcgctgggag	catgacaagt	900
tccgggaaga	tgagcaggcc	ccaaagtccc	gacaggagct	catgtctctt	tatggttatg	960
acattcgctc	agctcataat	cctgatgaca	tcaaacctcg	aagaatcccg	aaaccccgat	1020
atgggagatc	tccacaaaga	gatccaaact	ggaacgggtg	gcggctaaac	aagtctcatc	1080
gccaccaggg	tcttgggggc	accctaccac	caaggacatt	tattaacagg	aatgtctgag	1140
gtaccggccg	tatgtctgca	cccaggaatt	attctcgatc	tgggggcttc	aaggaaggtc	1200
gtgctgggtt	taggcctgtg	gaagctgggt	ggcagcatgg	tggccggctc	ggtgagactg	1260
ttaagcatga	gattagttac	cggtoacggc	gcctagagca	gaattctgtg	agggatccat	1320
ctocagaagc	agatgctcca	gtgcttgcca	gtcctgagaa	ggaagaggca	gcctcagagc	1380
caccagctgc	tgctcctgat	gtgcaccac	cacccctga	taggcccatt	gagaagaaat	1440
cctattcccc	ggcaagaaga	actogaacca	aagttggaga	tcagtcgaag	cttgcagagg	1500
agggtgcccc	tcctcctgaa	ggactgattc	cagcacctcc	agtcccagaa	accaccccaa	1560
ctccacctac	taagactggg	acctgggaag	ctccggtgga	ttctagtaca	agtggaactt	1620
agcaagatgt	ggcacaacta	aatatagcag	aacagaattg	gagtcggggg	cagccttctt	1680
tcctgcaacc	acgggaactt	cagggtatgc	ccaaccatat	acacatggga	gcaggacctc	1740
cacctcagtt	taaccggatg	gaagaaatgg	gtgtccaggg	tggctcagcc	aaacgctatt	1800
catcccagcg	gcaaagacct	gtgccagagc	cccccgcccc	tccagtgcac	atcagtatca	1860
tggaggggaca	ttactatgat	ccactgcagt	tccaggggacc	aatctatacc	catggtgaca	1920
gccctgcccc	gctgcctcca	cagggcagtc	ttgtgcagcc	aggaatgaac	cttccccacc	1980
cagggtttaca	tccccaccag	acaccagctc	ctctgcccc	tccaggcctc	tatccccacc	2040
cagtgtccat	gtctccagga	cagccaccac	ctcagcagtt	gcttgcctct	acttactttt	2100
ctgctccagg	cgctcatgaa	tttggtatc	ccagttaccc	ttatgctcca	ggggcactgc	2160
ctccccacc	acogcctcat	ctgtatccta	atacacaggc	cccatcacag	gtatatggag	2220
gagtgcacta	ctataacccc	gccagcagc	agggtgcagc	aaagccctcc	ccaccccgga	2280
ggactcccca	gccagtcacc	atcaagcccc	ctccacctga	ggttgtaagc	aggggttcca	2340
gttaatacaa	gtttctgaat	attttaaatc	ttaacatcat	ataaaaagca	gcagagggtga	2400
gaactcagaa	gagaaataca	gctggctatc	tactaccaga	agggttcaa	agatataggg	2460
tgtggctcct	accagcaaac	agctgaaaga	ggaggacccc	tgcttctctc	tgaggacagg	2520
ctctagagag	agggagaaac	aagtggacct	cgtcccatct	tcactcttca	cttgagttgg	2580
ctgtgttcgg	gggagcagag	agagccagac	agccccaagc	ttctgagctc	agatacagaa	2640
gcccattgtct	totgtgttct	ttcaactctg	ggaaattgaa	gtgtcttctg	ttcccaagga	2700
agctccttcc	tgtttgtttt	gtttttctaag	atgttccattt	ttaaagcctg	gcttcttctc	2760
cttaatatatta	ttttaatttt	ttctctttgt	ttctgtttct	tgtctctctc	cctgcctttt	2820
aaatgaaaca	agtctagctc	tctgggtttc	tagccctctc	ggattccctt	ttgactcttc	2880

	cgtgcacccc	agataatgga	gaatgtatca	gccagccttc	cccaccaagt	ctaaaaagac	2940
	ctggcccttc	acttttagtt	ggcatttggt	atcctcttgt	atacttgtag	tcccttaact	3000
	ctaaccctgt	ggaagcatgg	ctgtctgcac	agagggtccc	attgtgcaga	aaagctcaga	3060
5	gtagggtgggt	aggagccctt	ctotttgact	taggttttta	ggagtctgag	catccatcaa	3120
	tacctgtact	atgatgggct	tctgttctct	gctgagggcc	aataccctac	tgtggggaga	3180
	gatggcacac	cagatgcttt	tgtgagaaag	ggatggtgga	gtgagagcct	ttgcctttag	3240
	gggtgtgtat	tcacatagtc	ctcagggctc	agtcttttga	ggtaagtgga	attagagggc	3300
	cttgcttctc	ttctttccat	tottcttgtc	acaccccttt	tccagttgct	gtggaccaat	3360
10	gcactctctt	aaaggcaaat	attatccagc	aagcagtcct	ccctgtcctt	tgcaattgct	3420
	cttctccacg	tctttcctgc	tacaagtgtt	ttagatgtta	ctaccttatt	ttccccgaat	3480
	totatttttg	tccttgacga	cagaatataa	aaactcctgg	gcttaaggcc	taagggaagc	3540
	agtcaccttc	tgggcaagg	ctcctatctt	tcctccctat	ccatggcact	aaaccacttc	3600
	tctgctgcct	ctgtggaaga	gattcctatt	actgcagtac	atacgtctgc	caggggtaac	3660
	ctggccactg	tcctgtcct	totacagaac	ctgagggcaa	agatggtggc	tgtgtctctc	3720
15	cccggtaatg	tcactgtttt	tattccttcc	atctagcagc	tggcctaato	actctgagtc	3780
	acaggtgttg	gatggagagt	ggggagaggc	acttaatctg	taacccccaa	ggaggaaata	3840
	actaagagat	tottctaggg	gtagctgggt	gttgtgcctt	ttgtaggctg	ttccctttgc	3900
	otaaacctg	aagatgtctc	ctcaagcctg	tgggcagcat	gccagatttc	ccagacctta	3960
	agacactgtg	agagttgtct	ctgttggctc	actgtgttta	gttgcaagga	tttttccatg	4020
20	tgtggtggtg	ttttttgtta	ctgttttaaa	gggtgcccac	ttgtgatcag	cattgtgact	4080
	tggagataat	aaaatttaga	ctataaaact	gaaaaaaaa			4119
	<210> 18						
	<211> 2653						
25	<212> DNA						
	<213> Homo sapiens						
30	<400> 18						
	gagcgcggtg	ggagtttget	gctgccgctg	tgcagtttgt	tcaggggctt	gtggcggtga	60
	gtccgagagg	ctgcgtgtga	gagacgtgag	aaggatcctg	cactgaggag	gtggaaagaa	120
	gaggattgct	cgaggaggcc	tggggctctgt	gagacagcgg	agctgggtga	aggctgcggg	180
35	ttccggcgag	gcctgagctg	tgtgtctgtc	atgctcctaa	cccgatccca	ggcacaggct	240
	acaatcagtt	ttocaaaaag	gaagctgtct	cgggcattga	acaaagctaa	aaactccagt	300
	gatgccaaac	tagaaccaac	aatgttccaa	acogtaacct	gttctcctcg	tgtaaaagcc	360
	ctgcctctca	gccccaggaa	acgtctgggc	gatgacaacc	tatgcaacac	tccccattta	420
	cctccttggt	ctccaccaaa	gcaaggcaag	aaagagaatg	gtcccccctca	ctcacatata	480
40	cttaagggac	gaagattggt	atttgacaat	cagctgacaa	ttaagtctcc	tagcaaaaga	540
	gaactagcca	aagttcacca	aaacaaaata	ctttcttcag	ttagaaaaag	tcaagagatc	600
	acaacaaatt	ctgagcagag	atgtccactg	aagaaagaat	ctgcatgtgt	gagactatto	660
	aagcaagaag	gcacttgcta	ccagcaagca	aagctggtcc	tgaacacagc	tgtcccagat	720
	cggtgcctg	ccagggaag	ggagatggat	gtcatcagga	atttcttgag	ggaacacatc	780
	tgtgggaaaa	aagctggaag	cotttacctt	tctgggtgct	ctggaactgg	aaaaactgcc	840
45	tgtttaagcc	ggattctgca	agacctcaag	aaggaaactga	aaggctttaa	aactatcatg	900
	ctgaattgca	tgtccttgag	gactgccag	gctgtattcc	cagctattgc	tcaggagatt	960
	tgtcaggaag	aggtatccag	gccagctggg	aaggacatga	tgaggaaatt	ggaaaaacat	1020
	atgactgcag	agaagggcc	catgattgtg	ttggtattgg	acgagatgga	tcaactggac	1080
	agcaaggcc	aggatgtatt	gtacacgcta	tttgaatggc	catggctaag	caattctcac	1140
50	ttggtgtga	ttggtattgc	taataccctg	gatctcacag	atagaattct	acotaggctt	1200
	caagctagag	aaaaatgtaa	gccacagctg	ttgaacttcc	caocttatac	cagaaatcag	1260
	atagtacta	ttttgcaaga	tgcacttaac	caggtatcta	gagatcaggt	tctggacaat	1320
	gctgcagttc	aattctgtgc	cgcgaagtc	tctgctgttt	caggagatgt	tcgcaaagca	1380
	ctggatgttt	gcaggagagc	tattgaaatt	gtagagtcag	atgtcaaaag	ccagactatt	1440
55	ctcaaccac	tgtctgaatg	taaatcacct	tctgagcctc	tgattcccaa	gagggttggt	1500
	cttattcaca	tatcccaagt	catctcagaa	gttgatggta	acaggatgac	cttgagccaa	1560

	gagggagcac	aagattccctt	ccctcttcag	cagaagatct	tggtttgctc	tttgatgctc	1620
	ttgatcaggc	agttgaaaat	caaagaggtc	actctgggga	agttatatga	agcctacagt	1680
	aaagtctgtc	gcaaacagca	ggtggcggct	gtggaccagt	cagagtgttt	gtcactttca	1740
5	gggctcttgg	aagccagggg	catttttagga	ttaaagagaa	acaaggaaac	ccgtttgaca	1800
	aagggtgtttt	tcaagattga	agagaaagaa	atagaacatg	ctctgaaaga	taaagcttta	1860
	attggaaata	tcttagctac	tggattgcct	taaattcttc	tcttacaccc	caccogaaag	1920
	tattcagctg	gcatttagag	agctacagtc	ttcatttttag	tgctttacac	attcgggcct	1980
	gaaaacaaat	atgacctttt	ttacttgaag	ccaatgaatt	ttaatotata	gattctttaa	2040
10	tattagcaca	gaataatata	tttgggtctt	actattttta	cccataaaaag	tgaccaggta	2100
	gacctttttt	aattacatto	actacttcta	ccacttgtgt	atctctagcc	aatgtgcttg	2160
	caagtgtaca	gatctgtgta	gaggaaatgtg	tgtatattta	cctcttcggt	tgotcaaaca	2220
	tgagtgggta	tttttttggt	tgtttttttt	gttgttggtg	tttttgaggc	gcgtctcacc	2280
	ctgttgccca	ggctggagtg	caatggcgcg	ttctctgctc	actacagcac	ccgcttccca	2340
	gggtgaagtg	attctcttgc	ctcagcctcc	cgagttagctg	ggattacagg	tgcccaccac	2400
15	cgcgcccagc	taatttttta	atttttagta	gagacagggt	tttaccatgt	tgcccaggct	2460
	ggtcttgaac	tcctgacct	caagtgatct	gcccacottg	gcctccctaa	gtgctgggat	2520
	tataggcgtg	agccaccatg	ctcagccatt	aaggatattt	gttaagaact	ttaagttag	2580
	ggtaagaaga	atgaaaatga	tccagaaaaa	tgcaagcaag	tccacatgga	gatttgaggg	2640
	acactgggta	aag					2653
20	<210> 19						
	<211> 2907						
	<212> DNA						
25	<213> Homo sapiens						
30	<400> 19						
	gccatctggg	cccaggcccc	atgccccgag	gaggggtggt	ctgaagccca	ccagagcccc	60
	ctgcagact	gtctgcctcc	cttctgactg	tggccgcttg	gcatggccag	caacagcagc	120
	tcttgcccga	caactggggg	cgggcacctc	aatgggtacc	cgggtgcctcc	ctacgccttc	180
	ttcttccccc	ctatgetggg	tggactctcc	ccgccaggcg	ctctgaccac	tctccagcac	240
	cagcttccag	ttagtgata	tagcacacca	tccccagcca	ccattgagac	ccagagcagc	300
35	agttctgaag	agatagtgc	cagccctccc	tgcgcccccc	ctctaccccc	catctacaag	360
	ccttgctttg	tctgtcagga	caagtctcca	ggctaccact	atgggggtcag	cgctgtgag	420
	ggctgcaagg	gcttcttccg	ccgcagcctc	cagaagaaca	tggtgtacac	gtgtcacccg	480
	gacaagaact	gcattcatca	caaggtgacc	cggaaacgct	gccagtactg	ccgactgcag	540
	aagtgctttg	aagtgggcat	gtccaaggag	tctgtgagaa	acgaccgaaa	caagaagaag	600
40	aaggagggtg	ccaagcccga	gtgctctgag	agctacacgc	tgacgcocga	ggtgggggag	660
	ctcattgaga	aggtgcgcaa	agcgcaccag	gaaaccttcc	ctgcccctctg	ccagctgggc	720
	aaatacacta	cgaacaacag	ctcagaacaa	cgtgtctctc	tggacattga	cctctgggac	780
	aagttcagtg	aactctccac	caagtgcctc	attaagactg	tggagtctgc	caagcagctg	840
	cccggcttca	ccacctccac	catcgccgac	cagatcaccc	tcctcaaggc	tgctgccttg	900
45	gacatcctga	tcttgccgat	ctgcacgcgg	tacacgcccg	agcaggacac	catgaccttc	960
	tcggacgggc	tgacctgaa	coggaccccag	atgcacaacg	ctggcttcgg	ccccctcacc	1020
	gaactgggtc	ttgccttgc	caaccagctg	ctgcccctgg	agatggatga	tgccggagacg	1080
	gggctgctca	gcgccatctg	cctcatctgc	ggagaccgcc	aggacctgga	gcagccggac	1140
	cgggtggaca	tgtctcagga	gccctgctg	gaggcgctaa	aggtctacgt	gcggaagcgg	1200
	aggcccagcc	gccccacat	gttccccaa	atgctaatga	agattactga	cctgcgaagc	1260
50	atcagcggca	agggggctga	gcggtgatc	acgctgaaga	tggagatccc	gggctccatg	1320
	ccgcctctca	tccaggaaat	gttgagaaac	tcagagggcc	tggacactct	gagcggacag	1380
	ccgggggggtg	gggggcccga	cgggggtggc	ctggccccc	cggcaggcag	ctgtagcccc	1440
	agcctcagcc	ccagctccaa	cagaagcagc	ccggccaccc	actccccgtg	accgcccacg	1500
	ccacatggac	acagccctcg	ccctccggcc	cggcttttct	ctgcctttct	accgaccatg	1560
55	tgaccocgca	ccagccctgc	ccccacctgc	cctccggggc	agtactgggg	accttccctg	1620
	ggggacgggg	agggaggagg	cagcgactcc	ttggacagag	gcctggggccc	tcagtggact	1680

```

gectgctccc acagcctggg ctgacgtcag aggccgaggg caggaactga gtgaggcccc 1740
tggtcctggg tctcaggatg ggtcctgggg gcctcgtggt catcaagaca cccctctgcc 1800
cagctcacca catottcatc accagcaaac gccaggactt ggctccccc tcctcagaac 1860
5 tcacaagcca ttgtcccca gctggggaac ctcaacctcc cccctgcctc ggttggtgac 1920
agaggggggtg ggacaggggc ggggggttcc ccctgtacat accctgccat accaacccca 1980
ggtattaatt ctgctgggtt ttgtttttat ttttaatttt ttgttttgat ttttttaata 2040
agaattttca ttttaagcac atttatactg aaggaatttg tgctgtgtat tggggggagc 2100
tggatccaga gctggagggg gtgggtccgg gggaggagg ggctcggaag gggcccccac 2160
10 totcctttca tgtccctgtg ccccccagtt ctccctcctca gccttttctt cctcagtttt 2220
ctcttttaaaa ctgtgaagta ctaacttttc aaggcctgcc ttccctccc tccactgga 2280
gaagccgcca gcccctttct ccctctgcct gaccactggg tgtggaagg gtggggcagc 2340
cctgaaagga caggctcctg gccttggcac ttgcctgcac ccaccatgag gcatggagca 2400
gggcagagca agggcccccg gacagagttt tcccagacct ggctcctcgg cagagctgcc 2460
15 tccgctcagg gccacatca tctaggctcc ccagccccc ctgtgaagg gctggccagg 2520
ggcccagagc gcccccaccc cgggctcag ccaccagcac ccccataggg ccccagaca 2580
ccacacacat gcgctgctgc acacacacaa acacacacac actggacagt agatgggcoo 2640
acacacactt ggcccgagtt cctccatttc cctggcctgc ccccaacccc caacctgtcc 2700
caccctcgtg cccctcctt accccgcagg acgggcctac aggggggtct cccctcacco 2760
ctgcaccccc agctggggga gctggctctg ccccgacctc cttcaccagg ggttggggcc 2820
20 ccttcocctg gagcccggtg gtgcacctgt tactgttggg ctttccactg agatctactg 2880
gataaagaat aaagttctat ttattct 2907

```

<210> 20

<211> 2096

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<222> (23) .. (23)

<223> n=a, c, g or t

<220>

<221> misc_feature

<222> (27) .. (27)

<223> n=a, c, g or t

<220>

<221> misc_feature

<222> (80) .. (80)

<223> n=a, c, g or t

<220>

<221> misc_feature

<222> (120)..(120)

<223> n=a, c, g or t

<400> 20

agatgtttta	aaatactttg	atnctcngtt	tccaacctctc	ttaaattgtc	tttccctatg	60
ttaaataatac	agtcatacnc	ttgctgaaaa	aagttcgcaa	tgagaacaat	catctaaaaa	120
tggtctgaac	taggtcaggc	gcgggttgctc	atgcotgtaa	tcccaccact	ttgggaggcc	180
gaggcaattg	gatcacctga	ggtcaggatt	ttgagaccag	cttgaccaac	atggtggaat	240
cccatctcta	ctaaaaatac	aaaaaattag	cggggtgtgg	tgccacaccc	ctgtaatccc	300
acctactcag	gaggctgagg	caggaaaatc	ccttgaaccc	aggaggcaaa	ggttgcattg	360
agccgaaata	acaccactgc	actccagcct	ggacgataga	gtgagacccc	atctcaaaaa	420
aagagcagct	gtgacaaatg	cctgtattga	attgcaggtc	agtcttccac	ctccactacc	480
ggtgocaaaa	aaagggtctg	cccaaaaagg	actaaaaggg	atccagcttt	gaattctggg	540
gtctctcaaa	agcctgatcc	tgccaaaacc	aagaatcgcc	gcaaaaaggaa	gccatccact	600
tctgatgatt	ctgactctaa	ttttgagaaa	attgttttga	aagcagtcac	aagcaagggtg	660
agtgttgatc	ctagtccagtc	cttttgctgt	agatgtttctg	aaacacgtaa	ctaagccatt	720
gttctttaaaa	atttggcata	tctttaagaa	aattaactct	catattctgt	tagcttttac	780
tgtacatatt	tagttttaac	aaagttaaat	atgccactta	tttggccaat	ggaagagttg	840
gccttagatc	tgcttcttat	tacttggttag	aaaatagaaa	actccttgaa	tatagtgtct	900
tgatacattt	ttttacatta	caattatggt	gtcagattta	caatgtgcaa	gttacctggg	960
cttttctctt	ttagaaatcc	aagggggaga	gtgatgactt	ccatatggac	tttgactcag	1020
ctgtggctcc	tcgggcaaaa	tctgtacggg	caaagaaacc	tataaagtac	ctggaagagt	1080
cagatgaaga	tgatctgttt	taaaatgtga	ggcgattatt	ttaagtaatt	atcttaccac	1140
gccaagact	ggtttttaag	ttacctgaag	ctcttaactt	cctccctctt	gaatttagtt	1200
tggggaagg	gttttttagta	caagacatca	aagtgaagta	aagcccaagt	gttcttttagc	1260
tttttataat	actgtataaa	tagtgacat	ctcatgggca	ttgttttctt	ctctgctttg	1320
tctgtgtttt	gagtctgctt	cttttgtctt	taaaacctga	tttttaagtt	cttctgaact	1380
gtagaaatag	ctatctgac	acttcagcgt	aaagcagtg	gtttattaac	catccactaa	1440
gctaaaacta	gagcagtttg	atttaaaaagt	gtcactcttc	ctcctttctt	actttcagta	1500
gatattgagat	agagcataat	tatctgtttt	atcttagttt	tatacataat	ttaccatcag	1560
atagaacttt	atggtttctag	tacagatact	ctactacact	cagcctctta	tgtgccaagt	1620
ttttcttttaa	gcaatgagaa	attgctcatg	ttcttcatct	tctcaaatca	tcagaggccg	1680
aagaaaaaca	ctttggctgt	gtctataact	tgacacagtc	aatagaatga	agaaaattag	1740
agtagttatg	tgattatttc	agctcttgac	ctgtcccttc	tggtgcctc	tgagtctgaa	1800
tctoccaaag	agagaaacca	atttctaaga	ggactggatt	gcagaagact	cggggacaac	1860
atttgatcca	agatcttaaa	tggttatattg	ataaccatgc	tcagcaatga	gctattagat	1920
tcattttggg	aaatctccat	aatttcaatt	tgtaaaacttt	gttaagacct	gtctacattg	1980
ttatatgtgt	gtgacttgag	taatgttatc	aacgtttttg	taaatattta	ctatgttttt	2040
ctattagcta	aattccaaca	attttgtact	ttaataaaat	gttctaaaca	ttgaaa	2096

<210> 21

<211> 2160

<212> DNA

<213> Homo sapiens

<400> 21							
	agccccctgc	ccctgcgcgc	ccccgcgcgc	ctgcctgggc	cgggccgagg	atgoggogca	60
	gcgcctcggc	ggccagggtt	gtccccctcc	ggcacgcctg	ctaacttccc	ccgctaogtc	120
5	cccgcttgcc	ggcggggccg	ccccgtctcc	ccggggcctc	cggggtccggg	tcctccagga	180
	cgggccaggcc	gtgcgcgcgt	gtgcctcccg	ccgctcgcgc	gcgcgcgcgc	cgctccccgc	240
	ctgcgcgccag	cgccccgcgc	ccgcgcgccca	gtcctcgggc	ggcccatgct	gccccctctgc	300
	ctcgtggccg	ccctgctgct	ggcgcgcggg	ccggggccga	gcctggggcg	cgaagccatc	360
	cactgccccg	cctgctccga	ggagaagctg	gcgcgcctgc	gcccccccg	gggctgcgag	420
10	gagctgggtc	gagaggcggg	ctgcggctgt	tgccgcaact	gcgcctcggg	cttggggatg	480
	ccctgcgggg	tgtaaccccc	ccgttgccgc	tcgggcctgc	gctgctaccc	gccccgaggg	540
	gtggagaagc	ccctgcacac	actgatgcac	gggcaaggcg	tgatcatgga	gctggcggag	600
	atcgaggcca	tcaggaag	cctgcagccc	totgacaagg	acgagggtga	ccacccccac	660
	aacagcttca	gccccgttag	cgcccatgac	cgcagggtcc	tgagaagca	cttcgccaaa	720
	attcgagacc	ggagcaccag	tgggggcaag	atgaaggcca	atggggcgcc	ccggggagat	780
15	gccccgcctg	tgccccaggg	ctcctgccag	agcgagctgc	accgggcgct	ggagcggctg	840
	gocgcttcac	agagccgcac	ccacgaggac	ctotacttca	tcacatccc	caactgcgac	900
	cgcaacggca	acttccaccc	caagcagtgt	caaccagctc	tgatggggca	gcgtggcaag	960
	tgctggtgtg	tggaccggaa	gacgggggtg	aagcttccgg	ggggcctgga	gccaaagggg	1020
	gagctggaact	gccaccagct	ggctgacagc	tttcgagagt	gaggccctgc	agcaggccag	1080
20	ggactcagcg	tcacctgcta	ctcctgtgct	ctggaggctg	cagagctgac	ccagagtggg	1140
	gtctgagctc	gagtcctgtc	totgcctgcg	gcccagaagt	ttccctcaaa	tgccgctgtg	1200
	caogtgtgcg	tgtgcgtgcg	tgtgtgtgtg	tttgtgagca	tggtgtgtgc	cttggggtaa	1260
	gccagagcct	gggtgtttct	ctttggtgtt	acacagccca	agaggactga	gactggcact	1320
	tagcccaaga	ggtctgagcc	ctggtgtgtt	tcagatcga	tcctggattc	actcactcac	1380
25	tcattccttc	actcatccag	ccacctaaaa	acatttactg	accatgtact	acgtgccagc	1440
	tctagttttc	agccttggga	ggtttttatc	tgacttcctc	tgattttggc	atgtggagac	1500
	actcctataa	ggagagttca	agcctgtggg	agtagaaaaa	tcctattccc	agagtccag	1560
	gagaagagac	atgtaccttg	accatcgtcc	ttcctctcaa	gctagcccag	aggggtgggag	1620
	cctaagggaag	ogtggggtag	cagatggagt	aatgggtcacg	aggtccagac	ccactcccaa	1680
	agctcagaact	tgccaggctc	cctttctctt	cttccccagg	tccttccttt	aggtctggtt	1740
30	gttgaccat	ctgcttggtt	ggctggcagc	tgagagccct	gctgtgggag	agcgaagggg	1800
	gtcaaaggaa	gacttgaagc	acagagggct	agggagggtg	ggtacatttc	totgagcagt	1860
	caggggtggg	agaaagaatg	caagagtggg	ctgaatgtgc	ctaattggga	agaccacagt	1920
	gtaggggat	gaggggcttc	ctgggtcctg	ttccctacc	ccatttgtgg	tcacagccat	1980
	gaagtacccg	ggatgaacct	atccttcacg	tggtctgcctc	cctgtagctc	tgccctccctc	2040
35	tcocatcttc	cttccccctac	acctccctcc	ccacacctcc	ctactccctc	gggcatcttc	2100
	tggcttgact	ggatggaagg	agacttagga	acctaccagt	tgcccatgat	gtcttttctt	2160

<210> 22

<211> 2215

40

<212> DNA

<213> Homo sapiens

45

<400> 22

50

55

ctgcaggag	ccatgattgc	accactgcac	tcagcctgg	gcaacagagt	gagaccatgt	60
ctcaagaaaa	aaaaaaaaaga	aagaaaccac	tgctctaggc	taaatcccag	ccagagttag	120
agccaccag	ctaaactggc	ctgttttccc	ctatttcctt	ccccgaagg	atgcctgtgt	180
caagatgagg	tcacggacga	ttacatcgga	gacaacacca	cagtggacta	cactttgttc	240
gagtctttgt	gctccaagaa	ggacgtgogg	aactttaaag	cctggttcct	ccctatcatg	300
tactccatca	tttgtttcgt	gggcctactg	ggcaatgggc	tggtcgtgtt	gacctatata	360
tatttcaaga	ggctcaagac	catgaccgat	acctacctgc	tcaacctggc	ggtggcagac	420
atcctcttcc	tcctgacctt	tcctttctgg	gcctacagcg	oggccaagtc	ctgggtcttc	480
ggtgtccact	tttgcaagct	catctttgcc	atctacaaga	tgagcttctt	cagtggcatg	540
ctcctacttc	tttgcatcag	cattgaccgc	tacgtggcca	togtccaggc	tgtctcagct	600

	caccgccacc	gtgcccgcgt	ccttctcacc	agcaagctgt	cctgtgtggg	catctggata	660
	ctagccacag	tgctctccat	cccagagctc	ctgtacagt	acctccagag	gagcagcagt	720
5	gagcaagcga	tgogatgctc	tctcatcaca	gagcatgtgg	aggcctttat	caccatccag	780
	gtggcccaga	tggtgatcgg	cttcttggtc	cccctgctgg	ccatgagctt	ctgttacctt	840
	gtcatcatcc	gcacctgct	ccaggcaocg	aactttgagc	gcaacaaggc	catcaagggtg	900
	atcatcgctg	tggtcgtggg	cttcatagtc	ttccagctgc	cctacaatgg	ggtggtcctg	960
	gcccagacgg	tgggccaaactt	caacatcacc	agtagcacct	gtgagctcag	taagcaactc	1020
10	aacatcgctt	acgaogtcac	ctacagcctg	gcctgcgtcc	gctgctgctg	caaccctttc	1080
	ttgtacgcct	tcatogggct	caagtccgc	aacgatctct	tcaagctctt	caaggacctg	1140
	ggctgcctca	gccaggagca	gctccggcag	tggtcttcct	gtcggcacat	ccggcgctcc	1200
	tccatgagt	tgaggccga	gaccaccacc	acctctctcc	cataggcgac	tcttctgcct	1260
	ggactagagg	gacctctccc	agggtccctg	gggtggggat	agggagcaga	tgcaatgact	1320
	caggacatcc	ccccgccaaa	agctgctcag	ggaaaagcag	ctctccctcc	agagtgcagg	1380
15	ccctgcctca	gaagttagct	tcaccccaat	cccagctacc	tcaaccaatg	ccgaaaaaga	1440
	cagggtctgat	aagotaacac	cagacagaca	acactgggaa	acagaggcta	ttgtccccta	1500
	aaccaaaaac	tgaaagtga	agtcacagaa	ctgttcccac	ctgctggagt	gaagggggcca	1560
	aggagggtga	gtgcaagggg	cgtgggagtg	gcctgaagag	tcctctgaat	gaacctcttg	1620
	gcctcccaca	gactcaaatg	ctcagaccag	ctcttccgaa	aaccaggcct	tatctccaag	1680
20	accagagata	gtggggagac	ttcttggtt	ggtgaggaaa	agcggacatc	agctggtcaa	1740
	acaaactctc	tgaacccctc	cctccatcgt	tttcttact	gtcctccaag	ccagcgggaa	1800
	tggcagctgc	cacgcgcgcc	taaaagcaca	ctcatccctc	cacttgccgc	gtcgcctcc	1860
	caggctctca	acaggggaga	gtgtggtgtt	tcctgcaggc	caggccagct	gcctccgcgt	1920
	gatcaaaccc	acactctggg	ctccagagtg	gggatgacat	gcactcagct	cttggctcca	1980
25	ctgggatggg	aggagaggac	aagggaatg	tcaggggcgg	ggaggggtgac	agtggccgcc	2040
	caaggccacg	agcttggtct	ttgtctttt	tcacaggggac	tgaaaacctc	tcctcatgtt	2100
	ctgctttcga	ttcgttaaga	gagcaacatt	ttaccacac	acagataaag	ttttcccttg	2160
	aggaaacaac	agctttaaaa	gaaaaaagaa	aaaaaaagct	tggttaagtca	agtag	2215
	<210>	23					
30	<211>	958					
	<212>	DNA					
35	<213>	Homo sapiens					
	<400>	23					
40	ggggccggac	gcgagggggc	gggagagcgc	gggacaaagg	gaagcgaagc	cggagctgcg	60
	ggcgcttttt	ctgcccggcg	tgtctcagat	tcattcttaa	ggaactgaga	acttaatctt	120
	ccaaaatgtc	aaaaagacca	tcttatgccc	cacctccac	cccagctcct	gcaacacaaa	180
	tgcccagcac	accagggttt	gtgggataca	atccatacag	tcatctcgcc	tacaacaact	240
	acaggctggg	agggaacccg	agcaccaaca	gccgggtcac	ggcatacctc	ggtatcacga	300
	ttccaaaacc	cccaaagcca	ccagataagc	cgctgatgcc	ctacatgagg	tacagcagaa	360
45	aggtctggga	ccaagtaaag	gcttccaacc	ctgacctaaa	gttgtgggag	attggcaaga	420
	ttattggtgg	catgtggcga	gatctcactg	atgaagaaaa	acaagaatat	ttaaacgaat	480
	acgaagcaga	aaagatagag	tacaatgaat	ctatgaaggc	ctatcataat	ccccccgcgt	540
	accttgctta	cataaatgca	aaaagtcgtg	cagaagctgc	tttagaggaa	gaaagtcgac	600
	agagacaatc	tcgcatggag	aaaggagaac	ogtacatgag	cattcagcct	gctgaagatc	660
50	cagatgatta	tgatgatggc	ttttcaatga	agcatacagc	caccgcccgt	ttccagagaa	720
	accaccgcct	catcagtga	attcttagtg	agagtgtggg	gccagaogtt	cggtcagttg	780
	tcacaacagc	tagaatgcag	gtcctcaaac	ggcaggtcca	gtccttaatg	gttcatcagc	840
	gaaaactaga	agctgaactt	cttcaaatag	aggaacgcga	ccaggagaag	aagaggaaat	900
	tcctggaaag	cacagattca	tttaacaatg	aacttaaaag	gttgtgcggg	ctgaaagt	958
55							

<210> 24

<211> 6483

<212> DNA

<213> Homo sapiens

<400> 24

aagcttctaa	ttgcagttca	accacctgtt	acatatcttc	aggaaaaaat	cacaacctct	60
caacttcaac	ttcctcttct	ataaattaga	aataacaata	accacacctg	taaccccagc	120
actttgggag	gccaaggcag	gcagatcaag	aggtagggag	attgagacca	tcctggctaa	180
catgatgaaa	ccctgtctct	accaaaaaga	caaaaaatta	gccaggtagt	gtggcacaca	240
cctgtagtcc	cagctactcg	ggaggctgag	gcaggagaat	ggcgtgaacc	cgggagggtg	300
agcttgccagt	gagccgagat	ggcgcactg	cactccagcc	tgggcgacag	agcaagcctc	360
cgtctaaaaa	aaaaaaaaga	aagaaagaaa	gaaagaaaga	aaagaaataa	taataaccac	420
cattcctatc	tcaacagott	gttctagaaa	tttttaaagc	acagtatcac	aaacagcact	480
acataattgt	aaaacatgta	tgaatatata	catccaaaca	acagcaatgt	catagcctat	540
gggtagatat	aatccttatac	aatgtaccaa	aatcccaatt	tacttcaacta	gacaaaactgt	600
tataccaaat	tctgtacaca	gtatatccaa	gaaaatgtgt	tgtttttatt	gagaaactga	660
acctagcctg	ggaacacatg	tgcacagtct	agttcataat	atttgggtgca	agtatcatte	720
tctaatatag	atttacattt	ttgcaagcaa	atttttactt	gcaatcgtaa	catatccaaa	780
ttttcccttt	ttactcaatc	agaacttagt	gtaaaagtact	acaagtagt	tccttcggatt	840
tcattgctaag	aaaataatgc	agattttctg	cattattatg	gtcttcacag	aaaccttaac	900
tatgatgaat	ttaaaagtgc	aaaataatoc	aggataactt	tatgatttca	catttttttaa	960
tggtaaaaat	aatgcatcca	ttaattagaa	aattctaaaa	tcattacttc	cactttctta	1020
ggcaaaatat	caataatactc	tcatttgcca	aataaattaa	aagatctcct	acaaacacaa	1080
tctcctaaat	tgtggtttta	tggctttaat	gttttatgtg	tggcaactat	tgatgctagt	1140
taaaatttta	gaaactcttt	ctttttgatt	ccctacagtt	gtctacaaga	accttattgt	1200
agcatgatcc	tgcagacttt	tatactatct	gttgctccaa	ttaaaactgt	ttaaaacatg	1260
aatttgaaaa	atcttatttt	aactataatt	ttgtagotga	aacttttttt	tctaaaacttt	1320
gcaaacattc	tatgcaacct	gaattagtgc	tgagaaaatt	ggatcttaat	ggttgctcaa	1380
tggtcttcaa	caggtgaaaa	gcataataaa	acatgctcat	ctgaactcca	cccatcttca	1440
atttcaaat	agcatacctc	gtgtttatct	ttaggggcaa	ttcaaaattg	tacatattag	1500
gattgggttat	tactgaagat	aatttatgca	atcataagcc	aaagatgcta	agttggcaaa	1560
aagaaaacaa	tgtaagtaag	caaaccttaa	cacatgtgga	cacacctctc	cagtataata	1620
aggcttgcca	ctgtccttgg	tagcaggcac	tcctcgggct	aaacagcatc	accatgtctg	1680
ttcgatacag	ctcaagcaag	cactactctt	ccctccgcag	tggaggagga	ggaggaggag	1740
gggatgtggg	aggagggagg	ggagtgtcat	ccctaagaat	ttctagcagc	aaaggctccc	1800
ttggtggagg	atttagctca	gggggggtca	gtggtggctc	ttttagccgt	gggagctctg	1860
gtgggggatg	ctttgggggc	tcacagggtg	gctatggagg	attaggagggt	tttgggtggag	1920
gtagctttca	tggagactat	ggaagtagca	gctttggtgg	gagttatgga	ggcagctttg	1980
gagggggcaa	tttcggagggt	ggcagctttg	gtgggggcag	ctttggtgga	ggcggctttg	2040
gtggaggcgg	ctttggagga	ggctttgggtg	gtggatttgg	aggagatggg	ggccttctct	2100
ctggaaatga	aaaagtaacc	atgcagaatc	tgaatgaccg	cctggcttcc	tacttggaca	2160
aagtctgggc	tctggaagaa	tcaaaactatg	agctggaagg	caaaatcaag	gagtggatatg	2220
aaaagcatgg	caactcacat	caggggggagc	ctcgtgaacta	cagcaaatac	tacaaaacca	2280
tcgatgaoot	taaaaatcag	gtaagaggta	tttttaaato	cagctttaag	tatcttgtcc	2340
atgtaatoca	gacagatgaa	tcttaaatga	agcacaatgt	ggctgttcac	tatgcttacc	2400
catgttaactt	tcttccttca	aaaataacccc	agtctcatca	aagataaaca	tctgtgaaac	2460
tatggtcatg	gcaatcttca	tccagcaagt	gtgctacttg	tcttaagagg	atgggagatt	2520
tactaagcac	ttttgagggt	ttaattgagca	tacaatgagt	ccacagttaa	aatatgctag	2580
gctattttaca	aatgtagaaa	ctgaaaaaaa	aatcatgat	atgaatcaga	acaaaatgtt	2640
attcagactg	ataacaagcc	atattcagta	ccaacatggc	aagaaaaata	aattttccag	2700
tatgaaaatg	ggacactgct	tgcttctaag	gaatttctga	attgtacctc	ttgtgtacca	2760
gttcagagtg	tatttattta	ttagtattta	tcatgagtta	aacaaatgca	ggtgtgagtc	2820
agccaaagca	tggctgaaat	acatggaaat	cacatagtct	aaaagaggag	ggcacactta	2880

	caggaataca	tctatataat	tccagttagt	tttcagaaag	gaataattcg	tgtacagaaa	2940
	tacaagactg	gagaaattcc	aagagaacaa	ataattcaaa	gttaagtata	tgggtaagcc	3000
	tgcaatattt	catatttaaa	ataaaaaatt	ttcccaagat	tttgttaagag	aacaacataa	3060
5	aagtgcagag	tgcatctatg	tcactacaaa	agccatatct	gcatctgacc	tcttctcaaa	3120
	taactgtgcc	tctccctcca	gattctcaac	ctaacaactg	ataatgccaa	catcctgctt	3180
	cagatcgaca	atgccaggct	ggcagctgat	gacttcaggc	tgaagttaagt	taagtgatcg	3240
	ttgtataata	ctatcacaa	gaatacatca	gtgggtttta	acaatgactt	gggatgcctt	3300
	caataacatt	tacatttttc	tgaattcaac	caaagtttaa	tagtattgga	gttatctgag	3360
10	aaattttoca	tgtcagtggt	accttttttg	caatatttaa	ggaagaaaat	gcatatttaa	3420
	gtaactgcta	aggttttttc	cattaaacca	ctattacttc	taagagaact	gtacatgaca	3480
	aatattgcca	ttacatgaga	tcaactatgt	agttgctttt	taaatagtct	ctgcccagat	3540
	acatctcccc	tatataagtt	ataaccagta	ttgatcatcat	gcttgtttca	ggtagtgaaa	3600
	tgagtgtagct	ctgcgccaga	gcgtggaggc	tgacatcaac	ggcctgcgta	gggtgctgga	3660
15	tgagctgacc	ctgaccaagg	ctgacctgga	gatgcaaatt	gagagcctga	ctgaagagct	3720
	ggcctatctg	aagaagaacc	acgaggaggc	gacacaaaag	ttatactttt	cccagccaaa	3780
	agagagttca	ttatggtcct	cgtgtagcca	ataaatcttt	ctgttccctca	aacaggaaat	3840
	gaaagacott	cgaaatgtgt	ccactgggtga	tgtgaatgtg	gaaatgaatg	ctgccccggg	3900
	tgttgatctg	actcaacttc	tgaataacat	gagaagccaa	tatgaacaac	ttgctgaaca	3960
	aaaccgcaaa	gatgctgaag	cctggttcaa	tgaaaaggta	aagtaatott	ccttatagt	4020
20	aaactcatgg	aggttttatc	atttcagaat	ttcctcacc	ttttccttgt	ttttaatact	4080
	ctagagcaag	gaactgacta	cagaaattga	taataacatt	gaacagatat	ccagctataa	4140
	atctgagatt	actgaattga	gacgtaatgt	acaagctctg	gagatagaac	tacagtccca	4200
	actggccttg	gtatgttaac	tctcatgaaa	tgacttcaac	tttatcatac	aaagtttcat	4260
	gctcacctaa	gaatatgcaa	tgcaacaaaa	aatgacagag	ttggaggtaa	gaaagagaaa	4320
	acaaagtga	gctcatgtta	atggaggaaa	agtactacta	gtgttgatct	aaaagtgtct	4380
25	aaactgaaat	gggtgcatta	aacatacaac	aaattctgtt	cattttctta	ttctctata	4440
	taatgcctta	ctaaataatc	aaataagcgt	caccatactc	aactgaacaa	ggaagtcaact	4500
	aagccacaaa	aaaatccgtt	tcagaaaacaa	tcctctggaag	cctccttggc	agaaacagaa	4560
	ggctgcctact	gtgtgcagct	ctcacagatt	cacgcccaga	tatccgctct	ggaagaacag	4620
	ttgcaacaga	ttcgagctga	aaccgagctg	cagaatactg	aataccaaca	actcctggat	4680
30	attaagatcc	gactggagaa	tgaaattcaa	acctaccgca	gcctgctaga	aggagaggga	4740
	aggtaaatca	taacatgaaa	agttatocca	gtttctttta	ttcaatatcc	cagatagcaa	4800
	ggcttatcta	aaccccaaga	agatgccaga	gaatgagagg	aagggaggag	agagggtaga	4860
	gtacagaaaa	aggagtacgc	aaccgcaatc	tcactttctc	atgaatttgg	cccaaaatga	4920
	ttcttaagag	ttctgtgaac	ttaacattgt	tttcaaagga	tgggttttaa	aatatatacc	4980
	tggcagggtt	ttattttttc	aacacgtttt	gottattttc	taaattaacg	gcaactggaa	5040
35	agctaccac	cgtttttcaa	cgttagagat	aaccgaatgt	gacctcacc	cgttttagtt	5100
	cggaggcggc	ggacgcggcg	gcggaagttt	ogggcgcgcg	tacggcgcg	gaagctccgg	5160
	cggcggaagc	tccggcgcg	gctacggcg	cgccacggcg	ggcagttccg	gcggcggtca	5220
	cggaggcgga	agctccggcg	gcggaagctc	cggcgcgcg	tacggggcg	gaagctccag	5280
	cggcgccac	gcggcgggaa	gctccagcgg	cggccacggc	ggcagttcca	gcggcggtca	5340
40	cgggtgtggc	agttccggcg	gcggcgcgcg	cggctacggg	ggcggcagct	ccggcgcgcg	5400
	cagcagctcc	ggcgcgggat	acggcgcgcg	cagctccagc	ggaggccaca	agtcctcctc	5460
	ttccgggtcc	gtggcgagat	cttcacttaa	gggaccaagg	tcagcagaaa	ctagctgggg	5520
	taactctaga	ttagttttta	cttctgtga	tgggtttttt	gcgctttaag	ctctagagtt	5580
	gttttaaaaa	attaaaaatc	ttagagacgg	ttccgtttgc	atttggtcac	aaactactct	5640
	taacaccagc	cgtgaaaaat	ggcatgatca	aaatgtcata	ccttaagcat	ttttttgggc	5700
45	ttaacaatgt	aaagttgaaa	tttctctctt	tttacaatat	ttgcttggtta	attactaagg	5760
	atccctacag	actgttttaa	attttttttc	catcattcac	acagatacta	acaaaaccag	5820
	agtaatcaag	acaattattg	aagaggtggc	gcccagcgg	agagttcttt	catctatggt	5880
	tgaatcagaa	accaagaac	actactatta	aactgcatca	agaggaaaag	gtctcccttc	5940
	acacagacca	ttatttacag	atgcatggaa	aacaagctct	ccaagaaaac	actctgtct	6000
50	tgatggtcta	tggaaaataga	ccttgaaaat	aaggtgtcta	caaggtgttt	tgtggtttct	6060
	gtattttctc	ttttactttt	accacaaagt	gttctttaat	ggaaagaaaa	acaactttgt	6120
	gttctcattt	actaatgaat	ttcaataaac	tttcttactg	atgcaaaacta	tcccaatttg	6180
	tcagaattta	totttactta	agtacataat	actcttttaa	attaaagatt	agtaaccat	6240
	agcagttgaa	ggttgatgta	tcagaaaatt	cggagacag	aactattgtc	atgccttttc	6300
	taagtttttt	aatcatgtat	gttcagacca	cogtcagtaa	attcactgag	taaagtctgt	6360
55	aaatccccaa	tattactctt	taagatacac	aatatgtgga	aggctcccag	ctctctggct	6420

ttaaattatt tcaatcctgg aaattctgga atatctcaaa tataaccccc aaaataataa 6480
taa 6483

<210> 25

<211> 1871

<212> DNA

<213> Homo sapiens

<400> 25

agttgtggcc	accttcccca	ggccatggat	ctctccaaca	acaccatgtc	actctcagtg	60
cgcacccccg	gactgtcccc	gcggctctcc	tcgcagagtg	tgataggcag	acccaggggc	120
atgtctgctt	ccagtgttgg	aagtggttat	gggggaagtg	cctttggctt	tggagccagc	180
tgtgggggag	gcttttctgc	tgottccatg	tttggttota	gttccggctt	tgggggtggc	240
tcoggaagtt	ccatggcagg	aggactgggt	gctggttatg	ggagagccct	gggtggaggt	300
agctttggag	ggctggggat	gggatttggg	ggcagcccag	gaggtggctc	tctaggtatt	360
ctctcgggca	atgatggagg	ccttctttct	ggatcagaaa	aagaaactat	gcaaaatctt	420
aatgatagat	tagcttcccta	cctggataag	gtgcgagctc	tagaagaggc	taatactgag	480
ctagaaaata	aaattcgaga	atggtatgaa	acacgaggaa	ctgggactgc	agatgottca	540
cagagcgatt	acagcaaata	ttatccactg	attgaagacc	tcaggaataa	gatcatttca	600
gccagcattg	gaaatgccca	gctcctcttg	cagattgaca	atgcgagact	agctgctgag	660
gacttcagga	tgaagtatga	gaatgaactg	gccttgogcc	agggcgtaga	ggccgacatc	720
aatggcctgc	gccgggtgct	ggaogagctg	accctgacca	ggaccgacct	ggagatgcag	780
atcgagagcc	tgaacgagga	gctggcctac	atgaagaaga	accacgagga	tgagctccaa	840
agcttccggg	tgggcgggccc	aggcgaggctc	agcgtagaaa	tggacgctgc	ccccggagtg	900
gacctcacca	ggctcctcaa	tgatatgcgg	gcgcagtatg	aaaccatcgc	tgagcagaat	960
cggaaggacg	ctgaagcctg	gttcattgaa	aagagcgggg	agctccgtaa	ggagattagc	1020
accaacacog	agcagcttca	gtccagcaag	agcgagggtca	ccgacctgcg	tcgcgccttt	1080
cagaacctgg	agatcgagct	acagtcccag	ctcgccatga	agaaatccct	ggaggactcc	1140
ttggcogaag	cagaggcgga	ttactgcgcg	cagctgtccc	aggtgcagca	gctcatcagc	1200
aacctggagg	cacagctgct	ccagggtgcgc	gcggacgcag	agcgccagaa	cgtggaccac	1260
cagcggctgc	tgaatgtcaa	ggcccgccctg	gagctggaga	ttgagacctc	ccgccgcctg	1320
ctggacgggg	aggcccaagg	tgatggtttg	gaggaaagt	tatttgtgac	agactccaaa	1380
tcacaagcac	agtcaactga	ttcctctaaa	gacccaacca	aaacccgaaa	aatcaagaca	1440
gttgtgcagg	agatgggtgaa	tggtgaggtg	gtctcatctc	aagttcagga	aattgaagaa	1500
ctaattgtaaa	atttcacaag	atctgcccc	tgattggttc	cttaggaaca	agaaatttac	1560
aagtagaaat	tattcctttc	agagtaacat	gctgtattac	ttcaatccct	atttttgtct	1620
gttcattttt	ccttggtattc	cctattcaca	ttgaatcctt	tttgcccttc	tgaaacaata	1680
ttcagtcaca	agtcattttg	gtcatgttgg	tctttgtaac	aaatcaaaat	taccttatat	1740
ccttctggac	aactggagta	gtcttttaac	gaactttctt	ctggttaacc	ggaatatttt	1800
cttaatacata	gagctttact	caagtagtat	tgttttaata	gagttaattg	taataaaaga	1860
tgaaatggtaa	a					1871

<210> 26

<211> 1447

<212> DNA

<213> Homo sapiens

<400> 26
 ctgcaactgg ttctgcgagg gctccttcaa tggcagcgag aaggagacta tgcagttcct 60
 gaaogaccgc ctggccagct acctggagaa ggtgcgtcac gtggagcggg acaacgcgga 120
 5 gctggagaac ctcatccggg agcgggtctca gcagcaggag cccttgctgt gccccagcta 180
 ccagtcctac ttcaagaoca ttgaggagct ccagcagaag atcctgtgca gcaagtctga 240
 gaatgccagg ctggtggtgc agatcgacaa tgccaagctg gctgcagatg acttcagaac 300
 caagtaccag acggagcagt ccctgcggca gctggtggag tccgacatca acagcctgcg 360
 caggattctg gatgagctga ccctgtgcag gtctgacctg gaggcccaga tggagtccct 420
 10 gaaggaggag ctgctgtccc tcaagcagaa ccatgagcag gaagtcaaca ccttgcgctg 480
 ccagcttgga gaccgcctca acgtggaggt ggacgctgct ccgctgtgg acctgaacca 540
 ggtcctgaac gagaccagga atcagtatga ggccctggtg gaaaccaacc gcagggaagt 600
 ggagcaatgg ttccgccagc agaccgagga gctgaacaag cagggtggtat ccagctcgga 660
 gcagctgcag tcctaccagg cggagatcat cgagctgaga cgcacagtca atgccctgga 720
 15 gatcgagctg caggcccagc acaacctgcg atactctctg gaaaacacgc tgacagagag 780
 cgaggcccgcc tacagctccc agctgtccca ggtgcagagc ctgatcacca acgtggagtc 840
 ccagctggcg gagatccgca gtgacctgga gcggcagAAC caggagtatc aggtgctgct 900
 ggaogtgcg gcgcggtgg agtgtgagat caacacatac cggagcctgc tggagagcga 960
 ggactgcaag ctgcccctca acccctgcgc caccaccaat gcatgtgaaa agcccattgg 1020
 atcctgtgtc accaatcctt gtggtcctcg ttcccgctgt gggccttgca acaccttgg 1080
 20 gtaactagata ccctggggcc agcagaagta tagcatgaag acagaactac catcggtggg 1140
 ccagttctgc ctctctgaca accatcagcc accggacccc accccgaggc atcaccacaa 1200
 atcatggtct ggaaggagaa caaatgocca gcgtttgggt ctgactctga gcctagggct 1260
 actgatcctc ctcaccccag gtccctctoc tgtagtcatg ctgagttctg atggtcagag 1320
 gttggagctg tgacagtggc atacgaggtg ttttgttctc totgctgctt ctacctttat 1380
 25 tgcagttccc caaatgcct aataaacttt cctcttgcaa agcagacaaa aaaaaaaaaa 1440
 aaaaaaa 1447

<210> 27

<211> 261

<212> PRT

<213> Homo sapiens

<400> 27
 Met Asn Pro Asn Cys Ala Arg Cys Gly Lys Ile Val Tyr Pro Thr Glu
 1 5 10 15
 40 Lys Val Asn Cys Leu Asp Lys Phe Trp His Lys Ala Cys Phe His Cys
 20 25 30
 Glu Thr Cys Lys Met Thr Leu Asn Met Lys Asn Tyr Lys Gly Tyr Glu
 35 40 45
 Lys Lys Pro Tyr Cys Asn Ala His Tyr Pro Lys Gln Ser Phe Thr Met
 50 55 60
 45 Val Ala Asp Thr Pro Glu Asn Leu Arg Leu Lys Gln Gln Ser Glu Leu
 65 70 75 80
 Gln Ser Gln Val Arg Tyr Lys Glu Glu Phe Glu Lys Asn Lys Gly Lys
 85 90 95
 Gly Phe Ser Val Val Ala Asp Thr Pro Glu Leu Gln Arg Ile Lys Lys
 100 105 110
 50 Thr Gln Asp Gln Ile Ser Asn Ile Lys Tyr His Glu Glu Phe Glu Lys
 115 120 125
 Ser Arg Met Gly Pro Ser Gly Gly Glu Gly Met Glu Pro Glu Arg Arg
 130 135 140
 55 Asp Ser Gln Asp Gly Ser Ser Tyr Arg Arg Pro Leu Glu Gln Gln Gln
 145 150 155 160

Pro His His Ile Pro Thr Ser Ala Pro Val Tyr Gln Gln Pro Gln Gln
 165 170 175
 Gln Pro Val Ala Gln Ser Tyr Gly Gly Tyr Lys Glu Pro Ala Ala Pro
 180 185 190
 Val Ser Ile Gln Arg Ser Ala Pro Gly Gly Gly Lys Arg Tyr Arg
 195 200 205
 Ala Val Tyr Asp Tyr Ser Ala Ala Asp Glu Asp Glu Val Ser Phe Gln
 210 215 220
 Asp Gly Asp Thr Ile Val Asn Val Gln Gln Ile Asp Asp Gly Trp Met
 225 230 235 240
 Tyr Gly Thr Val Glu Arg Thr Gly Asp Thr Gly Met Leu Pro Ala Asn
 245 250 255
 Tyr Val Glu Ala Ile
 260
 <210> 28
 <211> 478
 <212> PRT
 <213> Homo sapiens
 <400> 28
 Met Val Gln Lys Thr Ser Met Ser Arg Gly Pro Tyr Pro Pro Ser Gln
 1 5 10 15
 Glu Ile Pro Met Glu Val Phe Asp Pro Ser Pro Gln Gly Lys Tyr Ser
 20 25 30
 Lys Arg Lys Gly Arg Phe Lys Arg Ser Asp Gly Ser Thr Ser Ser Asp
 35 40 45
 Thr Thr Ser Asn Ser Phe Val Arg Gln Gly Ser Ala Glu Ser Tyr Thr
 50 55 60
 Ser Arg Pro Ser Asp Ser Asp Val Ser Leu Glu Glu Asp Arg Glu Ala
 65 70 75 80
 Leu Arg Lys Glu Ala Glu Arg Gln Ala Leu Ala Gln Leu Glu Lys Ala
 85 90 95
 Lys Thr Lys Pro Val Ala Phe Ala Val Arg Thr Asn Val Gly Tyr Asn
 100 105 110
 Pro Ser Pro Gly Asp Glu Val Pro Val Gln Gly Val Ala Ile Thr Phe
 115 120 125
 Glu Pro Lys Asp Phe Leu His Ile Lys Glu Lys Tyr Asn Asn Asp Trp
 130 135 140
 Trp Ile Gly Arg Leu Val Lys Glu Gly Cys Glu Val Gly Phe Ile Pro
 145 150 155 160
 Ser Pro Val Lys Leu Asp Ser Leu Arg Leu Leu Gln Glu Gln Lys Leu
 165 170 175
 Arg Gln Asn Arg Leu Gly Ser Ser Lys Ser Gly Asp Asn Ser Ser Ser
 180 185 190
 Ser Leu Gly Asp Val Val Thr Gly Thr Arg Arg Pro Thr Pro Pro Ala
 195 200 205
 Ser Ala Lys Gln Lys Gln Lys Ser Thr Glu His Val Pro Pro Tyr Asp
 210 215 220
 Val Val Pro Ser Met Arg Pro Ile Ile Leu Val Gly Pro Ser Leu Lys
 225 230 235 240
 Gly Tyr Glu Val Thr Asp Met Met Gln Lys Ala Leu Phe Asp Phe Leu
 245 250 255

Lys His Arg Phe Asp Gly Arg Ile Ser Ile Thr Arg Val Thr Ala Asp
 260 265 270
 Ile Ser Leu Ala Lys Arg Ser Val Leu Asn Asn Pro Ser Lys His Ile
 275 280 285
 Ile Ile Glu Arg Ser Asn Thr Arg Ser Ser Leu Ala Glu Val Gln Ser
 290 295 300
 Glu Ile Glu Arg Ile Phe Glu Leu Ala Arg Thr Leu Gln Leu Val Ala
 305 310 315 320
 Leu Asp Ala Asp Thr Ile Asn His Pro Ala Gln Leu Ser Lys Thr Ser
 325 330 335
 Leu Ala Pro Ile Ile Val Tyr Ile Lys Ile Thr Ser Pro Lys Val Leu
 340 345 350
 Gln Arg Leu Ile Lys Ser Arg Gly Lys Ser Gln Ser Lys His Leu Asn
 355 360 365
 Val Gln Ile Ala Ala Ser Glu Lys Leu Ala Gln Cys Pro Pro Glu Met
 370 375 380
 Phe Asp Ile Ile Leu Asp Glu Asn Gln Leu Glu Asp Ala Cys Glu His
 385 390 395 400
 Leu Ala Glu Tyr Leu Glu Ala Tyr Trp Lys Ala Thr His Pro Pro Ser
 405 410 415
 Ser Thr Pro Pro Asn Pro Leu Leu Asn Arg Thr Met Ala Thr Ala Ala
 420 425 430
 Leu Arg Arg Ser Pro Ala Pro Val Ser Asn Leu Gln Val Gln Val Leu
 435 440 445
 Thr Ser Leu Arg Arg Asn Leu Gly Phe Trp Gly Gly Leu Glu Ser Ser
 450 455 460
 Gln Arg Gly Ser Val Val Pro Gln Glu Gln Glu His Ala Met
 465 470 475

<210> 29

<211> 196

<212> PRT

<213> Homo sapiens

<400> 29

Met Ser Met Leu Arg Leu Gln Lys Arg Leu Ala Ser Ser Val Leu Arg
 1 5 10 15
 Cys Gly Lys Lys Lys Val Trp Leu Asp Pro Asn Glu Thr Asn Glu Ile
 20 25 30
 Ala Asn Ala Asn Ser Arg Gln Gln Ile Arg Lys Leu Ile Lys Asp Gly
 35 40 45
 Leu Ile Ile Arg Lys Pro Val Thr Val His Ser Arg Ala Arg Cys Arg
 50 55 60
 Lys Asn Thr Leu Ala Arg Arg Lys Gly Arg His Met Gly Ile Gly Lys
 65 70 75 80
 Arg Lys Gly Thr Ala Asn Ala Arg Met Pro Glu Lys Val Thr Trp Met
 85 90 95
 Arg Arg Met Arg Ile Leu Arg Arg Leu Leu Arg Arg Tyr Arg Glu Ser
 100 105 110
 Lys Lys Ile Asp Arg His Met Tyr His Ser Leu Tyr Leu Lys Val Lys
 115 120 125
 Gly Asn Val Phe Lys Asn Lys Arg Ile Leu Met Glu His Ile His Lys
 130 135 140

Leu Lys Ala Asp Lys Ala Arg Lys Lys Leu Leu Ala Asp Gln Ala Glu
 145 150 155 160
 Ala Arg Arg Ser Lys Thr Lys Glu Ala Arg Lys Arg Arg Glu Glu Arg
 165 170 175
 Leu Gln Ala Lys Lys Glu Glu Ile Ile Lys Thr Leu Ser Lys Glu Glu
 180 185 190
 Glu Thr Lys Lys
 195
 <210> 30
 <211> 1566
 <212> PRT
 <213> Homo sapiens
 <400> 30
 Met Ser Ser Leu Leu Glu Arg Leu His Ala Lys Phe Asn Gln Asn Arg
 1 5 10 15
 Pro Trp Ser Glu Thr Ile Lys Leu Val Arg Gln Val Met Glu Lys Arg
 20 25 30
 Val Val Met Ser Ser Gly Gly His Gln His Leu Val Ser Cys Leu Glu
 35 40 45
 Thr Leu Gln Lys Ala Leu Lys Val Thr Ser Leu Pro Ala Met Thr Asp
 50 55 60
 Arg Leu Glu Ser Ile Ala Gly Gln Asn Gly Leu Gly Ser His Leu Ser
 65 70 75 80
 Ala Ser Gly Thr Glu Cys Tyr Ile Thr Ser Asp Met Phe Tyr Val Glu
 85 90 95
 Val Gln Leu Asp Pro Ala Gly Gln Leu Cys Asp Val Lys Val Ala His
 100 105 110
 His Gly Glu Asn Pro Val Ser Cys Pro Glu Leu Val Gln Gln Leu Arg
 115 120 125
 Glu Lys Asn Ser Asp Glu Phe Ser Lys His Leu Lys Gly Leu Val Asn
 130 135 140
 Leu Tyr Asn Leu Pro Gly Asp Asn Lys Leu Lys Thr Lys Met Tyr Leu
 145 150 155 160
 Ala Leu Gln Ser Leu Glu Gln Asp Leu Ser Lys Met Ala Ile Met Tyr
 165 170 175
 Trp Lys Ala Thr Asn Ala Gly Pro Leu Asp Lys Ile Leu His Gly Ser
 180 185 190
 Val Gly Tyr Leu Thr Pro Arg Ser Gly Gly His Leu Met Asn Leu Lys
 195 200 205
 Tyr Tyr Val Ser Pro Ser Asp Leu Leu Asp Asp Lys Thr Ala Ser Pro
 210 215 220
 Ile Ile Leu His Glu Asn Asn Val Ser Arg Ser Leu Gly Met Asn Ala
 225 230 235 240
 Ser Val Thr Ile Glu Gly Thr Ser Ala Val Tyr Lys Leu Pro Ile Ala
 245 250 255
 Pro Leu Ile Met Gly Ser His Pro Val Asp Asn Lys Trp Thr Pro Ser
 260 265 270
 Phe Ser Ser Ile Thr Ser Ala Asn Ser Val Asp Leu Pro Ala Cys Phe
 275 280 285
 Phe Leu Lys Phe Pro Gln Pro Ile Pro Val Ser Arg Ala Phe Val Gln
 290 295 300

Lys Leu Gln Asn Cys Thr Gly Ile Pro Leu Phe Glu Thr Gln Pro Thr
 305 310 315 320
 Tyr Ala Pro Leu Tyr Glu Leu Ile Thr Gln Phe Glu Leu Ser Lys Asp
 325 330 335
 Pro Asp Pro Ile Pro Leu Asn His Asn Met Arg Phe Tyr Ala Ala Leu
 340 345 350
 Pro Gly Gln Gln His Cys Tyr Phe Leu Asn Lys Asp Ala Pro Leu Pro
 355 360 365
 Asp Gly Arg Ser Leu Gln Gly Thr Leu Val Ser Lys Ile Thr Phe Gln
 370 375 380
 His Pro Gly Arg Val Pro Leu Ile Leu Asn Leu Ile Arg His Gln Val
 385 390 395 400
 Ala Tyr Asn Thr Leu Ile Gly Ser Cys Val Lys Arg Thr Ile Leu Lys
 405 410 415
 Glu Asp Ser Pro Gly Leu Leu Gln Phe Glu Val Cys Pro Leu Ser Glu
 420 425 430
 Ser Arg Phe Ser Val Ser Phe Gln His Pro Val Asn Asp Ser Leu Val
 435 440 445
 Cys Val Val Met Asp Val Gln Gly Leu Thr His Val Ser Cys Lys Leu
 450 455 460
 Tyr Lys Gly Leu Ser Asp Ala Leu Ile Cys Thr Asp Asp Phe Ile Ala
 465 470 475 480
 Lys Val Val Gln Arg Cys Met Ser Ile Pro Val Thr Met Arg Ala Ile
 485 490 495
 Arg Arg Lys Ala Glu Thr Ile Gln Ala Asp Thr Pro Ala Leu Ser Leu
 500 505 510
 Ile Ala Glu Thr Val Glu Asp Met Val Lys Lys Asn Leu Pro Pro Ala
 515 520 525
 Ser Ser Pro Gly Tyr Gly Met Thr Thr Gly Asn Asn Pro Met Ser Gly
 530 535 540
 Thr Thr Thr Ser Thr Asn Thr Phe Pro Gly Gly Pro Ile Ala Thr Leu
 545 550 555 560
 Phe Asn Met Ser Met Ser Ile Lys Asp Arg His Glu Ser Val Gly His
 565 570 575
 Gly Glu Asp Phe Ser Lys Val Ser Gln Asn Pro Ile Leu Thr Ser Leu
 580 585 590
 Leu Gln Ile Thr Gly Asn Gly Gly Ser Thr Ile Gly Ser Ser Pro Thr
 595 600 605
 Pro Pro His His Thr Pro Pro Pro Val Ser Ser Met Ala Gly Asn Thr
 610 615 620
 Lys Asn His Pro Met Leu Met Asn Leu Leu Lys Asp Asn Pro Ala Gln
 625 630 635 640
 Asp Phe Ser Thr Leu Tyr Gly Ser Ser Pro Leu Glu Arg Gln Asn Ser
 645 650 655
 Ser Ser Gly Ser Pro Arg Met Glu Ile Cys Ser Gly Ser Asn Lys Thr
 660 665 670
 Lys Lys Lys Lys Ser Ser Arg Leu Pro Pro Glu Lys Pro Lys His Gln
 675 680 685
 Thr Glu Asp Asp Phe Gln Arg Glu Leu Phe Ser Met Asp Val Asp Ser
 690 695 700
 Gln Asn Pro Ile Phe Asp Val Asn Met Thr Ala Asp Thr Leu Asp Thr
 705 710 715 720
 Pro His Ile Thr Pro Ala Pro Ser Gln Cys Ser Thr Pro Pro Thr Thr
 725 730 735
 Tyr Pro Gln Pro Val Pro His Pro Gln Pro Ser Ile Gln Arg Met Val
 740 745 750
 Arg Leu Ser Ser Ser Asp Ser Ile Gly Pro Asp Val Thr Asp Ile Leu
 755 760 765

Ser Asp Ile Ala Glu Glu Ala Ser Lys Leu Pro Ser Thr Ser Asp Asp
 770 775 780
 Cys Pro Ala Ile Gly Thr Pro Leu Arg Asp Ser Ser Ser Gly His
 785 790 795 800
 Ser Gln Ser Thr Leu Phe Asp Ser Asp Val Phe Gln Thr Asn Asn Asn
 805 810 815
 Glu Asn Pro Tyr Thr Asp Pro Ala Asp Leu Ile Ala Asp Ala Ala Gly
 820 825 830
 Ser Pro Ser Ser Asp Ser Pro Thr Asn His Phe Phe His Asp Gly Val
 835 840 845
 Asp Phe Asn Pro Asp Leu Leu Asn Ser Gln Ser Gln Ser Gly Phe Gly
 850 855 860
 Glu Glu Tyr Phe Asp Glu Ser Ser Gln Ser Gly Asp Asn Asp Asp Phe
 865 870 875 880
 Lys Gly Phe Ala Ser Gln Ala Leu Asn Thr Leu Gly Val Pro Met Leu
 885 890 895
 Gly Gly Asp Asn Gly Glu Thr Lys Phe Lys Gly Asn Asn Gln Ala Asp
 900 905 910
 Thr Val Asp Phe Ser Ile Ile Ser Val Ala Gly Lys Ala Leu Ala Pro
 915 920 925

 Ala Asp Leu Met Glu His His Ser Gly Ser Gln Gly Pro Leu Leu Thr
 930 935 940
 Thr Gly Asp Leu Gly Lys Glu Lys Thr Gln Lys Arg Val Lys Glu Gly
 945 950 955 960
 Asn Gly Thr Ser Asn Ser Thr Leu Ser Gly Pro Gly Leu Asp Ser Lys
 965 970 975
 Pro Gly Lys Arg Ser Arg Thr Pro Ser Asn Asp Gly Lys Ser Lys Asp
 980 985 990
 Lys Pro Pro Lys Arg Lys Lys Ala Asp Thr Glu Gly Lys Ser Pro Ser
 995 1000 1005
 His Ser Ser Ser Asn Arg Pro Phe Thr Pro Pro Thr Ser Thr Gly
 1010 1015 1020
 Gly Ser Lys Ser Pro Gly Ser Ala Gly Arg Ser Gln Thr Pro Pro
 1025 1030 1035
 Gly Val Ala Thr Pro Pro Ile Pro Lys Ile Thr Ile Gln Ile Pro
 1040 1045 1050
 Lys Gly Thr Val Met Val Gly Lys Pro Ser Ser His Ser Gln Tyr
 1055 1060 1065
 Thr Ser Ser Gly Ser Val Ser Ser Ser Gly Ser Lys Ser His His
 1070 1075 1080
 Ser His Ser Ser Ser Ser Ser Ser Ser Ala Ser Thr Ser Gly Lys
 1085 1090 1095
 Met Lys Ser Ser Lys Ser Glu Gly Ser Ser Ser Ser Lys Leu Ser
 1100 1105 1110
 Ser Ser Met Tyr Ser Ser Gln Gly Ser Ser Gly Ser Ser Gln Ser
 1115 1120 1125
 Lys Asn Ser Ser Gln Ser Gly Gly Lys Pro Gly Ser Ser Pro Ile
 1130 1135 1140
 Thr Lys His Gly Leu Ser Ser Gly Ser Ser Ser Thr Lys Met Lys
 1145 1150 1155
 Pro Gln Gly Lys Pro Ser Ser Leu Met Asn Pro Ser Leu Ser Lys
 1160 1165 1170
 Pro Asn Ile Ser Pro Ser His Ser Arg Pro Pro Gly Gly Ser Asp
 1175 1180 1185
 Lys Leu Ala Ser Pro Met Lys Pro Val Pro Gly Thr Pro Pro Ser
 1190 1195 1200
 Ser Lys Ala Lys Ser Pro Ile Ser Ser Gly Ser Gly Gly Ser His
 1205 1210 1215

	Met	Ser	Gly	Thr	Ser	Ser	Ser	Ser	Gly	Met	Lys	Ser	Ser	Ser	Gly
	1220						1225					1230			
5	Leu	Gly	Ser	Ser	Gly	Ser	Leu	Ser	Gln	Lys	Thr	Pro	Pro	Ser	Ser
	1235						1240					1245			
	Asn	Ser	Cys	Thr	Ala	Ser	Ser	Ser	Ser	Phe	Ser	Ser	Ser	Gly	Ser
	1250						1255					1260			
	Ser	Met	Ser	Ser	Ser	Gln	Asn	Gln	His	Gly	Ser	Ser	Lys	Gly	Lys
	1265						1270					1275			
10	Ser	Pro	Ser	Arg	Asn	Lys	Lys	Pro	Ser	Leu	Thr	Ala	Val	Ile	Asp
	1280						1285					1290			
	Lys	Leu	Lys	His	Gly	Val	Val	Thr	Ser	Gly	Pro	Gly	Gly	Glu	Asp
	1295						1300					1305			
	Pro	Leu	Asp	Gly	Gln	Met	Gly	Val	Ser	Thr	Asn	Ser	Ser	Ser	His
	1310						1315					1320			
15	Pro	Met	Ser	Ser	Lys	His	Asn	Met	Ser	Gly	Gly	Glu	Phe	Gln	Gly
	1325						1330					1335			
	Lys	Arg	Glu	Lys	Ser	Asp	Lys	Asp	Lys	Ser	Lys	Val	Ser	Thr	Ser
	1340						1345					1350			
	Gly	Ser	Ser	Val	Asp	Ser	Ser	Lys	Lys	Thr	Ser	Glu	Ser	Lys	Asn
	1355						1360					1365			
20	Val	Gly	Ser	Thr	Gly	Val	Ala	Lys	Ile	Ile	Ile	Ser	Lys	His	Asp
	1370						1375					1380			
	Gly	Gly	Ser	Pro	Ser	Ile	Lys	Ala	Lys	Val	Thr	Leu	Gln	Lys	Pro
	1385						1390					1395			
	Gly	Glu	Ser	Ser	Gly	Glu	Gly	Leu	Arg	Pro	Gln	Met	Ala	Ser	Ser
25	1400						1405					1410			
	Lys	Asn	Tyr	Gly	Ser	Pro	Leu	Ile	Ser	Gly	Ser	Thr	Pro	Lys	His
	1415						1420					1425			
	Glu	Arg	Gly	Ser	Pro	Ser	His	Ser	Lys	Ser	Pro	Ala	Tyr	Thr	Pro
	1430						1435					1440			
30	Gln	Asn	Leu	Asp	Ser	Glu	Ser	Glu	Ser	Gly	Ser	Ser	Ile	Ala	Glu
	1445						1450					1455			
	Lys	Ser	Tyr	Gln	Asn	Ser	Pro	Ser	Ser	Asp	Asp	Gly	Ile	Arg	Pro
	1460						1465					1470			
	Leu	Pro	Glu	Tyr	Ser	Thr	Glu	Lys	His	Lys	Lys	His	Lys	Lys	Glu
	1475						1480					1485			
35	Lys	Lys	Lys	Val	Lys	Asp	Lys	Asp	Arg	Asp	Arg	Asp	Arg	Asp	Lys
	1490						1495					1500			
	Asp	Arg	Asp	Lys	Lys	Lys	Ser	His	Ser	Ile	Lys	Pro	Glu	Ser	Trp
	1505						1510					1515			
	Ser	Lys	Ser	Pro	Ile	Ser	Ser	Asp	Gln	Ser	Leu	Ser	Met	Thr	Ser
	1520						1525					1530			
40	Asn	Thr	Ile	Leu	Ser	Ala	Asp	Arg	Pro	Ser	Arg	Leu	Ser	Pro	Asp
	1535						1540					1545			
	Phe	Met	Ile	Gly	Glu	Glu	Asp	Asp	Asp	Leu	Met	Asp	Val	Ala	Leu
	1550						1555					1560			
45	Ile	Gly	Asn												
	1565														

<210> 31

<211> 1490

50

<212> PRT

<213> Homo sapiens

55

<400> 31
 Met Pro Asn Ser Glu Arg His Gly Gly Lys Lys Asp Gly Ser Gly Gly
 1 5 10 15
 5 Ala Ser Gly Thr Leu Gln Pro Ser Ser Gly Gly Gly Ser Ser Asn Ser
 20 25 30
 Arg Glu Arg His Arg Leu Val Ser Lys His Lys Arg His Lys Ser Lys
 35 40 45
 His Ser Lys Asp Met Gly Leu Val Thr Pro Glu Ala Ala Ser Leu Gly
 50 55 60
 10 Thr Val Ile Lys Pro Leu Val Glu Tyr Asp Asp Ile Ser Ser Asp Ser
 65 70 75 80
 Asp Thr Phe Ser Asp Asp Met Ala Phe Lys Leu Asp Arg Arg Glu Asn
 85 90 95
 15 Asp Glu Arg Arg Gly Ser Asp Arg Ser Asp Arg Leu His Lys His Arg
 100 105 110
 His His Gln His Arg Arg Ser Arg Asp Leu Leu Lys Ala Lys Gln Thr
 115 120 125
 Glu Lys Glu Lys Ser Gln Glu Val Ser Ser Lys Ser Gly Ser Met Lys
 130 135 140
 20 Asp Arg Ile Ser Gly Ser Ser Lys Arg Ser Asn Glu Glu Thr Asp Asp
 145 150 155 160
 Tyr Gly Lys Ala Gln Val Ala Lys Ser Ser Ser Lys Glu Ser Arg Ser
 165 170 175
 Ser Lys Leu His Lys Glu Lys Thr Arg Lys Glu Arg Glu Leu Lys Ser
 180 185 190
 25 Gly His Lys Asp Arg Ser Lys Ser His Arg Lys Arg Glu Thr Pro Lys
 195 200 205
 Ser Tyr Lys Thr Val Asp Ser Pro Lys Arg Arg Ser Arg Ser Pro His
 210 215 220
 Arg Lys Trp Ser Asp Ser Ser Lys Gln Asp Asp Ser Pro Ser Gly Ala
 225 230 235 240
 30 Ser Tyr Gly Gln Asp Tyr Asp Leu Ser Pro Ser Arg Ser His Thr Ser
 245 250 255
 Ser Asn Tyr Asp Ser Tyr Lys Lys Ser Pro Gly Ser Thr Ser Arg Arg
 260 265 270
 35 Gln Ser Val Ser Pro Pro Tyr Lys Glu Pro Ser Ala Tyr Gln Ser Ser
 275 280 285
 Thr Arg Ser Pro Ser Pro Tyr Ser Arg Arg Gln Arg Ser Val Ser Pro
 290 295 300
 Tyr Ser Arg Arg Arg Ser Ser Ser Tyr Glu Arg Ser Gly Ser Tyr Ser
 305 310 315 320
 40 Gly Arg Ser Pro Ser Pro Tyr Gly Arg Arg Arg Ser Ser Ser Pro Phe
 325 330 335
 Leu Ser Lys Arg Ser Leu Ser Arg Ser Pro Leu Pro Ser Arg Lys Ser
 340 345 350
 Met Lys Ser Arg Ser Arg Ser Pro Ala Tyr Ser Arg His Ser Ser Ser
 355 360 365
 45 His Ser Lys Lys Lys Arg Ser Ser Ser Arg Ser Arg His Ser Ser Ile
 370 375 380
 Ser Pro Val Arg Leu Pro Leu Asn Ser Ser Leu Gly Ala Glu Leu Ser
 385 390 395 400
 Arg Lys Lys Lys Glu Arg Ala Ala Ala Ala Ala Ala Lys Met Asp
 405 410 415
 50 Gly Lys Glu Ser Lys Gly Ser Pro Val Phe Leu Pro Arg Lys Glu Asn
 420 425 430
 Ser Ser Val Glu Ala Lys Asp Ser Gly Leu Glu Ser Lys Lys Leu Pro
 435 440 445
 55 Arg Ser Val Lys Leu Glu Lys Ser Ala Pro Asp Thr Glu Leu Val Asn
 450 455 460

Val Thr His Leu Asn Thr Glu Val Lys Asn Ser Ser Asp Thr Gly Lys
 465 470 475 480
 Val Lys Leu Asp Glu Asn Ser Glu Lys His Leu Val Lys Asp Leu Lys
 485 490 495
 Ala Gln Gly Thr Arg Asp Ser Lys Pro Ile Ala Leu Lys Glu Glu Ile
 500 505 510
 Val Thr Pro Lys Glu Thr Glu Thr Ser Glu Lys Glu Thr Pro Pro Pro
 515 520 525
 Leu Pro Thr Ile Ala Ser Pro Pro Pro Leu Pro Thr Thr Thr Pro
 530 535 540
 Pro Pro Gln Thr Pro Pro Leu Pro Pro Leu Pro Pro Ile Pro Ala Leu
 545 550 555 560
 Pro Gln Gln Pro Pro Leu Pro Pro Ser Gln Pro Ala Phe Ser Gln Val
 565 570 575
 Pro Ala Ser Ser Thr Ser Thr Leu Pro Ser Thr His Ser Lys Thr
 580 585 590
 Ser Ala Val Ser Ser Gln Ala Asn Ser Gln Pro Pro Val Gln Val Ser
 595 600 605
 Val Lys Thr Gln Val Ser Val Thr Ala Ala Ile Pro His Leu Lys Thr
 610 615 620
 Ser Thr Leu Pro Pro Leu Pro Leu Pro Pro Leu Pro Gly Gly Asp
 625 630 635 640
 Asp Met Asp Ser Pro Lys Glu Thr Leu Pro Ser Lys Pro Val Lys Lys
 645 650 655
 Glu Lys Glu Gln Arg Thr Arg His Leu Leu Thr Asp Leu Pro Leu Pro
 660 665 670
 Pro Glu Leu Pro Gly Gly Asp Leu Ser Pro Pro Asp Ser Pro Glu Pro
 675 680 685
 Lys Ala Ile Thr Pro Pro Gln Gln Pro Tyr Lys Lys Arg Pro Lys Ile
 690 695 700
 Cys Cys Pro Arg Tyr Gly Glu Arg Arg Gln Thr Glu Ser Asp Trp Gly
 705 710 715 720
 Lys Arg Cys Val Asp Lys Phe Asp Ile Ile Gly Ile Ile Gly Glu Gly
 725 730 735
 Thr Tyr Gly Gln Val Tyr Lys Ala Arg Asp Lys Asp Thr Gly Glu Leu
 740 745 750
 Val Ala Leu Lys Lys Val Arg Leu Asp Asn Glu Lys Glu Gly Phe Pro
 755 760 765
 Ile Thr Ala Ile Arg Glu Ile Lys Ile Leu Arg Gln Leu Ile His Arg
 770 775 780
 Ser Val Val Asn Met Lys Glu Ile Val Thr Asp Lys Gln Asp Ala Leu
 785 790 795 800
 Asp Phe Lys Lys Asp Lys Gly Ala Phe Tyr Leu Val Phe Glu Tyr Met
 805 810 815
 Asp His Asp Leu Met Gly Leu Leu Glu Ser Gly Leu Val His Phe Ser
 820 825 830
 Glu Asp His Ile Lys Ser Phe Met Lys Gln Leu Met Glu Gly Leu Glu
 835 840 845
 Tyr Cys His Lys Lys Asn Phe Leu His Arg Asp Ile Lys Cys Ser Asn
 850 855 860
 Ile Leu Leu Asn Asn Ser Gly Gln Ile Lys Leu Ala Asp Phe Gly Leu
 865 870 875 880
 Ala Arg Leu Tyr Asn Ser Glu Glu Ser Arg Pro Tyr Thr Asn Lys Val
 885 890 895
 Ile Thr Leu Trp Tyr Arg Pro Pro Glu Leu Leu Leu Gly Glu Glu Arg
 900 905 910
 Tyr Thr Pro Ala Ile Asp Val Trp Ser Cys Gly Cys Ile Leu Gly Glu
 915 920 925

Leu Phe Thr Lys Lys Pro Ile Phe Gln Ala Asn Leu Glu Leu Ala Gln
 930 935 940
 Leu Glu Leu Ile Ser Arg Leu Cys Gly Ser Pro Cys Pro Ala Val Trp
 5 945 950 955 960
 Pro Asp Val Ile Lys Leu Pro Tyr Phe Asn Thr Met Lys Pro Lys Lys
 965 970 975
 Gln Tyr Arg Arg Arg Leu Arg Glu Glu Phe Ser Phe Ile Pro Ser Ala
 980 985 990
 10 Ala Leu Asp Leu Leu Asp His Met Leu Thr Leu Asp Pro Ser Lys Arg
 995 1000 1005
 Cys Thr Ala Glu Gln Thr Leu Gln Ser Asp Phe Leu Lys Asp Val
 1010 1015 1020
 Glu Leu Ser Lys Met Ala Pro Pro Asp Leu Pro His Trp Gln Asp
 1025 1030 1035
 15 Cys His Glu Leu Trp Ser Lys Lys Arg Arg Arg Gln Arg Gln Ser
 1040 1045 1050
 Gly Val Val Val Glu Glu Pro Pro Pro Ser Lys Thr Ser Arg Lys
 1055 1060 1065
 Glu Thr Thr Ser Gly Thr Ser Thr Glu Pro Val Lys Asn Ser Ser
 1070 1075 1080
 20 Pro Ala Pro Pro Gln Pro Ala Pro Gly Lys Val Glu Ser Gly Ala
 1085 1090 1095
 Gly Asp Ala Ile Gly Leu Ala Asp Ile Thr Gln Gln Leu Asn Gln
 1100 1105 1110
 Ser Glu Leu Ala Val Leu Leu Asn Leu Leu Gln Ser Gln Thr Asp
 25 1115 1120 1125
 Leu Ser Ile Pro Gln Met Ala Gln Leu Leu Asn Ile His Ser Asn
 1130 1135 1140
 Pro Glu Met Gln Gln Gln Leu Glu Ala Leu Asn Gln Ser Ile Ser
 1145 1150 1155
 30 Ala Leu Thr Glu Ala Thr Ser Gln Gln Gln Asp Ser Glu Thr Met
 1160 1165 1170
 Ala Pro Glu Glu Ser Leu Lys Glu Ala Pro Ser Ala Pro Val Ile
 1175 1180 1185
 Leu Pro Ser Ala Glu Gln Met Thr Leu Glu Ala Ser Ser Thr Pro
 1190 1195 1200
 35 Ala Asp Met Gln Asn Ile Leu Ala Val Leu Leu Ser Gln Leu Met
 1205 1210 1215
 Lys Thr Gln Glu Pro Ala Gly Ser Leu Glu Glu Asn Asn Ser Asp
 1220 1225 1230
 Lys Asn Ser Gly Pro Gln Gly Pro Arg Arg Thr Pro Thr Met Pro
 1235 1240 1245
 40 Gln Glu Glu Ala Ala Ala Cys Pro Pro His Ile Leu Pro Pro Glu
 1250 1255 1260
 Lys Arg Pro Pro Glu Pro Pro Gly Pro Pro Pro Pro Pro Pro Pro
 1265 1270 1275
 Pro Pro Leu Val Glu Gly Asp Leu Ser Ser Ala Pro Gln Glu Leu
 45 1280 1285 1290
 Asn Pro Ala Val Thr Ala Ala Leu Leu Gln Leu Leu Ser Gln Pro
 1295 1300 1305
 Glu Ala Glu Pro Pro Gly His Leu Pro His Glu His Gln Ala Leu
 1310 1315 1320
 Arg Pro Met Glu Tyr Ser Thr Arg Pro Arg Pro Asn Arg Thr Tyr
 50 1325 1330 1335
 Gly Asn Thr Asp Gly Pro Glu Thr Gly Phe Ser Ala Ile Asp Thr
 1340 1345 1350
 Asp Glu Arg Asn Ser Gly Pro Ala Leu Thr Glu Ser Leu Val Gln
 1355 1360 1365

Thr Leu Val Lys Asn Arg Thr Phe Ser Gly Ser Leu Ser His Leu
 1370 1375 1380
 Gly Glu Ser Ser Ser Tyr Gln Gly Thr Gly Ser Val Gln Phe Pro
 1385 1390 1395
 Gly Asp Gln Asp Leu Arg Phe Ala Arg Val Pro Leu Ala Leu His
 1400 1405 1410
 Pro Val Val Gly Gln Pro Phe Leu Lys Ala Glu Gly Ser Ser Asn
 1415 1420 1425
 Ser Val Val His Ala Glu Thr Lys Leu Gln Asn Tyr Gly Glu Leu
 1430 1435 1440
 Gly Pro Gly Thr Thr Gly Ala Ser Ser Ser Gly Ala Gly Leu His
 1445 1450 1455
 Trp Gly Gly Pro Thr Gln Ser Ser Ala Tyr Gly Lys Leu Tyr Arg
 1460 1465 1470
 Gly Pro Thr Arg Val Pro Pro Arg Gly Gly Arg Gly Arg Gly Val
 1475 1480 1485
 Pro Tyr
 1490

 <210> 32

 <211> 381

 <212> PRT

 <213> Homo sapiens

 <400> 32
 Met Leu Thr Arg Leu Phe Ser Glu Pro Gly Leu Leu Ser Asp Val Pro
 1 5 10 15
 Lys Phe Ala Ser Trp Gly Asp Gly Glu Asp Asp Glu Pro Arg Ser Asp
 20 25 30
 Lys Gly Asp Ala Pro Pro Pro Pro Pro Pro Ala Pro Gly Pro Gly Ala
 35 40 45
 Pro Gly Pro Ala Arg Ala Ala Lys Pro Val Pro Leu Arg Gly Glu Glu
 50 55 60
 Gly Thr Glu Ala Thr Leu Ala Glu Val Lys Glu Glu Gly Glu Leu Gly
 65 70 75 80
 Gly Glu Glu Glu Glu Glu Glu Glu Glu Glu Glu Gly Leu Asp Glu Ala
 85 90 95
 Glu Gly Glu Arg Pro Lys Lys Arg Gly Pro Lys Lys Arg Lys Met Thr
 100 105 110
 Lys Ala Arg Leu Glu Arg Ser Lys Leu Arg Arg Gln Lys Ala Asn Ala
 115 120 125
 Arg Glu Arg Asn Arg Met His Asp Leu Asn Ala Ala Leu Asp Asn Leu
 130 135 140
 Arg Lys Val Val Pro Cys Tyr Ser Lys Thr Gln Lys Leu Ser Lys Ile
 145 150 155 160
 Glu Thr Leu Arg Leu Ala Lys Asn Tyr Ile Trp Ala Leu Ser Glu Ile
 165 170 175
 Leu Arg Ser Gly Lys Arg Pro Asp Leu Val Ser Tyr Val Gln Thr Leu
 180 185 190
 Cys Lys Gly Leu Ser Gln Pro Thr Thr Asn Leu Val Ala Gly Cys Leu
 195 200 205
 Gln Leu Asn Ser Arg Asn Phe Leu Thr Glu Gln Gly Ala Asp Gly Ala
 210 215 220

Gly Arg Phe His Gly Ser Gly Gly Pro Phe Ala Met His Pro Tyr Pro
 225 230 235 240
 Tyr Pro Cys Ser Arg Leu Ala Gly Ala Gln Cys Gln Ala Ala Gly Gly
 245 250 255
 Leu Gly Gly Gly Ala Ala His Ala Leu Arg Thr His Gly Tyr Cys Ala
 260 265 270
 Ala Tyr Glu Thr Leu Tyr Ala Ala Gly Gly Gly Gly Ala Ser Pro
 275 280 285
 Asp Tyr Asn Ser Ser Glu Tyr Glu Gly Pro Leu Ser Pro Pro Leu Cys
 290 295 300
 Leu Asn Gly Asn Phe Ser Leu Lys Gln Asp Ser Ser Pro Asp His Glu
 305 310 315 320
 Lys Ser Tyr His Tyr Ser Met His Tyr Ser Ala Leu Pro Gly Ser Arg
 325 330 335
 His Gly His Gly Leu Val Phe Gly Ser Ser Ala Val Arg Gly Gly Val
 340 345 350
 His Ser Glu Asn Leu Leu Ser Tyr Asp Met His Leu His His Asp Arg
 355 360 365
 Gly Pro Met Tyr Glu Glu Leu Asn Ala Phe Phe His Asn
 370 375 380

 <210> 33

 <211> 445

 <212> PRT

 <213> Homo sapiens

 <400> 33
 Met Ser Lys Leu Pro Arg Glu Leu Thr Arg Asp Leu Glu Arg Ser Leu
 1 5 10 15
 Pro Ala Val Ala Ser Leu Gly Ser Ser Leu Ser His Ser Gln Ser Leu
 20 25 30
 Ser Ser His Leu Leu Pro Pro Glu Lys Arg Arg Ala Ile Ser Asp
 35 40 45
 Val Arg Arg Thr Phe Cys Leu Phe Val Thr Phe Asp Leu Leu Phe Ile
 50 55 60
 Ser Leu Leu Trp Ile Ile Glu Leu Asn Thr Asn Thr Gly Ile Arg Lys
 65 70 75 80
 Asn Leu Glu Gln Glu Ile Ile Gln Tyr Asn Phe Lys Thr Ser Phe Phe
 85 90 95
 Asp Ile Phe Val Leu Ala Phe Phe Arg Phe Ser Gly Leu Leu Leu Gly
 100 105 110
 Tyr Ala Val Leu Gln Leu Arg His Trp Trp Val Ile Ala Val Thr Thr
 115 120 125
 Leu Val Ser Ser Ala Phe Leu Ile Val Lys Val Ile Leu Ser Glu Leu
 130 135 140
 Leu Ser Lys Gly Ala Phe Gly Tyr Leu Leu Pro Ile Val Ser Phe Val
 145 150 155 160
 Leu Ala Trp Leu Glu Thr Trp Phe Leu Asp Phe Lys Val Leu Pro Gln
 165 170 175
 Glu Ala Glu Glu Glu Arg Trp Tyr Leu Ala Ala Gln Val Ala Val Ala
 180 185 190
 Arg Gly Pro Leu Leu Phe Ser Gly Ala Leu Ser Glu Gly Gln Phe Tyr
 195 200 205

Asp Arg Gln Glu Val Ala Glu Ile Thr Lys Gln Leu Pro Pro Val Val
 130 135 140
 Pro Val Ser Lys Pro Gly Ala Leu Arg Arg Ser Leu Ser Arg Ser Met
 145 150 155 160
 Ser Gln Glu Ala Gln Arg Gly
 165

<210> 35

<211> 282

<212> PRT

<213> Homo sapiens

<400> 35

Met Ser Gly Ala Asp Arg Ser Pro Asn Ala Gly Ala Ala Pro Asp Ser
 1 5 10 15
 Ala Pro Gly Gln Ala Ala Val Ala Ser Ala Tyr Gln Arg Phe Glu Pro
 20 20 25 30
 Arg Ala Tyr Leu Arg Asn Asn Tyr Ala Pro Pro Arg Gly Asp Leu Cys
 35 40 45
 Asn Pro Asn Gly Val Gly Pro Trp Lys Leu Arg Cys Leu Ala Gln Thr
 25 50 55 60
 Phe Ala Thr Gly Glu Val Ser Gly Arg Thr Leu Ile Asp Ile Gly Ser
 65 70 75 80
 Gly Pro Thr Val Tyr Gln Leu Leu Ser Ala Cys Ser His Phe Glu Asp
 85 90 95
 Ile Thr Met Thr Asp Phe Leu Glu Val Asn Arg Gln Glu Leu Gly Arg
 30 100 105 110
 Trp Leu Gln Glu Glu Pro Gly Ala Phe Asn Trp Ser Met Tyr Ser Gln
 115 120 125
 His Ala Cys Leu Ile Glu Gly Lys Gly Glu Cys Trp Gln Asp Lys Glu
 130 135 140
 Arg Gln Leu Arg Ala Arg Val Lys Arg Val Leu Pro Ile Asp Val His
 35 145 150 155 160
 Gln Pro Gln Pro Leu Gly Ala Gly Ser Pro Ala Pro Leu Pro Ala Asp
 165 170 175
 Ala Leu Val Ser Ala Phe Cys Leu Glu Ala Val Ser Pro Asp Leu Ala
 40 180 185 190
 Ser Phe Gln Arg Ala Leu Asp His Ile Thr Thr Leu Leu Arg Pro Gly
 195 200 205
 Gly His Leu Leu Leu Ile Gly Ala Leu Glu Glu Ser Trp Tyr Leu Ala
 210 215 220
 Gly Glu Ala Arg Leu Thr Val Val Pro Val Ser Glu Glu Glu Val Arg
 45 225 230 235 240
 Glu Ala Leu Val Arg Ser Gly Tyr Lys Val Arg Asp Leu Arg Thr Tyr
 245 250 255
 Ile Met Pro Ala His Leu Gln Thr Gly Val Asp Asp Val Lys Gly Val
 260 265 270
 Phe Phe Ala Trp Ala Gln Lys Val Gly Leu
 275 280

<210> 36

<211> 1255

<212> PRT

<213> Homo sapiens

<400> 36

Met Glu Leu Ala Ala Leu Cys Arg Trp Gly Leu Leu Leu Ala Leu Leu
 1 5 10 15
 Pro Pro Gly Ala Ala Ser Thr Gln Val Cys Thr Gly Thr Asp Met Lys
 20 25 30
 Leu Arg Leu Pro Ala Ser Pro Glu Thr His Leu Asp Met Leu Arg His
 35 40 45
 Leu Tyr Gln Gly Cys Gln Val Val Gln Gly Asn Leu Glu Leu Thr Tyr
 50 55 60
 Leu Pro Thr Asn Ala Ser Leu Ser Phe Leu Gln Asp Ile Gln Glu Val
 65 70 75 80
 Gln Gly Tyr Val Leu Ile Ala His Asn Gln Val Arg Gln Val Pro Leu
 85 90 95
 Gln Arg Leu Arg Ile Val Arg Gly Thr Gln Leu Phe Glu Asp Asn Tyr
 100 105 110
 Ala Leu Ala Val Leu Asp Asn Gly Asp Pro Leu Asn Asn Thr Thr Pro
 115 120 125
 Val Thr Gly Ala Ser Pro Gly Gly Leu Arg Glu Leu Gln Leu Arg Ser
 130 135 140
 Leu Thr Glu Ile Leu Lys Gly Gly Val Leu Ile Gln Arg Asn Pro Gln
 145 150 155 160
 Leu Cys Tyr Gln Asp Thr Ile Leu Trp Lys Asp Ile Phe His Lys Asn
 165 170 175
 Asn Gln Leu Ala Leu Thr Leu Ile Asp Thr Asn Arg Ser Arg Ala Cys
 180 185 190
 His Pro Cys Ser Pro Met Cys Lys Gly Ser Arg Cys Trp Gly Glu Ser
 195 200 205
 Ser Glu Asp Cys Gln Ser Leu Thr Arg Thr Val Cys Ala Gly Gly Cys
 210 215 220
 Ala Arg Cys Lys Gly Pro Leu Pro Thr Asp Cys Cys His Glu Gln Cys
 225 230 235 240
 Ala Ala Gly Cys Thr Gly Pro Lys His Ser Asp Cys Leu Ala Cys Leu
 245 250 255
 His Phe Asn His Ser Gly Ile Cys Glu Leu His Cys Pro Ala Leu Val
 260 265 270
 Thr Tyr Asn Thr Asp Thr Phe Glu Ser Met Pro Asn Pro Glu Gly Arg
 275 280 285
 Tyr Thr Phe Gly Ala Ser Cys Val Thr Ala Cys Pro Tyr Asn Tyr Leu
 290 295 300
 Ser Thr Asp Val Gly Ser Cys Thr Leu Val Cys Pro Leu His Asn Gln
 305 310 315 320
 Glu Val Thr Ala Glu Asp Gly Thr Gln Arg Cys Glu Lys Cys Ser Lys
 325 330 335
 Pro Cys Ala Arg Val Cys Tyr Gly Leu Gly Met Glu His Leu Arg Glu
 340 345 350
 Val Arg Ala Val Thr Ser Ala Asn Ile Gln Glu Phe Ala Gly Cys Lys
 355 360 365
 Lys Ile Phe Gly Ser Leu Ala Phe Leu Pro Glu Ser Phe Asp Gly Asp
 370 375 380

Pro Ala Ser Asn Thr Ala Pro Leu Gln Pro Glu Gln Leu Gln Val Phe
 385 390 395 400
 Glu Thr Leu Glu Glu Ile Thr Gly Tyr Leu Tyr Ile Ser Ala Trp Pro
 405 410 415
 Asp Ser Leu Pro Asp Leu Ser Val Phe Gln Asn Leu Gln Val Ile Arg
 420 425 430

 Gly Arg Ile Leu His Asn Gly Ala Tyr Ser Leu Thr Leu Gln Gly Leu
 435 440 445
 Gly Ile Ser Trp Leu Gly Leu Arg Ser Leu Arg Glu Leu Gly Ser Gly
 450 455 460
 Leu Ala Leu Ile His His Asn Thr His Leu Cys Phe Val His Thr Val
 465 470 475 480
 Pro Trp Asp Gln Leu Phe Arg Asn Pro His Gln Ala Leu Leu His Thr
 485 490 495
 Ala Asn Arg Pro Glu Asp Glu Cys Val Gly Glu Gly Leu Ala Cys His
 500 505 510
 Gln Leu Cys Ala Arg Gly His Cys Trp Gly Pro Gly Pro Thr Gln Cys
 515 520 525
 Val Asn Cys Ser Gln Phe Leu Arg Gly Gln Glu Cys Val Glu Glu Cys
 530 535 540
 Arg Val Leu Gln Gly Leu Pro Arg Glu Tyr Val Asn Ala Arg His Cys
 545 550 555 560
 Leu Pro Cys His Pro Glu Cys Gln Pro Gln Asn Gly Ser Val Thr Cys
 565 570 575
 Phe Gly Pro Glu Ala Asp Gln Cys Val Ala Cys Ala His Tyr Lys Asp
 580 585 590
 Pro Pro Phe Cys Val Ala Arg Cys Pro Ser Gly Val Lys Pro Asp Leu
 595 600 605
 Ser Tyr Met Pro Ile Trp Lys Phe Pro Asp Glu Glu Gly Ala Cys Gln
 610 615 620
 Pro Cys Pro Ile Asn Cys Thr His Ser Cys Val Asp Leu Asp Asp Lys
 625 630 635 640
 Gly Cys Pro Ala Glu Gln Arg Ala Ser Pro Leu Thr Ser Ile Val Ser
 645 650 655
 Ala Val Val Gly Ile Leu Leu Val Val Val Leu Gly Val Val Phe Gly
 660 665 670
 Ile Leu Ile Lys Arg Arg Gln Gln Lys Ile Arg Lys Tyr Thr Met Arg
 675 680 685
 Arg Leu Leu Gln Glu Thr Glu Leu Val Glu Pro Leu Thr Pro Ser Gly
 690 695 700
 Ala Met Pro Asn Gln Ala Gln Met Arg Ile Leu Lys Glu Thr Glu Leu
 705 710 715 720
 Arg Lys Val Lys Val Leu Gly Ser Gly Ala Phe Gly Thr Val Tyr Lys
 725 730 735
 Gly Ile Trp Ile Pro Asp Gly Glu Asn Val Lys Ile Pro Val Ala Ile
 740 745 750
 Lys Val Leu Arg Glu Asn Thr Ser Pro Lys Ala Asn Lys Glu Ile Leu
 755 760 765
 Asp Glu Ala Tyr Val Met Ala Gly Val Gly Ser Pro Tyr Val Ser Arg
 770 775 780
 Leu Leu Gly Ile Cys Leu Thr Ser Thr Val Gln Leu Val Thr Gln Leu
 785 790 795 800
 Met Pro Tyr Gly Cys Leu Leu Asp His Val Arg Glu Asn Arg Gly Arg
 805 810 815
 Leu Gly Ser Gln Asp Leu Leu Asn Trp Cys Met Gln Ile Ala Lys Gly
 820 825 830
 Met Ser Tyr Leu Glu Asp Val Arg Leu Val His Arg Asp Leu Ala Ala
 835 840 845

Arg Asn Val Leu Val Lys Ser Pro Asn His Val Lys Ile Thr Asp Phe
 850 855 860
 Gly Leu Ala Arg Leu Leu Asp Ile Asp Glu Thr Glu Tyr His Ala Asp
 865 870 875 880
 Gly Gly Lys Val Pro Ile Lys Trp Met Ala Leu Glu Ser Ile Leu Arg
 885 890 895
 Arg Arg Phe Thr His Gln Ser Asp Val Trp Ser Tyr Gly Val Thr Val
 900 905 910
 Trp Glu Leu Met Thr Phe Gly Ala Lys Pro Tyr Asp Gly Ile Pro Ala
 915 920 925
 Arg Glu Ile Pro Asp Leu Leu Glu Lys Gly Glu Arg Leu Pro Gln Pro
 930 935 940
 Pro Ile Cys Thr Ile Asp Val Tyr Met Ile Met Val Lys Cys Trp Met
 945 950 955 960
 Ile Asp Ser Glu Cys Arg Pro Arg Phe Arg Glu Leu Val Ser Glu Phe
 965 970 975
 Ser Arg Met Ala Arg Asp Pro Gln Arg Phe Val Val Ile Gln Asn Glu
 980 985 990
 Asp Leu Gly Pro Ala Ser Pro Leu Asp Ser Thr Phe Tyr Arg Ser Leu
 995 1000 1005
 Leu Glu Asp Asp Asp Met Gly Asp Leu Val Asp Ala Glu Glu Tyr
 1010 1015 1020
 Leu Val Pro Gln Gln Gly Phe Phe Cys Pro Asp Pro Ala Pro Gly
 1025 1030 1035
 Ala Gly Gly Met Val His His Arg His Arg Ser Ser Ser Thr Arg
 1040 1045 1050
 Ser Gly Gly Gly Asp Leu Thr Leu Gly Leu Glu Pro Ser Glu Glu
 1055 1060 1065
 Glu Ala Pro Arg Ser Pro Leu Ala Pro Ser Glu Gly Ala Gly Ser
 1070 1075 1080
 Asp Val Phe Asp Gly Asp Leu Gly Met Gly Ala Ala Lys Gly Leu
 1085 1090 1095
 Gln Ser Leu Pro Thr His Asp Pro Ser Pro Leu Gln Arg Tyr Ser
 1100 1105 1110
 Glu Asp Pro Thr Val Pro Leu Pro Ser Glu Thr Asp Gly Tyr Val
 1115 1120 1125
 Ala Pro Leu Thr Cys Ser Pro Gln Pro Glu Tyr Val Asn Gln Pro
 1130 1135 1140
 Asp Val Arg Pro Gln Pro Pro Ser Pro Arg Glu Gly Pro Leu Pro
 1145 1150 1155
 Ala Ala Arg Pro Ala Gly Ala Thr Leu Glu Arg Ala Lys Thr Leu
 1160 1165 1170
 Ser Pro Gly Lys Asn Gly Val Val Lys Asp Val Phe Ala Phe Gly
 1175 1180 1185
 Gly Ala Val Glu Asn Pro Glu Tyr Leu Thr Pro Gln Gly Gly Ala
 1190 1195 1200
 Ala Pro Gln Pro His Pro Pro Pro Ala Phe Ser Pro Ala Phe Asp
 1205 1210 1215
 Asn Leu Tyr Tyr Trp Asp Gln Asp Pro Pro Glu Arg Gly Ala Pro
 1220 1225 1230
 Pro Ser Thr Phe Lys Gly Thr Pro Thr Ala Glu Asn Pro Glu Tyr
 1235 1240 1245
 Leu Gly Leu Asp Val Pro Val
 1250 1255

<210> 37

<211> 532

<212> PRT

<213> Homo sapiens

<400> 37

Met	Glu	Leu	Asp	Leu	Ser	Pro	Pro	His	Leu	Ser	Ser	Ser	Pro	Glu	Asp
1			5						10					15	
Leu	Trp	Pro	Ala	Pro	Gly	Thr	Pro	Pro	Gly	Thr	Pro	Arg	Pro	Pro	Asp
			20					25					30		
Thr	Pro	Leu	Pro	Glu	Glu	Val	Lys	Arg	Ser	Gln	Pro	Leu	Leu	Ile	Pro
		35					40					45			
Thr	Thr	Gly	Arg	Lys	Leu	Arg	Glu	Glu	Glu	Arg	Arg	Ala	Thr	Ser	Leu
	50					55					60				
Pro	Ser	Ile	Pro	Asn	Pro	Phe	Pro	Glu	Leu	Cys	Ser	Pro	Pro	Ser	Gln
65				70						75					80
Ser	Pro	Ile	Leu	Gly	Gly	Pro	Ser	Ser	Ala	Arg	Gly	Leu	Leu	Pro	Arg
			85						90					95	
Asp	Ala	Ser	Arg	Pro	His	Val	Val	Lys	Val	Tyr	Ser	Glu	Asp	Gly	Ala
			100					105					110		
Cys	Arg	Ser	Val	Glu	Val	Ala	Ala	Gly	Ala	Thr	Ala	Arg	His	Val	Cys
		115					120					125			
Glu	Met	Leu	Val	Gln	Arg	Ala	His	Ala	Leu	Ser	Asp	Glu	Thr	Trp	Gly
	130					135					140				
Leu	Val	Glu	Cys	His	Pro	His	Leu	Ala	Leu	Glu	Arg	Gly	Leu	Glu	Asp
145					150					155					160
His	Glu	Ser	Val	Val	Glu	Val	Gln	Ala	Ala	Trp	Pro	Val	Gly	Gly	Asp
			165						170					175	
Ser	Arg	Phe	Val	Phe	Arg	Lys	Asn	Phe	Ala	Lys	Tyr	Glu	Leu	Phe	Lys
		180						185					190		
Ser	Ser	Pro	His	Ser	Leu	Phe	Pro	Glu	Lys	Met	Val	Ser	Ser	Cys	Leu
		195					200					205			
Asp	Ala	His	Thr	Gly	Ile	Ser	His	Glu	Asp	Leu	Ile	Gln	Asn	Phe	Leu
	210					215						220			
Asn	Ala	Gly	Ser	Phe	Pro	Glu	Ile	Gln	Gly	Phe	Leu	Gln	Leu	Arg	Gly
225				230						235					240
Ser	Gly	Arg	Lys	Leu	Trp	Lys	Arg	Phe	Phe	Cys	Phe	Leu	Arg	Arg	Ser
			245						250					255	
Gly	Leu	Tyr	Tyr	Ser	Thr	Lys	Gly	Thr	Ser	Lys	Asp	Pro	Arg	His	Leu
		260						265					270		
Gln	Tyr	Val	Ala	Asp	Val	Asn	Glu	Ser	Asn	Val	Tyr	Val	Val	Thr	Gln
		275					280					285			
Gly	Arg	Lys	Leu	Tyr	Gly	Met	Pro	Thr	Asp	Phe	Gly	Phe	Cys	Val	Lys
	290					295					300				
Pro	Asn	Lys	Leu	Arg	Asn	Gly	His	Lys	Gly	Leu	Arg	Ile	Phe	Cys	Ser
305					310					315					320
Glu	Asp	Glu	Gln	Ser	Arg	Thr	Cys	Trp	Leu	Ala	Ala	Phe	Arg	Leu	Phe
			325						330					335	
Lys	Tyr	Gly	Val	Gln	Leu	Tyr	Lys	Asn	Tyr	Gln	Gln	Ala	Gln	Ser	Arg
		340						345					350		
His	Leu	His	Pro	Ser	Cys	Leu	Gly	Ser	Pro	Pro	Leu	Arg	Ser	Ala	Ser
		355					360					365			
Asp	Asn	Thr	Leu	Val	Ala	Met	Asp	Phe	Ser	Gly	His	Ala	Gly	Arg	Val
	370					375					380				

Ile Glu Asn Pro Arg Glu Ala Leu Ser Val Ala Leu Glu Glu Ala Gln
 385 390 395 400
 Ala Trp Arg Lys Lys Thr Asn His Arg Leu Ser Leu Pro Met Pro Ala
 405 410 415
 Ser Gly Thr Ser Leu Ser Ala Ala Ile His Arg Thr Gln Leu Trp Phe
 420 425 430
 His Gly Arg Ile Ser Arg Glu Glu Ser Gln Arg Leu Ile Gly Gln Gln
 435 440 445
 Gly Leu Val Asp Gly Leu Phe Leu Val Arg Glu Ser Gln Arg Asn Pro
 450 455 460
 Gln Gly Phe Val Leu Ser Leu Cys His Leu Gln Lys Val Lys His Tyr
 465 470 475 480
 Leu Ile Leu Pro Ser Glu Glu Glu Gly Arg Leu Tyr Phe Ser Met Asp
 485 490 495
 Asp Gly Gln Thr Arg Phe Thr Asp Leu Leu Gln Leu Val Glu Phe His
 500 505 510
 Gln Leu Asn Arg Gly Ile Leu Pro Cys Leu Leu Arg His Cys Cys Thr
 515 520 525
 Arg Val Ala Leu
 530
 <210> 38
 <211> 534
 <212> PRT
 <213> Homo sapiens
 <400> 38
 Met Lys Gln Glu Gly Ser Ala Arg Arg Arg Gly Ala Asp Lys Ala Lys
 1 5 10 15
 Pro Pro Pro Gly Gly Gly Glu Gln Glu Pro Pro Pro Pro Pro Ala Pro
 20 25 30
 Gln Asp Val Glu Met Lys Glu Glu Ala Ala Thr Gly Gly Gly Ser Thr
 35 40 45
 Gly Glu Ala Asp Gly Lys Thr Ala Ala Ala Val Glu His Ser Gln
 50 55 60
 Arg Glu Leu Asp Thr Val Thr Leu Glu Asp Ile Lys Glu His Val Lys
 65 70 75 80
 Gln Leu Glu Lys Ala Val Ser Gly Lys Glu Pro Arg Phe Val Leu Arg
 85 90 95
 Ala Leu Arg Met Leu Pro Ser Thr Ser Arg Arg Leu Asn His Tyr Val
 100 105 110
 Leu Tyr Lys Ala Val Gln Gly Phe Phe Thr Ser Asn Asn Ala Thr Arg
 115 120 125
 Asp Phe Leu Leu Pro Phe Leu Glu Glu Pro Met Asp Thr Glu Ala Asp
 130 135 140
 Leu Gln Phe Arg Pro Arg Thr Gly Lys Ala Ala Ser Thr Pro Leu Leu
 145 150 155 160
 Pro Glu Val Glu Ala Tyr Leu Gln Leu Leu Val Val Ile Phe Met Met
 165 170 175
 Asn Ser Lys Arg Tyr Lys Glu Ala Gln Lys Ile Ser Asp Asp Leu Met
 180 185 190
 Gln Lys Ile Ser Thr Gln Asn Arg Arg Ala Leu Asp Leu Val Ala Ala
 195 200 205

Lys Cys Tyr Tyr Tyr His Ala Arg Val Tyr Glu Phe Leu Asp Lys Leu
 210 215 220
 5 Asp Val Val Arg Ser Phe Leu His Ala Arg Leu Arg Thr Ala Thr Leu
 225 230 235 240
 Arg His Asp Ala Asp Gly Gln Ala Thr Leu Leu Asn Leu Leu Leu Arg
 245 250 255
 Asn Tyr Leu His Tyr Ser Leu Tyr Asp Gln Ala Glu Lys Leu Val Ser
 260 265 270
 10 Lys Ser Val Phe Pro Glu Gln Ala Asn Asn Asn Glu Trp Ala Arg Tyr
 275 280 285
 Leu Tyr Tyr Thr Gly Arg Ile Lys Ala Ile Gln Leu Glu Tyr Ser Glu
 290 295 300
 Ala Arg Arg Thr Met Thr Asn Ala Leu Arg Lys Ala Pro Gln His Thr
 305 310 315 320
 15 Ala Val Gly Phe Lys Gln Thr Val His Lys Leu Leu Ile Val Val Glu
 325 330 335
 Leu Leu Leu Gly Glu Ile Pro Asp Arg Leu Gln Phe Arg Gln Pro Ser
 340 345 350
 Leu Lys Arg Ser Leu Met Pro Tyr Phe Leu Leu Thr Gln Ala Val Arg
 355 360 365
 20 Thr Gly Asn Leu Ala Lys Phe Asn Gln Val Leu Asp Gln Phe Gly Glu
 370 375 380
 Lys Phe Gln Ala Asp Gly Thr Tyr Thr Leu Ile Ile Arg Leu Arg His
 385 390 395 400
 25 Asn Val Ile Lys Thr Gly Val Arg Met Ile Ser Leu Ser Tyr Ser Arg
 405 410 415
 Ile Ser Leu Ala Asp Ile Ala Gln Lys Leu Gln Leu Asp Ser Pro Glu
 420 425 430
 Asp Ala Glu Phe Ile Val Ala Lys Ala Ile Arg Asp Gly Val Ile Glu
 435 440 445
 30 Ala Ser Ile Asn His Glu Lys Gly Tyr Val Gln Ser Lys Glu Met Ile
 450 455 460
 Asp Ile Tyr Ser Thr Arg Glu Pro Gln Leu Ala Phe His Gln Arg Ile
 465 470 475 480
 Ser Phe Cys Leu Asp Ile His Asn Met Ser Val Lys Ala Met Arg Phe
 485 490 495
 35 Pro Pro Lys Ser Tyr Asn Lys Asp Leu Glu Ser Ala Glu Glu Arg Arg
 500 505 510
 Glu Arg Glu Gln Gln Asp Leu Glu Phe Ala Lys Glu Met Ala Glu Asp
 515 520 525
 40 Asp Asp Asp Ser Phe Pro
 530
 <210> 39
 <211> 207
 45 <212> PRT
 <213> Homo sapiens
 50
 <400> 39
 Met Ala Gly Pro Ala Thr Gln Ser Pro Met Lys Leu Met Ala Leu Gln
 1 5 10 15
 Leu Leu Leu Trp His Ser Ala Leu Trp Thr Val Gln Glu Ala Thr Pro
 20 25 30
 55

Leu Gly Pro Ala Ser Ser Leu Pro Gln Ser Phe Leu Leu Lys Cys Leu
 35 40 45
 Glu Gln Val Arg Lys Ile Gln Gly Asp Gly Ala Ala Leu Gln Glu Lys
 5 50 55 60
 Leu Val Ser Glu Cys Ala Thr Tyr Lys Leu Cys His Pro Glu Glu Leu
 65 70 75 80
 Val Leu Leu Gly His Ser Leu Gly Ile Pro Trp Ala Pro Leu Ser Ser
 85 90 95
 Cys Pro Ser Gln Ala Leu Gln Leu Ala Gly Cys Leu Ser Gln Leu His
 10 100 105 110
 Ser Gly Leu Phe Leu Tyr Gln Gly Leu Leu Gln Ala Leu Glu Gly Ile
 115 120 125
 Ser Pro Glu Leu Gly Pro Thr Leu Asp Thr Leu Gln Leu Asp Val Ala
 130 135 140
 Asp Phe Ala Thr Thr Ile Trp Gln Gln Met Glu Glu Leu Gly Met Ala
 15 145 150 155 160
 Pro Ala Leu Gln Pro Thr Gln Gly Ala Met Pro Ala Phe Ala Ser Ala
 165 170 175
 Phe Gln Arg Arg Ala Gly Gly Val Leu Val Ala Ser His Leu Gln Ser
 180 185 190
 Phe Leu Glu Val Ser Tyr Arg Val Leu Arg His Leu Ala Gln Pro
 195 200 205

<210> 40

<211> 989

<212> PRT

<213> Homo sapiens

<400> 40

Met Lys Val Val Asn Leu Lys Gln Ala Ile Leu Gln Ala Trp Lys Glu
 1 5 10 15
 Arg Trp Ser Tyr Tyr Gln Trp Ala Ile Asn Met Lys Lys Phe Phe Pro
 20 25 30
 Lys Gly Ala Thr Trp Asp Ile Leu Asn Leu Ala Asp Ala Leu Leu Glu
 35 40 45
 Gln Ala Met Ile Gly Pro Ser Pro Asn Pro Leu Ile Leu Ser Tyr Leu
 50 55 60
 Lys Tyr Ala Ile Ser Ser Gln Met Val Ser Tyr Ser Ser Val Leu Thr
 65 70 75 80
 Ala Ile Ser Lys Phe Asp Asp Phe Ser Arg Asp Leu Cys Val Gln Ala
 85 90 95
 Leu Leu Asp Ile Met Asp Met Phe Cys Asp Arg Leu Ser Cys His Gly
 100 105 110
 Lys Ala Glu Glu Cys Ile Gly Leu Cys Arg Ala Leu Leu Ser Ala Leu
 115 120 125
 His Trp Leu Leu Arg Cys Thr Ala Ala Ser Ala Glu Arg Leu Arg Glu
 130 135 140
 Gly Leu Glu Ala Gly Thr Pro Ala Ala Gly Glu Lys Gln Leu Ala Met
 145 150 155 160
 Cys Leu Gln Arg Leu Glu Lys Thr Leu Ser Ser Thr Lys Asn Arg Ala
 165 170 175
 Leu Leu His Ile Ala Lys Leu Glu Glu Ala Ser Ser Trp Thr Ala Ile
 180 185 190

	Glu	His	Ser	Leu	Leu	Lys	Leu	Gly	Glu	Ile	Leu	Thr	Asn	Leu	Ser	Asn
			195					200					205			
5	Pro	Gln	Leu	Arg	Ser	Gln	Ala	Glu	Gln	Cys	Gly	Thr	Leu	Ile	Arg	Ser
		210					215					220				
	Ile	Pro	Thr	Met	Leu	Ser	Val	His	Ala	Glu	Gln	Met	His	Lys	Thr	Gly
	225					230					235					240
	Phe	Pro	Thr	Val	His	Ala	Val	Ile	Leu	Leu	Glu	Gly	Thr	Met	Asn	Leu
				245						250					255	
10	Thr	Gly	Glu	Thr	Gln	Ser	Leu	Val	Glu	Gln	Leu	Thr	Met	Val	Lys	Arg
			260						265					270		
	Met	Gln	His	Ile	Pro	Thr	Pro	Leu	Phe	Val	Leu	Glu	Ile	Trp	Lys	Ala
			275					280					285			
	Cys	Phe	Val	Gly	Leu	Ile	Glu	Ser	Pro	Glu	Gly	Thr	Glu	Glu	Leu	Lys
	290						295					300				
15	Trp	Thr	Ala	Phe	Thr	Phe	Leu	Lys	Ile	Pro	Gln	Val	Leu	Val	Lys	Leu
	305					310					315					320
	Lys	Lys	Tyr	Ser	His	Gly	Asp	Lys	Asp	Phe	Thr	Glu	Asp	Val	Asn	Cys
				325						330					335	
	Ala	Phe	Glu	Phe	Leu	Leu	Lys	Leu	Thr	Pro	Leu	Leu	Asp	Lys	Ala	Asp
				340					345					350		
20	Gln	Arg	Cys	Asn	Cys	Asp	Cys	Thr	Asn	Phe	Leu	Leu	Gln	Glu	Cys	Gly
			355					360					365			
	Lys	Gln	Gly	Leu	Leu	Ser	Glu	Ala	Ser	Val	Asn	Asn	Leu	Met	Ala	Lys
		370					375					380				
25	Arg	Lys	Ala	Asp	Arg	Glu	His	Ala	Pro	Gln	Gln	Lys	Ser	Gly	Glu	Asn
	385					390					395					400
	Ala	Asn	Ile	Gln	Pro	Asn	Ile	Gln	Leu	Ile	Leu	Arg	Ala	Glu	Pro	Thr
				405						410					415	
	Val	Thr	Asn	Ile	Leu	Lys	Thr	Met	Asp	Ala	Asp	His	Ser	Lys	Ser	Pro
				420					425					430		
30	Glu	Gly	Leu	Leu	Gly	Val	Leu	Gly	His	Met	Leu	Ser	Gly	Lys	Ser	Leu
		435						440					445			
	Asp	Leu	Leu	Leu	Ala	Ala	Ala	Ala	Thr	Gly	Lys	Leu	Lys	Ser	Ser	Phe
		450					455				460					
	Ala	Arg	Lys	Phe	Ile	Asn	Leu	Asn	Glu	Phe	Thr	Thr	Tyr	Gly	Ser	Glu
	465					470					475					480
35	Glu	Ser	Thr	Lys	Pro	Ala	Ser	Val	Arg	Ala	Leu	Leu	Phe	Asp	Ile	Ser
				485					490						495	
	Phe	Leu	Met	Leu	Cys	His	Val	Ala	Gln	Thr	Tyr	Gly	Ser	Glu	Val	Ile
				500					505					510		
	Leu	Ser	Glu	Ser	Arg	Thr	Gly	Ala	Glu	Val	Pro	Phe	Phe	Glu	Thr	Trp
			515					520					525			
40	Met	Gln	Thr	Cys	Met	Pro	Glu	Glu	Gly	Lys	Ile	Leu	Asn	Pro	Asp	His
		530					535					540				
	Pro	Cys	Phe	Arg	Pro	Asp	Ser	Thr	Lys	Val	Glu	Ser	Leu	Val	Ala	Leu
	545					550					555					560
45	Leu	Asn	Asn	Ser	Ser	Glu	Met	Lys	Leu	Val	Gln	Met	Lys	Trp	His	Glu
				565						570					575	
	Ala	Cys	Leu	Ser	Ile	Ser	Ala	Ala	Ile	Leu	Glu	Ile	Leu	Asn	Ala	Trp
				580					585					590		
	Glu	Asn	Gly	Val	Leu	Ala	Phe	Glu	Ser	Ile	Gln	Lys	Ile	Thr	Asp	Asn
			595					600					605			
50	Ile	Lys	Gly	Lys	Val	Cys	Ser	Leu	Ala	Val	Cys	Ala	Val	Ala	Trp	Leu
		610					615					620				
	Val	Ala	His	Val	Arg	Met	Leu	Gly	Leu	Asp	Glu	Arg	Glu	Lys	Ser	Leu
	625					630					635					640
	Gln	Met	Ile	Arg	Gln	Leu	Ala	Gly	Pro	Leu	Phe	Ser	Glu	Asn	Thr	Leu
					645					650					655	

	Gln	Phe	Tyr	Asn	Glu	Arg	Val	Val	Ile	Met	Asn	Ser	Ile	Leu	Glu	Arg
				660					665					670		
5	Met	Cys	Ala	Asp	Val	Leu	Gln	Gln	Thr	Ala	Thr	Gln	Ile	Lys	Phe	Pro
			675					680					685			
	Ser	Thr	Gly	Val	Asp	Thr	Met	Pro	Tyr	Trp	Asn	Leu	Leu	Pro	Pro	Lys
			690				695					700				
	Arg	Pro	Ile	Lys	Glu	Val	Leu	Thr	Asp	Ile	Phe	Ala	Lys	Val	Leu	Glu
						710					715					720
10	Lys	Gly	Trp	Val	Asp	Ser	Arg	Ser	Ile	His	Ile	Phe	Asp	Thr	Leu	Leu
					725					730						735
	His	Met	Gly	Gly	Val	Tyr	Trp	Phe	Cys	Asn	Asn	Leu	Ile	Lys	Glu	Leu
				740					745					750		
	Leu	Lys	Glu	Thr	Arg	Lys	Glu	His	Thr	Leu	Arg	Ala	Val	Glu	Leu	Leu
				755				760					765			
15	Tyr	Ser	Ile	Phe	Cys	Leu	Asp	Met	Gln	Gln	Val	Thr	Leu	Val	Leu	Leu
		770					775						780			
	Gly	His	Ile	Leu	Pro	Gly	Leu	Leu	Thr	Asp	Ser	Ser	Lys	Trp	His	Ser
		785				790					795					800
	Leu	Met	Asp	Pro	Pro	Gly	Thr	Ala	Leu	Ala	Lys	Leu	Ala	Val	Trp	Cys
					805					810					815	
20	Ala	Leu	Ser	Ser	Tyr	Ser	Ser	His	Lys	Gly	Gln	Ala	Ser	Thr	Arg	Gln
				820					825					830		
	Lys	Lys	Arg	His	Arg	Glu	Asp	Ile	Glu	Asp	Tyr	Ile	Ser	Leu	Phe	Pro
				835				840					845			
	Leu	Asp	Asp	Val	Gln	Pro	Ser	Lys	Leu	Met	Arg	Leu	Leu	Ser	Ser	Asn
							855					860				
25	Glu	Asp	Asp	Ala	Asn	Ile	Leu	Ser	Ser	Pro	Thr	Asp	Arg	Ser	Met	Ser
						870					875					880
	Ser	Ser	Leu	Ser	Ala	Ser	Gln	Leu	His	Thr	Val	Asn	Met	Arg	Asp	Pro
					885					890					895	
30	Leu	Asn	Arg	Val	Leu	Ala	Asn	Leu	Phe	Leu	Leu	Ile	Ser	Ser	Ile	Leu
				900					905					910		
	Gly	Ser	Arg	Thr	Ala	Gly	Pro	His	Thr	Gln	Phe	Val	Gln	Trp	Phe	Met
								920					925			
	Glu	Glu	Cys	Val	Asp	Cys	Leu	Glu	Gln	Gly	Gly	Arg	Gly	Ser	Val	Leu
							935					940				
35	Gln	Phe	Met	Pro	Phe	Thr	Thr	Val	Ser	Glu	Leu	Val	Lys	Val	Ser	Ala
						950					955					960
	Met	Ser	Ser	Pro	Lys	Val	Val	Leu	Ala	Ile	Thr	Asp	Leu	Ser	Leu	Pro
					965					970					975	
	Leu	Gly	Arg	Gln	Val	Ala	Ala	Lys	Ala	Ile	Ala	Ala	Leu			
40				980						985						

<210> 41

<211> 490

<212> PRT

<213> Homo sapiens

<400> 41

Met Glu Gln Lys Pro Ser Lys Val Glu Cys Gly Ser Asp Pro Glu Glu
1 5 10 15
Asn Ser Ala Arg Ser Pro Asp Gly Lys Arg Lys Arg Lys Asn Gly Gln
20 25 30

Cys Ser Leu Lys Thr Ser Met Ser Gly Tyr Ile Pro Ser Tyr Leu Asp
 35 40 45
 Lys Asp Glu Gln Cys Val Val Cys Gly Asp Lys Ala Thr Gly Tyr His
 50 55 60
 Tyr Arg Cys Ile Thr Cys Glu Gly Cys Lys Gly Phe Phe Arg Arg Thr
 65 70 75 80
 Ile Gln Lys Asn Leu His Pro Thr Tyr Ser Cys Lys Tyr Asp Ser Cys
 85 90 95
 Cys Val Ile Asp Lys Ile Thr Arg Asn Gln Cys Gln Leu Cys Arg Phe
 100 105 110
 Lys Lys Cys Ile Ala Val Gly Met Ala Met Asp Leu Val Leu Asp Asp
 115 120 125
 Ser Lys Arg Val Ala Lys Arg Lys Leu Ile Glu Gln Asn Arg Glu Arg
 130 135 140
 Arg Arg Lys Glu Glu Met Ile Arg Ser Leu Gln Gln Arg Pro Glu Pro
 145 150 155 160
 Thr Pro Glu Glu Trp Asp Leu Ile His Ile Ala Thr Glu Ala His Arg
 165 170 175
 Ser Thr Asn Ala Gln Gly Ser His Trp Lys Gln Arg Arg Lys Phe Leu
 180 185 190
 Pro Asp Asp Ile Gly Gln Ser Pro Ile Val Ser Met Pro Asp Gly Asp
 195 200 205
 Lys Val Asp Leu Glu Ala Phe Ser Glu Phe Thr Lys Ile Ile Thr Pro
 210 215 220
 Ala Ile Thr Arg Val Val Asp Phe Ala Lys Lys Leu Pro Met Phe Ser
 225 230 235 240
 Glu Leu Pro Cys Glu Asp Gln Ile Ile Leu Leu Lys Gly Cys Cys Met
 245 250 255
 Glu Ile Met Ser Leu Arg Ala Ala Val Arg Tyr Asp Pro Glu Ser Asp
 260 265 270
 Thr Leu Thr Leu Ser Gly Glu Met Ala Val Lys Arg Glu Gln Leu Lys
 275 280 285
 Asn Gly Gly Leu Gly Val Val Ser Asp Ala Ile Phe Glu Leu Gly Lys
 290 295 300
 Ser Leu Ser Ala Phe Asn Leu Asp Asp Thr Glu Val Ala Leu Leu Gln
 305 310 315 320
 Ala Val Leu Leu Met Ser Thr Asp Arg Ser Gly Leu Leu Cys Val Asp
 325 330 335
 Lys Ile Glu Lys Ser Gln Glu Ala Tyr Leu Leu Ala Phe Glu His Tyr
 340 345 350
 Val Asn His Arg Lys His Asn Ile Pro His Phe Trp Pro Lys Leu Leu
 355 360 365
 Met Lys Glu Arg Glu Val Gln Ser Ser Ile Leu Tyr Lys Gly Ala Ala
 370 375 380
 Ala Glu Gly Arg Pro Gly Gly Ser Leu Gly Val His Pro Glu Gly Gln
 385 390 395 400
 Gln Leu Leu Gly Met His Val Val Gln Gly Pro Gln Val Arg Gln Leu
 405 410 415
 Glu Gln Gln Leu Gly Glu Ala Gly Ser Leu Gln Gly Pro Val Leu Gln
 420 425 430
 His Gln Ser Pro Lys Ser Pro Gln Gln Arg Leu Leu Glu Leu Leu His
 435 440 445
 Arg Ser Gly Ile Leu His Ala Arg Ala Val Cys Gly Glu Asp Asp Ser
 450 455 460
 Ser Glu Ala Asp Ser Pro Ser Ser Ser Glu Glu Glu Pro Glu Val Cys
 465 470 475 480
 Glu Asp Leu Ala Gly Asn Ala Ala Ser Pro
 485 490

<210> 42

<211> 614

<212> PRT

<213> Homo sapiens

<400> 42

Met Thr Thr Leu Asp Ser Asn Asn Asn Thr Gly Gly Val Ile Thr Tyr
 1 5 10 15
 Ile Gly Ser Ser Gly Ser Ser Pro Ser Arg Thr Ser Pro Glu Ser Leu
 20 25 30
 Tyr Ser Asp Asn Ser Asn Gly Ser Phe Gln Ser Leu Thr Gln Gly Cys
 35 40 45
 Pro Thr Tyr Phe Pro Pro Ser Pro Thr Gly Ser Leu Thr Gln Asp Pro
 50 55 60
 Ala Arg Ser Phe Gly Ser Ile Pro Pro Ser Leu Ser Asp Asp Gly Ser
 65 70 75 80
 Pro Ser Ser Ser Ser Ser Ser Ser Ser Ser Ser Ser Ser Phe Tyr Asn
 85 90 95
 Gly Ser Pro Pro Gly Ser Leu Gln Val Ala Met Glu Asp Ser Ser Arg
 100 105 110
 Val Ser Pro Ser Lys Ser Thr Ser Asn Ile Thr Lys Leu Asn Gly Met
 115 120 125
 Val Leu Leu Cys Lys Val Cys Gly Asp Val Ala Ser Gly Phe His Tyr
 130 135 140
 Gly Val Leu Ala Cys Glu Gly Cys Lys Gly Phe Phe Arg Arg Ser Ile
 145 150 155 160
 Gln Gln Asn Ile Gln Tyr Lys Arg Cys Leu Lys Asn Glu Asn Cys Ser
 165 170 175
 Ile Val Arg Ile Asn Arg Asn Arg Cys Gln Gln Cys Arg Phe Lys Lys
 180 185 190
 Cys Leu Ser Val Gly Met Ser Arg Asp Ala Val Arg Phe Gly Arg Ile
 195 200 205
 Pro Lys Arg Glu Lys Gln Arg Met Leu Ala Glu Met Gln Ser Ala Met
 210 215 220
 Asn Leu Ala Asn Asn Gln Leu Ser Ser Gln Cys Pro Leu Glu Thr Ser
 225 230 235 240
 Pro Thr Gln His Pro Thr Pro Gly Pro Met Gly Pro Ser Pro Pro Pro
 245 250 255
 Ala Pro Val Pro Ser Pro Leu Val Gly Phe Ser Gln Phe Pro Gln Gln
 260 265 270
 Leu Thr Pro Pro Arg Ser Pro Ser Pro Glu Pro Thr Val Glu Asp Val
 275 280 285
 Ile Ser Gln Val Ala Arg Ala His Arg Glu Ile Phe Thr Tyr Ala His
 290 295 300
 Asp Lys Leu Gly Ser Ser Pro Gly Asn Phe Asn Ala Asn His Ala Ser
 305 310 315 320
 Gly Ser Pro Pro Ala Thr Thr Pro His Arg Trp Glu Asn Gln Gly Cys
 325 330 335
 Pro Pro Ala Pro Asn Asp Asn Asn Thr Leu Ala Ala Gln Arg His Asn
 340 345 350
 Glu Ala Leu Asn Gly Leu Arg Gln Ala Pro Ser Ser Tyr Pro Pro Thr
 355 360 365

Trp Pro Pro Gly Pro Ala His His Ser Cys His Gln Ser Asn Ser Asn
 370 375 380
 Gly His Arg Leu Cys Pro Thr His Val Tyr Ala Ala Pro Glu Gly Lys
 385 390 395 400
 Ala Pro Ala Asn Ser Pro Arg Gln Gly Asn Ser Lys Asn Val Leu Leu
 405 410 415
 Ala Cys Pro Met Asn Met Tyr Pro His Gly Arg Ser Gly Arg Thr Val
 420 425 430
 Gln Glu Ile Trp Glu Asp Phe Ser Met Ser Phe Thr Pro Ala Val Arg
 435 440 445
 Glu Val Val Glu Phe Ala Lys His Ile Pro Gly Phe Arg Asp Leu Ser
 450 455 460
 Gln His Asp Gln Val Thr Leu Leu Lys Ala Gly Thr Phe Glu Val Leu
 465 470 475 480
 Met Val Arg Phe Ala Ser Leu Phe Asn Val Lys Asp Gln Thr Val Met
 485 490 495
 Phe Leu Ser Arg Thr Thr Tyr Ser Leu Gln Glu Leu Gly Ala Met Gly
 500 505 510
 Met Gly Asp Leu Leu Ser Ala Met Phe Asp Phe Ser Glu Lys Leu Asn
 515 520 525
 Ser Leu Ala Leu Thr Glu Glu Leu Gly Leu Phe Thr Ala Val Val
 530 535 540
 Leu Val Ser Ala Asp Arg Ser Gly Met Glu Asn Ser Ala Ser Val Glu
 545 550 555 560
 Gln Leu Gln Glu Thr Leu Leu Arg Ala Leu Arg Ala Leu Val Leu Lys
 565 570 575
 Asn Arg Pro Leu Glu Thr Ser Arg Phe Thr Lys Leu Leu Leu Lys Leu
 580 585 590
 Pro Asp Leu Arg Thr Leu Asn Asn Met His Ser Glu Lys Leu Leu Ser
 595 600 605
 Phe Arg Val Asp Ala Gln
 610

 <210> 43

 <211> 703

 <212> PRT

 <213> Homo sapiens

 <400> 43
 Met Ala Asp Arg Arg Arg Gln Arg Ala Ser Gln Asp Thr Glu Asp Glu
 1 5 10 15
 Glu Ser Gly Ala Ser Gly Ser Asp Ser Gly Gly Ser Pro Leu Arg Gly
 20 25 30
 Gly Gly Ser Cys Ser Gly Ser Ala Gly Gly Gly Gly Ser Gly Ser Leu
 35 40 45
 Pro Ser Gln Arg Gly Gly Arg Thr Gly Ala Leu His Leu Arg Arg Val
 50 55 60
 Glu Ser Gly Gly Ala Lys Ser Ala Glu Glu Ser Glu Cys Glu Ser Glu
 65 70 75 80
 Asp Gly Ile Glu Gly Asp Ala Val Leu Ser Asp Tyr Glu Ser Ala Glu
 85 90 95
 Asp Ser Glu Gly Glu Glu Gly Glu Tyr Ser Glu Glu Glu Asn Ser Lys
 100 105 110

	Val	Glu	Leu	Lys	Ser	Glu	Ala	Asn	Asp	Ala	Val	Asn	Ser	Ser	Thr	Lys
			115					120					125			
5	Glu	Glu	Lys	Gly	Glu	Glu	Lys	Pro	Asp	Thr	Lys	Ser	Thr	Val	Thr	Gly
		130					135					140				
	Glu	Arg	Gln	Ser	Gly	Asp	Gly	Gln	Glu	Ser	Thr	Glu	Pro	Val	Glu	Asn
		145				150					155					160
	Lys	Val	Gly	Lys	Lys	Gly	Pro	Lys	His	Leu	Asp	Asp	Asp	Glu	Asp	Arg
					165					170						175
10	Lys	Asn	Pro	Ala	Tyr	Ile	Pro	Arg	Lys	Gly	Leu	Phe	Phe	Glu	His	Asp
				180					185					190		
	Leu	Arg	Gly	Gln	Thr	Gln	Glu	Glu	Glu	Val	Arg	Pro	Lys	Gly	Arg	Gln
			195					200					205			
	Arg	Lys	Leu	Trp	Lys	Asp	Glu	Gly	Arg	Trp	Glu	His	Asp	Lys	Phe	Arg
		210					215					220				
15	Glu	Asp	Glu	Gln	Ala	Pro	Lys	Ser	Arg	Gln	Glu	Leu	Ile	Ala	Leu	Tyr
		225					230				235					240
	Gly	Tyr	Asp	Ile	Arg	Ser	Ala	His	Asn	Pro	Asp	Asp	Ile	Lys	Pro	Arg
					245					250					255	
	Arg	Ile	Arg	Lys	Pro	Arg	Tyr	Gly	Ser	Pro	Pro	Gln	Arg	Asp	Pro	Asn
				260					265					270		
20	Trp	Asn	Gly	Glu	Arg	Leu	Asn	Lys	Ser	His	Arg	His	Gln	Gly	Leu	Gly
			275					280					285			
	Gly	Thr	Leu	Pro	Pro	Arg	Thr	Phe	Ile	Asn	Arg	Asn	Ala	Ala	Gly	Thr
		290					295					300				
25	Gly	Arg	Met	Ser	Ala	Pro	Arg	Asn	Tyr	Ser	Arg	Ser	Gly	Gly	Phe	Lys
		305					310				315					320
	Glu	Gly	Arg	Ala	Gly	Phe	Arg	Pro	Val	Glu	Ala	Gly	Gly	Gln	His	Gly
					325					330					335	
	Gly	Arg	Ser	Gly	Glu	Thr	Val	Lys	His	Glu	Ile	Ser	Tyr	Arg	Ser	Arg
				340					345					350		
30	Arg	Leu	Glu	Gln	Thr	Ser	Val	Arg	Asp	Pro	Ser	Pro	Glu	Ala	Asp	Ala
			355					360					365			
	Pro	Val	Leu	Gly	Ser	Pro	Glu	Lys	Glu	Glu	Ala	Ala	Ser	Glu	Pro	Pro
		370					375					380				
	Ala	Ala	Ala	Pro	Asp	Ala	Ala	Pro	Pro	Pro	Pro	Asp	Arg	Pro	Ile	Glu
		385					390				395					400
35	Lys	Lys	Ser	Tyr	Ser	Arg	Ala	Arg	Arg	Thr	Arg	Thr	Lys	Val	Gly	Asp
					405					410					415	
	Ala	Val	Lys	Leu	Ala	Glu	Glu	Val	Pro	Pro	Pro	Pro	Glu	Gly	Leu	Ile
				420					425				430			
	Pro	Ala	Pro	Pro	Val	Pro	Glu	Thr	Thr	Pro	Thr	Pro	Pro	Thr	Lys	Thr
				435				440					445			
40	Gly	Thr	Trp	Glu	Ala	Pro	Val	Asp	Ser	Ser	Thr	Ser	Gly	Leu	Glu	Gln
		450					455					460				
	Asp	Val	Ala	Gln	Leu	Asn	Ile	Ala	Glu	Gln	Asn	Trp	Ser	Pro	Gly	Gln
		465				470					475					480
45	Pro	Ser	Phe	Leu	Gln	Pro	Arg	Glu	Leu	Arg	Gly	Met	Pro	Asn	His	Ile
					485				490						495	
	His	Met	Gly	Ala	Gly	Pro	Pro	Pro	Gln	Phe	Asn	Arg	Met	Glu	Glu	Met
				500					505					510		
	Gly	Val	Gln	Gly	Gly	Arg	Ala	Lys	Arg	Tyr	Ser	Ser	Gln	Arg	Gln	Arg
			515					520					525			
50	Pro	Val	Pro	Glu	Pro	Pro	Ala	Pro	Pro	Val	His	Ile	Ser	Ile	Met	Glu
		530					535					540				
	Gly	His	Tyr	Tyr	Asp	Pro	Leu	Gln	Phe	Gln	Gly	Pro	Ile	Tyr	Thr	His
		545				550					555					560
	Gly	Asp	Ser	Pro	Ala	Pro	Leu	Pro	Pro	Gln	Gly	Met	Leu	Val	Gln	Pro
					565					570					575	

Gly Met Asn Leu Pro His Pro Gly Leu His Pro His Gln Thr Pro Ala
 580 585 590
 Pro Leu Pro Asn Pro Gly Leu Tyr Pro Pro Pro Val Ser Met Ser Pro
 595 600 605
 Gly Gln Pro Pro Pro Gln Gln Leu Leu Ala Pro Thr Tyr Phe Ser Ala
 610 615 620
 Pro Gly Val Met Asn Phe Gly Asn Pro Ser Tyr Pro Tyr Ala Pro Gly
 625 630 635 640
 Ala Leu Pro Pro Pro Pro Pro Pro His Leu Tyr Pro Asn Thr Gln Ala
 645 650 655
 Pro Ser Gln Val Tyr Gly Gly Val Thr Tyr Tyr Asn Pro Ala Gln Gln
 660 665 670
 Gln Val Gln Pro Lys Pro Ser Pro Pro Arg Arg Thr Pro Gln Pro Val
 675 680 685
 Thr Ile Lys Pro Pro Pro Pro Glu Val Val Ser Arg Gly Ser Ser
 690 695 700

<210> 44

<211> 560

<212> PRT

<213> Homo sapiens

<400> 44

Met Pro Gln Thr Arg Ser Gln Ala Gln Ala Thr Ile Ser Phe Pro Lys
 1 5 10 15
 Arg Lys Leu Ser Arg Ala Leu Asn Lys Ala Lys Asn Ser Ser Asp Ala
 20 25 30
 Lys Leu Glu Pro Thr Asn Val Gln Thr Val Thr Cys Ser Pro Arg Val
 35 40 45
 Lys Ala Leu Pro Leu Ser Pro Arg Lys Arg Leu Gly Asp Asp Asn Leu
 50 55 60
 Cys Asn Thr Pro His Leu Pro Pro Cys Ser Pro Pro Lys Gln Gly Lys
 65 70 75 80
 Lys Glu Asn Gly Pro Pro His Ser His Thr Leu Lys Gly Arg Arg Leu
 85 90 95
 Val Phe Asp Asn Gln Leu Thr Ile Lys Ser Pro Ser Lys Arg Glu Leu
 100 105 110
 Ala Lys Val His Gln Asn Lys Ile Leu Ser Ser Val Arg Lys Ser Gln
 115 120 125
 Glu Ile Thr Thr Asn Ser Glu Gln Arg Cys Pro Leu Lys Lys Glu Ser
 130 135 140
 Ala Cys Val Arg Leu Phe Lys Gln Glu Gly Thr Cys Tyr Gln Gln Ala
 145 150 155 160
 Lys Leu Val Leu Asn Thr Ala Val Pro Asp Arg Leu Pro Ala Arg Glu
 165 170 175
 Arg Glu Met Asp Val Ile Arg Asn Phe Leu Arg Glu His Ile Cys Gly
 180 185 190
 Lys Lys Ala Gly Ser Leu Tyr Leu Ser Gly Ala Pro Gly Thr Gly Lys
 195 200 205
 Thr Ala Cys Leu Ser Arg Ile Leu Gln Asp Leu Lys Lys Glu Leu Lys
 210 215 220
 Gly Phe Lys Thr Ile Met Leu Asn Cys Met Ser Leu Arg Thr Ala Gln
 225 230 235 240

Ala Val Phe Pro Ala Ile Ala Gln Glu Ile Cys Gln Glu Glu Val Ser
 245 250 255
 Arg Pro Ala Gly Lys Asp Met Met Arg Lys Leu Glu Lys His Met Thr
 260 265 270
 Ala Glu Lys Gly Pro Met Ile Val Leu Val Leu Asp Glu Met Asp Gln
 275 280 285
 Leu Asp Ser Lys Gly Gln Asp Val Leu Tyr Thr Leu Phe Glu Trp Pro
 290 295 300
 Trp Leu Ser Asn Ser His Leu Val Leu Ile Gly Ile Ala Asn Thr Leu
 305 310 315 320
 Asp Leu Thr Asp Arg Ile Leu Pro Arg Leu Gln Ala Arg Glu Lys Cys
 325 330 335
 Lys Pro Gln Leu Leu Asn Phe Pro Pro Tyr Thr Arg Asn Gln Ile Val
 340 345 350
 Thr Ile Leu Gln Asp Arg Leu Asn Gln Val Ser Arg Asp Gln Val Leu
 355 360 365
 Asp Asn Ala Ala Val Gln Phe Cys Ala Arg Lys Val Ser Ala Val Ser
 370 375 380
 Gly Asp Val Arg Lys Ala Leu Asp Val Cys Arg Arg Ala Ile Glu Ile
 385 390 395 400
 Val Glu Ser Asp Val Lys Ser Gln Thr Ile Leu Lys Pro Leu Ser Glu
 405 410 415
 Cys Lys Ser Pro Ser Glu Pro Leu Ile Pro Lys Arg Val Gly Leu Ile
 420 425 430
 His Ile Ser Gln Val Ile Ser Glu Val Asp Gly Asn Arg Met Thr Leu
 435 440 445
 Ser Gln Glu Gly Ala Gln Asp Ser Phe Pro Leu Gln Gln Lys Ile Leu
 450 455 460
 Val Cys Ser Leu Met Leu Leu Ile Arg Gln Leu Lys Ile Lys Glu Val
 465 470 475 480
 Thr Leu Gly Lys Leu Tyr Glu Ala Tyr Ser Lys Val Cys Arg Lys Gln
 485 490 495
 Gln Val Ala Ala Val Asp Gln Ser Glu Cys Leu Ser Leu Ser Gly Leu
 500 505 510
 Leu Glu Ala Arg Gly Ile Leu Gly Leu Lys Arg Asn Lys Glu Thr Arg
 515 520 525
 Leu Thr Lys Val Phe Phe Lys Ile Glu Glu Lys Glu Ile Glu His Ala
 530 535 540
 Leu Lys Asp Lys Ala Leu Ile Gly Asn Ile Leu Ala Thr Gly Leu Pro
 545 550 555 560

 <210> 45
 <211> 462
 <212> PRT
 <213> Homo sapiens

 <400> 45
 Met Ala Ser Asn Ser Ser Ser Cys Pro Thr Pro Gly Gly Gly His Leu
 1 5 10 15
 Asn Gly Tyr Pro Val Pro Pro Tyr Ala Phe Phe Phe Pro Pro Met Leu
 20 25 30
 Gly Gly Leu Ser Pro Pro Gly Ala Leu Thr Thr Leu Gln His Gln Leu
 35 40 45

	Pro	Val	Ser	Gly	Tyr	Ser	Thr	Pro	Ser	Pro	Ala	Thr	Ile	Glu	Thr	Gln
	50						55					60				
5	Ser	Ser	Ser	Ser	Glu	Glu	Ile	Val	Pro	Ser	Pro	Pro	Ser	Pro	Pro	Pro
	65				70						75					80
	Leu	Pro	Arg	Ile	Tyr	Lys	Pro	Cys	Phe	Val	Cys	Gln	Asp	Lys	Ser	Ser
					85					90					95	
	Gly	Tyr	His	Tyr	Gly	Val	Ser	Ala	Cys	Glu	Gly	Cys	Lys	Gly	Phe	Phe
					100				105					110		
10	Arg	Arg	Ser	Ile	Gln	Lys	Asn	Met	Val	Tyr	Thr	Cys	His	Arg	Asp	Lys
					115			120					125			
	Asn	Cys	Ile	Ile	Asn	Lys	Val	Thr	Arg	Asn	Arg	Cys	Gln	Tyr	Cys	Arg
	130						135					140				
	Leu	Gln	Lys	Cys	Phe	Glu	Val	Gly	Met	Ser	Lys	Glu	Ser	Val	Arg	Asn
15	145					150					155					160
	Asp	Arg	Asn	Lys	Lys	Lys	Lys	Glu	Val	Pro	Lys	Pro	Glu	Cys	Ser	Glu
					165					170					175	
	Ser	Tyr	Thr	Leu	Thr	Pro	Glu	Val	Gly	Glu	Leu	Ile	Glu	Lys	Val	Arg
					180				185					190		
20	Lys	Ala	His	Gln	Glu	Thr	Phe	Pro	Ala	Leu	Cys	Gln	Leu	Gly	Lys	Tyr
					195			200					205			
	Thr	Thr	Asn	Asn	Ser	Ser	Glu	Gln	Arg	Val	Ser	Leu	Asp	Ile	Asp	Leu
							215					220				
	Trp	Asp	Lys	Phe	Ser	Glu	Leu	Ser	Thr	Lys	Cys	Ile	Ile	Lys	Thr	Val
	225					230					235					240
25	Glu	Phe	Ala	Lys	Gln	Leu	Pro	Gly	Phe	Thr	Thr	Leu	Thr	Ile	Ala	Asp
					245					250					255	
	Gln	Ile	Thr	Leu	Leu	Lys	Ala	Ala	Cys	Leu	Asp	Ile	Leu	Ile	Leu	Arg
					260				265					270		
	Ile	Cys	Thr	Arg	Tyr	Thr	Pro	Glu	Gln	Asp	Thr	Met	Thr	Phe	Ser	Asp
30					275			280					285			
	Gly	Leu	Thr	Leu	Asn	Arg	Thr	Gln	Met	His	Asn	Ala	Gly	Phe	Gly	Pro
					290		295					300				
	Leu	Thr	Asp	Leu	Val	Phe	Ala	Phe	Ala	Asn	Gln	Leu	Leu	Pro	Leu	Glu
	305					310					315					320
35	Met	Asp	Asp	Ala	Glu	Thr	Gly	Leu	Leu	Ser	Ala	Ile	Cys	Leu	Ile	Cys
					325					330					335	
	Gly	Asp	Arg	Gln	Asp	Leu	Glu	Gln	Pro	Asp	Arg	Val	Asp	Met	Leu	Gln
					340				345					350		
	Glu	Pro	Leu	Leu	Glu	Ala	Leu	Lys	Val	Tyr	Val	Arg	Lys	Arg	Arg	Pro
					355			360					365			
40	Ser	Arg	Pro	His	Met	Phe	Pro	Lys	Met	Leu	Met	Lys	Ile	Thr	Asp	Leu
							375					380				
	Arg	Ser	Ile	Ser	Ala	Lys	Gly	Ala	Glu	Arg	Val	Ile	Thr	Leu	Lys	Met
	385					390					395					400
	Glu	Ile	Pro	Gly	Ser	Met	Pro	Pro	Leu	Ile	Gln	Glu	Met	Leu	Glu	Asn
45					405						410				415	
	Ser	Glu	Gly	Leu	Asp	Thr	Leu	Ser	Gly	Gln	Pro	Gly	Gly	Gly	Gly	Arg
					420				425					430		
	Asp	Gly	Gly	Gly	Leu	Ala	Pro	Pro	Pro	Gly	Ser	Cys	Ser	Pro	Ser	Leu
					435			440					445			
50	Ser	Pro	Ser	Ser	Asn	Arg	Ser	Ser	Pro	Ala	Thr	His	Ser	Pro		
	450						455					460				

55

<210> 46

<211> 1531

<212> PRT

<213> Homo sapiens

<400> 46

```

Met Glu Val Ser Pro Leu Gln Pro Val Asn Glu Asn Met Gln Val Asn
1      5      10      15
Lys Ile Lys Lys Asn Glu Asp Ala Lys Lys Arg Leu Ser Val Glu Arg
20      25      30
Ile Tyr Gln Lys Lys Thr Gln Leu Glu His Ile Leu Leu Arg Pro Asp
35      40      45
Thr Tyr Ile Gly Ser Val Glu Leu Val Thr Gln Gln Met Trp Val Tyr
50      55      60
Asp Glu Asp Val Gly Ile Asn Tyr Arg Glu Val Thr Phe Val Pro Gly
65      70      75      80
Leu Tyr Lys Ile Phe Asp Glu Ile Leu Val Asn Ala Ala Asp Asn Lys
85      90      95
Gln Arg Asp Pro Lys Met Ser Cys Ile Arg Val Thr Ile Asp Pro Glu
100     105     110
Asn Asn Leu Ile Ser Ile Trp Asn Asn Gly Lys Gly Ile Pro Val Val
115     120     125
Glu His Lys Val Glu Lys Met Tyr Val Pro Ala Leu Ile Phe Gly Gln
130     135     140
Leu Leu Thr Ser Ser Asn Tyr Asp Asp Asp Glu Lys Lys Val Thr Gly
145     150     155     160
Gly Arg Asn Gly Tyr Gly Ala Lys Leu Cys Asn Ile Phe Ser Thr Lys
165     170     175
Phe Thr Val Glu Thr Ala Ser Arg Glu Tyr Lys Lys Met Phe Lys Gln
180     185     190
Thr Trp Met Asp Asn Met Gly Arg Ala Gly Glu Met Glu Leu Lys Pro
195     200     205
Phe Asn Gly Glu Asp Tyr Thr Cys Ile Thr Phe Gln Pro Asp Leu Ser
210     215     220
Lys Phe Lys Met Gln Ser Leu Asp Lys Asp Ile Val Ala Leu Met Val
225     230     235     240
Arg Arg Ala Tyr Asp Ile Ala Gly Ser Thr Lys Asp Val Lys Val Phe
245     250     255
Leu Asn Gly Asn Lys Leu Pro Val Lys Gly Phe Arg Ser Tyr Val Asp
260     265     270
Met Tyr Leu Lys Asp Lys Leu Asp Glu Thr Gly Asn Ser Leu Lys Val
275     280     285
Ile His Glu Gln Val Asn His Arg Trp Glu Val Cys Leu Thr Met Ser
290     295     300
Glu Lys Gly Phe Gln Gln Ile Ser Phe Val Asn Ser Ile Ala Thr Ser
305     310     315     320
Lys Gly Gly Arg His Val Asp Tyr Val Ala Asp Gln Ile Val Thr Lys
325     330     335
Leu Val Asp Val Val Lys Lys Lys Asn Lys Gly Gly Val Ala Val Lys
340     345     350
Ala His Gln Val Lys Asn His Met Trp Ile Phe Val Asn Ala Leu Ile
355     360     365
Glu Asn Pro Thr Phe Asp Ser Gln Thr Lys Glu Asn Met Thr Leu Gln
370     375     380

```

Pro Lys Ser Phe Gly Ser Thr Cys Gln Leu Ser Glu Lys Phe Ile Lys
 385 390 395 400
 Ala Ala Ile Gly Cys Gly Ile Val Glu Ser Ile Leu Asn Trp Val Lys
 405 410 415
 Phe Lys Ala Gln Val Gln Leu Asn Lys Lys Cys Ser Ala Val Lys His
 420 425 430
 Asn Arg Ile Lys Gly Ile Pro Lys Leu Asp Asp Ala Asn Asp Ala Gly
 435 440 445
 Gly Arg Asn Ser Thr Glu Cys Thr Leu Ile Leu Thr Glu Gly Asp Ser
 450 455 460
 Ala Lys Thr Leu Ala Val Ser Gly Leu Gly Val Val Gly Arg Asp Lys
 465 470 475 480
 Tyr Gly Val Phe Pro Leu Arg Gly Lys Ile Leu Asn Val Arg Glu Ala
 485 490 495
 Ser His Lys Gln Ile Met Glu Asn Ala Glu Ile Asn Asn Ile Ile Lys
 500 505 510
 Ile Val Gly Leu Gln Tyr Lys Lys Asn Tyr Glu Asp Glu Asp Ser Leu
 515 520 525
 Lys Thr Leu Arg Tyr Gly Lys Ile Met Ile Met Thr Asp Gln Asp Gln
 530 535 540
 Asp Gly Ser His Ile Lys Gly Leu Leu Ile Asn Phe Ile His His Asn
 545 550 555 560
 Trp Pro Ser Leu Leu Arg His Arg Phe Leu Glu Glu Phe Ile Thr Pro
 565 570 575
 Ile Val Lys Val Ser Lys Asn Lys Gln Glu Met Ala Phe Tyr Ser Leu
 580 585 590
 Pro Glu Phe Glu Glu Trp Lys Ser Ser Thr Pro Asn His Lys Lys Trp
 595 600 605
 Lys Val Lys Tyr Tyr Lys Gly Leu Gly Thr Ser Thr Ser Lys Glu Ala
 610 615 620
 Lys Gln Tyr Phe Ala Asp Met Lys Arg His Arg Ile Gln Phe Lys Tyr
 625 630 635 640
 Ser Gly Pro Glu Asp Asp Ala Ala Ile Ser Leu Ala Phe Ser Lys Lys
 645 650 655
 Gln Ile Asp Asp Arg Lys Glu Trp Leu Thr Asn Phe Met Glu Asp Arg
 660 665 670
 Arg Gln Arg Lys Leu Leu Gly Leu Pro Glu Asp Tyr Leu Tyr Gly Gln
 675 680 685
 Thr Thr Thr Tyr Leu Thr Tyr Asp Asp Phe Ile Asn Lys Glu Leu Ile
 690 695 700
 Leu Phe Ser Asn Ser Asp Asn Glu Arg Ser Ile Pro Ser Met Val Asp
 705 710 715 720
 Gly Leu Lys Pro Gly Gln Arg Lys Val Leu Phe Thr Cys Phe Lys Arg
 725 730 735
 Asn Asp Lys Arg Glu Val Lys Val Ala Gln Leu Ala Gly Ser Val Ala
 740 745 750
 Glu Met Ser Ser Tyr His His Gly Glu Met Ser Leu Met Met Thr Ile
 755 760 765
 Ile Asn Leu Ala Gln Asn Phe Val Gly Ser Asn Asn Leu Asn Leu Leu
 770 775 780
 Gln Pro Ile Gly Gln Phe Gly Thr Arg Leu His Gly Gly Lys Asp Ser
 785 790 795 800
 Ala Ser Pro Arg Tyr Ile Phe Thr Met Leu Ser Ser Leu Ala Arg Leu
 805 810 815
 Leu Phe Pro Pro Lys Asp Asp His Thr Leu Lys Phe Leu Tyr Asp Asp
 820 825 830
 Asn Gln Arg Val Glu Pro Glu Trp Tyr Ile Pro Ile Ile Pro Met Val
 835 840 845

Leu Ile Asn Gly Ala Glu Gly Ile Gly Thr Gly Trp Ser Cys Lys Ile
 850 855 860
 Pro Asn Phe Asp Val Arg Glu Ile Val Asn Asn Ile Arg Arg Leu Met
 865 870 875 880
 5 Asp Gly Glu Glu Pro Leu Pro Met Leu Pro Ser Tyr Lys Asn Phe Lys
 885 890 895
 Gly Thr Ile Glu Glu Leu Ala Pro Asn Gln Tyr Val Ile Ser Gly Glu
 900 905 910
 10 Val Ala Ile Leu Asn Ser Thr Thr Ile Glu Ile Ser Glu Leu Pro Val
 915 920 925
 Arg Thr Trp Thr Gln Thr Tyr Lys Glu Gln Val Leu Glu Pro Met Leu
 930 935 940
 Asn Gly Thr Glu Lys Thr Pro Pro Leu Ile Thr Asp Tyr Arg Glu Tyr
 945 950 955 960
 15 His Thr Asp Thr Thr Val Lys Phe Val Val Lys Met Thr Glu Glu Lys
 965 970 975
 Leu Ala Glu Ala Glu Arg Val Gly Leu His Lys Val Phe Lys Leu Gln
 980 985 990
 Thr Ser Leu Thr Cys Asn Ser Met Val Leu Phe Asp His Val Gly Cys
 995 1000 1005
 20 Leu Lys Lys Tyr Asp Thr Val Leu Asp Ile Leu Arg Asp Phe Phe
 1010 1015 1020
 Glu Leu Arg Leu Lys Tyr Tyr Gly Leu Arg Lys Glu Trp Leu Leu
 1025 1030 1035
 25 Gly Met Leu Gly Ala Glu Ser Ala Lys Leu Asn Asn Gln Ala Arg
 1040 1045 1050
 Phe Ile Leu Glu Lys Ile Asp Gly Lys Ile Ile Ile Glu Asn Lys
 1055 1060 1065
 Pro Lys Lys Glu Leu Ile Lys Val Leu Ile Gln Arg Gly Tyr Asp
 1070 1075 1080
 30 Ser Asp Pro Val Lys Ala Trp Lys Glu Ala Gln Gln Lys Val Pro
 1085 1090 1095
 Asp Glu Glu Glu Asn Glu Glu Ser Asp Asn Glu Lys Glu Thr Glu
 1100 1105 1110
 Lys Ser Asp Ser Val Thr Asp Ser Gly Pro Thr Phe Asn Tyr Leu
 1115 1120 1125
 35 Leu Asp Met Pro Leu Trp Tyr Leu Thr Lys Glu Lys Lys Asp Glu
 1130 1135 1140
 Leu Cys Arg Leu Arg Asn Glu Lys Glu Gln Glu Leu Asp Thr Leu
 1145 1150 1155
 Lys Arg Lys Ser Pro Ser Asp Leu Trp Lys Glu Asp Leu Ala Thr
 1160 1165 1170
 40 Phe Ile Glu Glu Leu Glu Ala Val Glu Ala Lys Glu Lys Gln Asp
 1175 1180 1185
 Glu Gln Val Gly Leu Pro Gly Lys Gly Gly Lys Ala Lys Gly Lys
 1190 1195 1200
 45 Lys Thr Gln Met Ala Glu Val Leu Pro Ser Pro Arg Gly Gln Arg
 1205 1210 1215
 Val Ile Pro Arg Ile Thr Ile Glu Met Lys Ala Glu Ala Glu Lys
 1220 1225 1230
 Lys Asn Lys Lys Lys Ile Lys Asn Glu Asn Thr Glu Gly Ser Pro
 1235 1240 1245
 50 Gln Glu Asp Gly Val Glu Leu Glu Gly Leu Lys Gln Arg Leu Glu
 1250 1255 1260
 Lys Lys Gln Lys Arg Glu Pro Gly Thr Lys Thr Lys Lys Gln Thr
 1265 1270 1275
 Thr Leu Ala Phe Lys Pro Ile Lys Lys Gly Lys Lys Arg Asn Pro
 1280 1285 1290
 55

Trp Ser Asp Ser Glu Ser Asp Arg Ser Ser Asp Glu Ser Asn Phe
 1295 1300 1305
 Asp Val Pro Pro Arg Glu Thr Glu Pro Arg Arg Ala Ala Thr Lys
 1310 1315 1320
 Thr Lys Phe Thr Met Asp Leu Asp Ser Asp Glu Asp Phe Ser Asp
 1325 1330 1335
 Phe Asp Glu Lys Thr Asp Asp Glu Asp Phe Val Pro Ser Asp Ala
 1340 1345 1350
 Ser Pro Pro Lys Thr Lys Thr Ser Pro Lys Leu Ser Asn Lys Glu
 1355 1360 1365
 Leu Lys Pro Gln Lys Ser Val Val Ser Asp Leu Glu Ala Asp Asp
 1370 1375 1380
 Val Lys Gly Ser Val Pro Leu Ser Ser Ser Pro Pro Ala Thr His
 1385 1390 1395
 Phe Pro Asp Glu Thr Glu Ile Thr Asn Pro Val Pro Lys Lys Asn
 1400 1405 1410
 Val Thr Val Lys Lys Thr Ala Ala Lys Ser Gln Ser Ser Thr Ser
 1415 1420 1425
 Thr Thr Gly Ala Lys Lys Arg Ala Ala Pro Lys Gly Thr Lys Arg
 1430 1435 1440
 Asp Pro Ala Leu Asn Ser Gly Val Ser Gln Lys Pro Asp Pro Ala
 1445 1450 1455
 Lys Thr Lys Asn Arg Arg Lys Arg Lys Pro Ser Thr Ser Asp Asp
 1460 1465 1470
 Ser Asp Ser Asn Phe Glu Lys Ile Val Ser Lys Ala Val Thr Ser
 1475 1480 1485
 Lys Lys Ser Lys Gly Glu Ser Asp Asp Phe His Met Asp Phe Asp
 1490 1495 1500
 Ser Ala Val Ala Pro Arg Ala Lys Ser Val Arg Ala Lys Lys Pro
 1505 1510 1515
 Ile Lys Tyr Leu Glu Glu Ser Asp Glu Asp Asp Leu Phe
 1520 1525 1530

<210> 47

<211> 258

<212> PRT

<213> Homo sapiens

<400> 47

Met Leu Pro Leu Cys Leu Val Ala Ala Leu Leu Leu Ala Ala Gly Pro
 1 5 10 15
 Gly Pro Ser Leu Gly Asp Glu Ala Ile His Cys Pro Pro Cys Ser Glu
 20 25 30
 Glu Lys Leu Ala Arg Cys Arg Pro Pro Val Gly Cys Glu Glu Leu Val
 35 40 45
 Arg Glu Pro Gly Cys Gly Cys Cys Ala Thr Cys Ala Leu Gly Leu Gly
 50 55 60
 Met Pro Cys Gly Val Tyr Thr Pro Arg Cys Gly Ser Gly Leu Arg Cys
 65 70 75 80
 Tyr Pro Pro Arg Gly Val Glu Lys Pro Leu His Thr Leu Met His Gly
 85 90 95
 Gln Gly Val Cys Met Glu Leu Ala Glu Ile Glu Ala Ile Gln Glu Ser
 100 105 110

Leu Gln Pro Ser Asp Lys Asp Glu Gly Asp His Pro Asn Asn Ser Phe
 115 120 125
 Ser Pro Cys Ser Ala His Asp Arg Arg Cys Leu Gln Lys His Phe Ala
 130 135 140
 Lys Ile Arg Asp Arg Ser Thr Ser Gly Gly Lys Met Lys Val Asn Gly
 145 150 155 160
 Ala Pro Arg Glu Asp Ala Arg Pro Val Pro Gln Gly Ser Cys Gln Ser
 165 170 175
 Glu Leu His Arg Ala Leu Glu Arg Leu Ala Ala Ser Gln Ser Arg Thr
 180 185 190
 His Glu Asp Leu Tyr Ile Ile Pro Ile Pro Asn Cys Asp Arg Asn Gly
 195 200 205
 Asn Phe His Pro Lys Gln Cys His Pro Ala Leu Asp Gly Gln Arg Gly
 210 215 220
 Lys Cys Trp Cys Val Asp Arg Lys Thr Gly Val Lys Leu Pro Gly Gly
 225 230 235 240
 Leu Glu Pro Lys Gly Glu Leu Asp Cys His Gln Leu Ala Asp Ser Phe
 245 250 255
 Arg Glu

<210> 48

<211> 378

<212> PRT

<213> Homo sapiens

<400> 48

Met Asp Leu Gly Lys Pro Met Lys Ser Val Leu Val Val Ala Leu Leu
 1 5 10 15
 Val Ile Phe Gln Val Cys Leu Cys Gln Asp Glu Val Thr Asp Asp Tyr
 20 25 30
 Ile Gly Asp Asn Thr Thr Val Asp Tyr Thr Leu Phe Glu Ser Leu Cys
 35 40 45
 Ser Lys Lys Asp Val Arg Asn Phe Lys Ala Trp Phe Leu Pro Ile Met
 50 55 60
 Tyr Ser Ile Ile Cys Phe Val Gly Leu Leu Gly Asn Gly Leu Val Val
 65 70 75 80
 Leu Thr Tyr Ile Tyr Phe Lys Arg Leu Lys Thr Met Thr Asp Thr Tyr
 85 90 95
 Leu Leu Asn Leu Ala Val Ala Asp Ile Leu Phe Leu Leu Thr Leu Pro
 100 105 110
 Phe Trp Ala Tyr Ser Ala Ala Lys Ser Trp Val Phe Gly Val His Phe
 115 120 125
 Cys Lys Leu Ile Phe Ala Ile Tyr Lys Met Ser Phe Phe Ser Gly Met
 130 135 140
 Leu Leu Leu Leu Cys Ile Ser Ile Asp Arg Tyr Val Ala Ile Val Gln
 145 150 155 160
 Ala Val Ser Ala His Arg His Arg Ala Arg Val Leu Leu Ile Ser Lys
 165 170 175
 Leu Ser Cys Val Gly Ile Trp Ile Leu Ala Thr Val Leu Ser Ile Pro
 180 185 190
 Glu Leu Leu Tyr Ser Asp Leu Gln Arg Ser Ser Ser Glu Gln Ala Met
 195 200 205

Arg Cys Ser Leu Ile Thr Glu His Val Glu Ala Phe Ile Thr Ile Gln
 210 215 220
 Val Ala Gln Met Val Ile Gly Phe Leu Val Pro Leu Leu Ala Met Ser
 225 230 235
 Phe Cys Tyr Leu Val Ile Ile Arg Thr Leu Leu Gln Ala Arg Asn Phe
 245 250 255
 Glu Arg Asn Lys Ala Ile Lys Val Ile Ile Ala Val Val Val Val Phe
 260 265 270
 Ile Val Phe Gln Leu Pro Tyr Asn Gly Val Val Leu Ala Gln Thr Val
 275 280 285
 Ala Asn Phe Asn Ile Thr Ser Ser Thr Cys Glu Leu Ser Lys Gln Leu
 290 295 300
 Asn Ile Ala Tyr Asp Val Thr Tyr Ser Leu Ala Cys Val Arg Cys Cys
 305 310 315 320
 Val Asn Pro Phe Leu Tyr Ala Phe Ile Gly Val Lys Phe Arg Asn Asp
 325 330 335
 Leu Phe Lys Leu Phe Lys Asp Leu Gly Cys Leu Ser Gln Glu Gln Leu
 340 345 350
 Arg Gln Trp Ser Ser Cys Arg His Ile Arg Arg Ser Ser Met Ser Val
 355 360 365
 Glu Ala Glu Thr Thr Thr Thr Phe Ser Pro
 370 375

<210> 49

<211> 411

<212> PRT

<213> Homo sapiens

<400> 49

Met Ser Lys Arg Pro Ser Tyr Ala Pro Pro Pro Thr Pro Ala Pro Ala
 1 5 10 15
 Thr Gln Met Pro Ser Thr Pro Gly Phe Val Gly Tyr Asn Pro Tyr Ser
 20 25 30
 His Leu Ala Tyr Asn Asn Tyr Arg Leu Gly Gly Asn Pro Ser Thr Asn
 35 40 45
 Ser Arg Val Thr Ala Ser Ser Gly Ile Thr Ile Pro Lys Pro Pro Lys
 50 55 60
 Pro Pro Asp Lys Pro Leu Met Pro Tyr Met Arg Tyr Ser Arg Lys Val
 65 70 75 80
 Trp Asp Gln Val Lys Ala Ser Asn Pro Asp Leu Lys Leu Trp Glu Ile
 85 90 95
 Gly Lys Ile Ile Gly Gly Met Trp Arg Asp Leu Thr Asp Glu Glu Lys
 100 105 110
 Gln Glu Tyr Leu Asn Glu Tyr Glu Ala Glu Lys Ile Glu Tyr Asn Glu
 115 120 125
 Ser Met Lys Ala Tyr His Asn Ser Pro Ala Tyr Leu Ala Tyr Ile Asn
 130 135 140
 Ala Lys Ser Arg Ala Glu Ala Ala Leu Glu Glu Ser Arg Gln Arg
 145 150 155 160
 Gln Ser Arg Met Glu Lys Gly Glu Pro Tyr Met Ser Ile Gln Pro Ala
 165 170 175
 Glu Asp Pro Asp Asp Tyr Asp Asp Gly Phe Ser Met Lys His Thr Ala
 180 185 190

Thr Ala Arg Phe Gln Arg Asn His Arg Leu Ile Ser Glu Ile Leu Ser
 195 200 205
 Glu Ser Val Val Pro Asp Val Arg Ser Val Val Thr Thr Ala Arg Met
 210 215 220
 Gln Val Leu Lys Arg Gln Val Gln Ser Leu Met Val His Gln Arg Lys
 225 230 235 240
 Leu Glu Ala Glu Leu Leu Gln Ile Glu Glu Arg His Gln Glu Lys Lys
 245 250 255
 Arg Lys Phe Leu Glu Ser Thr Asp Ser Phe Asn Asn Glu Leu Lys Arg
 260 265 270
 Leu Cys Gly Leu Lys Val Glu Val Asp Met Glu Lys Ile Ala Ala Glu
 275 280 285
 Ile Ala Gln Ala Glu Glu Gln Ala Arg Lys Arg Gln Glu Glu Arg Glu
 290 295 300
 Lys Glu Ala Ala Glu Gln Ala Glu Arg Ser Gln Ser Ser Ile Val Pro
 305 310 315 320
 Glu Glu Glu Gln Ala Ala Asn Lys Gly Glu Glu Lys Lys Asp Asp Glu
 325 330 335
 Asn Ile Pro Met Glu Thr Glu Glu Thr His Leu Glu Glu Thr Thr Glu
 340 345 350
 Ser Gln Gln Asn Gly Glu Glu Gly Thr Ser Thr Pro Glu Asp Lys Glu
 355 360 365
 Ser Gly Gln Glu Gly Val Asp Ser Met Ala Glu Glu Gly Thr Ser Asp
 370 375 380
 Ser Asn Thr Gly Ser Glu Ser Asn Ser Ala Thr Val Glu Glu Pro Pro
 385 390 395 400
 Thr Asp Pro Ile Pro Glu Asp Glu Lys Lys Glu
 405 410

<210> 50

<211> 593

<212> PRT

<213> Homo sapiens

<400> 50

Met Ser Val Arg Tyr Ser Ser Ser Lys His Tyr Ser Ser Ser Arg Ser
 1 5 10 15
 Gly Gly Gly Gly Gly Gly Gly Cys Gly Gly Gly Gly Gly Val Ser
 20 25 30
 Ser Leu Arg Ile Ser Ser Ser Lys Gly Ser Leu Gly Gly Gly Phe Ser
 35 40 45
 Ser Gly Gly Phe Ser Gly Gly Ser Phe Ser Arg Gly Ser Ser Gly Gly
 50 55 60
 Gly Cys Phe Gly Gly Ser Ser Gly Gly Tyr Gly Gly Leu Gly Gly Phe
 65 70 75 80
 Gly Gly Gly Ser Phe His Gly Ser Tyr Gly Ser Ser Ser Phe Gly Gly
 85 90 95
 Ser Tyr Gly Gly Ser Phe Gly Gly Gly Asn Phe Gly Gly Gly Ser Phe
 100 105 110
 Gly Gly Gly Ser Phe Gly Gly Gly Gly Phe Gly Gly Gly Phe Gly
 115 120 125
 Gly Gly Phe Gly Gly Gly Phe Gly Gly Asp Gly Gly Leu Leu Ser Gly
 130 135 140

Asn Glu Lys Val Thr Met Gln Asn Leu Asn Asp Arg Leu Ala Ser Tyr
 145 150 155 160
 Leu Asp Lys Val Arg Ala Leu Glu Glu Ser Asn Tyr Glu Leu Glu Gly
 165 170 175
 Lys Ile Lys Glu Trp Tyr Glu Lys His Gly Asn Ser His Gln Gly Glu
 180 185 190
 Pro Arg Asp Tyr Ser Lys Tyr Tyr Lys Thr Ile Asp Asp Leu Lys Asn
 195 200 205
 Gln Ile Leu Asn Leu Thr Thr Asp Asn Ala Asn Ile Leu Leu Gln Ile
 210 215 220
 Asp Asn Ala Arg Leu Ala Ala Asp Asp Phe Arg Leu Lys Tyr Glu Asn
 225 230 235 240
 Glu Val Ala Leu Arg Gln Ser Val Glu Ala Asp Ile Asn Gly Leu Arg
 245 250 255
 Arg Val Leu Asp Glu Leu Thr Leu Thr Lys Ala Asp Leu Glu Met Gln
 260 265 270
 Ile Glu Ser Leu Thr Glu Glu Leu Ala Tyr Leu Lys Lys Asn His Glu
 275 280 285
 Glu Glu Met Lys Asp Leu Arg Asn Val Ser Thr Gly Asp Val Asn Val
 290 295 300
 Glu Met Asn Ala Ala Pro Gly Val Asp Leu Thr Gln Leu Leu Asn Asn
 305 310 315 320
 Met Arg Ser Gln Tyr Glu Gln Leu Ala Glu Gln Asn Arg Lys Asp Ala
 325 330 335
 Glu Ala Trp Phe Asn Glu Lys Ser Lys Glu Leu Thr Thr Glu Ile Asp
 340 345 350
 Asn Asn Ile Glu Gln Ile Ser Ser Tyr Lys Ser Glu Ile Thr Glu Leu
 355 360 365
 Arg Arg Asn Val Gln Ala Leu Glu Ile Glu Leu Gln Ser Gln Leu Ala
 370 375 380
 Leu Lys Gln Ser Leu Glu Ala Ser Leu Ala Glu Thr Glu Gly Arg Tyr
 385 390 395 400
 Cys Val Gln Leu Ser Gln Ile His Ala Gln Ile Ser Ala Leu Glu Glu
 405 410 415
 Gln Leu Gln Gln Ile Arg Ala Glu Thr Glu Cys Gln Asn Thr Glu Tyr
 420 425 430
 Gln Gln Leu Leu Asp Ile Lys Ile Arg Leu Glu Asn Glu Ile Gln Thr
 435 440 445
 Tyr Arg Ser Leu Leu Glu Gly Glu Gly Ser Ser Gly Gly Gly Arg
 450 455 460
 Gly Gly Gly Ser Phe Gly Gly Gly Tyr Gly Gly Gly Ser Ser Gly Gly
 465 470 475 480
 Gly Ser Ser Gly Gly Gly Tyr Gly Gly Gly His Gly Gly Ser Ser Gly
 485 490 495
 Gly Gly Tyr Gly Gly Gly Ser Ser Gly Gly Gly Ser Ser Gly Gly
 500 505 510
 Tyr Gly Gly Gly Ser Ser Ser Gly Gly His Gly Gly Gly Ser Ser Ser
 515 520 525
 Gly Gly His Gly Gly Ser Ser Ser Gly Gly Tyr Gly Gly Gly Ser Ser
 530 535 540
 Gly Gly Gly Gly Gly Gly Tyr Gly Gly Gly Ser Ser Gly Gly Gly Ser
 545 550 555 560
 Ser Ser Gly Gly Gly Tyr Gly Gly Gly Ser Ser Ser Gly Gly His Lys
 565 570 575
 Ser Ser Ser Ser Gly Ser Val Gly Glu Ser Ser Ser Lys Gly Pro Arg
 580 585 590
 Tyr

<210> 51

<211> 494

<212> PRT

<213> Homo sapiens

<400> 51

Met	Asp	Leu	Ser	Asn	Asn	Thr	Met	Ser	Leu	Ser	Val	Arg	Thr	Pro	Gly
1				5					10					15	
Leu	Ser	Arg	Arg	Leu	Ser	Ser	Gln	Ser	Val	Ile	Gly	Arg	Pro	Arg	Gly
			20					25					30		
Met	Ser	Ala	Ser	Ser	Val	Gly	Ser	Gly	Tyr	Gly	Gly	Ser	Ala	Phe	Gly
		35					40					45			
Phe	Gly	Ala	Ser	Cys	Gly	Gly	Gly	Phe	Ser	Ala	Ala	Ser	Met	Phe	Gly
	50					55					60				
Ser	Ser	Ser	Gly	Phe	Gly	Gly	Gly	Ser	Gly	Ser	Ser	Met	Ala	Gly	Gly
	65				70					75				80	
Leu	Gly	Ala	Gly	Tyr	Gly	Arg	Ala	Leu	Gly	Gly	Gly	Ser	Phe	Gly	Gly
				85					90					95	
Leu	Gly	Met	Gly	Phe	Gly	Gly	Ser	Pro	Gly	Gly	Gly	Ser	Leu	Gly	Ile
			100					105					110		
Leu	Ser	Gly	Asn	Asp	Gly	Gly	Leu	Leu	Ser	Gly	Ser	Glu	Lys	Glu	Thr
		115					120					125			
Met	Gln	Asn	Leu	Asn	Asp	Arg	Leu	Ala	Ser	Tyr	Leu	Asp	Lys	Val	Arg
	130					135						140			
Ala	Leu	Glu	Glu	Ala	Asn	Thr	Glu	Leu	Glu	Asn	Lys	Ile	Arg	Glu	Trp
	145				150					155					160
Tyr	Glu	Thr	Arg	Gly	Thr	Gly	Thr	Ala	Asp	Ala	Ser	Gln	Ser	Asp	Tyr
				165					170					175	
Ser	Lys	Tyr	Tyr	Pro	Leu	Ile	Glu	Asp	Leu	Arg	Asn	Lys	Ile	Ile	Ser
			180					185					190		
Ala	Ser	Ile	Gly	Asn	Ala	Gln	Leu	Leu	Leu	Gln	Ile	Asp	Asn	Ala	Arg
		195					200					205			
Leu	Ala	Ala	Glu	Asp	Phe	Arg	Met	Lys	Tyr	Glu	Asn	Glu	Leu	Ala	Leu
	210					215						220			
Arg	Gln	Gly	Val	Glu	Ala	Asp	Ile	Asn	Gly	Leu	Arg	Arg	Val	Leu	Asp
	225				230					235					240
Glu	Leu	Thr	Leu	Thr	Arg	Thr	Asp	Leu	Glu	Met	Gln	Ile	Glu	Ser	Leu
				245					250					255	
Asn	Glu	Glu	Leu	Ala	Tyr	Met	Lys	Lys	Asn	His	Glu	Asp	Glu	Leu	Gln
			260					265					270		
Ser	Phe	Arg	Val	Gly	Gly	Pro	Gly	Glu	Val	Ser	Val	Glu	Met	Asp	Ala
			275				280						285		
Ala	Pro	Gly	Val	Asp	Leu	Thr	Arg	Leu	Leu	Asn	Asp	Met	Arg	Ala	Gln
		290				295						300			
Tyr	Glu	Thr	Ile	Ala	Glu	Gln	Asn	Arg	Lys	Asp	Ala	Glu	Ala	Trp	Phe
	305				310					315					320
Ile	Glu	Lys	Ser	Gly	Glu	Leu	Arg	Lys	Glu	Ile	Ser	Thr	Asn	Thr	Glu
				325					330					335	
Gln	Leu	Gln	Ser	Ser	Lys	Ser	Glu	Val	Thr	Asp	Leu	Arg	Arg	Ala	Phe
			340				345						350		
Gln	Asn	Leu	Glu	Ile	Glu	Leu	Gln	Ser	Gln	Leu	Ala	Met	Lys	Lys	Ser
		355				360						365			
Leu	Glu	Asp	Ser	Leu	Ala	Glu	Ala	Glu	Gly	Asp	Tyr	Cys	Ala	Gln	Leu
		370				375					380				

Ser Gln Val Gln Gln Leu Ile Ser Asn Leu Glu Ala Gln Leu Leu Gln
 385 390 395 400
 Val Arg Ala Asp Ala Glu Arg Gln Asn Val Asp His Gln Arg Leu Leu
 405 410 415
 5 Asn Val Lys Ala Arg Leu Glu Leu Glu Ile Glu Thr Tyr Arg Arg Leu
 420 425 430
 Leu Asp Gly Glu Ala Gln Gly Asp Gly Leu Glu Glu Ser Leu Phe Val
 435 440 445
 10 Thr Asp Ser Lys Ser Gln Ala Gln Ser Thr Asp Ser Ser Lys Asp Pro
 450 455 460
 Thr Lys Thr Arg Lys Ile Lys Thr Val Val Gln Glu Met Val Asn Gly
 465 470 475 480
 Glu Val Val Ser Ser Gln Val Gln Glu Ile Glu Glu Leu Met
 485 490
 15
 <210> 52
 <211> 361
 <212> PRT
 20
 <213> Homo sapiens
 25
 <400> 52
 Cys Asn Trp Phe Cys Glu Gly Ser Phe Asn Gly Ser Glu Lys Glu Thr
 1 5 10 15
 Met Gln Phe Leu Asn Asp Arg Leu Ala Ser Tyr Leu Glu Lys Val Arg
 20 25 30
 30 His Val Glu Arg Asp Asn Ala Glu Leu Glu Asn Leu Ile Arg Glu Arg
 35 35 40 45
 Ser Gln Gln Gln Glu Pro Leu Leu Cys Pro Ser Tyr Gln Ser Tyr Phe
 50 55 60
 Lys Thr Ile Glu Glu Leu Gln Gln Lys Ile Leu Cys Ser Lys Ser Glu
 65 70 75 80
 35 Asn Ala Arg Leu Val Val Gln Ile Asp Asn Ala Lys Leu Ala Ala Asp
 85 90 95
 Asp Phe Arg Thr Lys Tyr Gln Thr Glu Gln Ser Leu Arg Gln Leu Val
 100 105 110
 Glu Ser Asp Ile Asn Ser Leu Arg Arg Ile Leu Asp Glu Leu Thr Leu
 115 120 125
 40 Cys Arg Ser Asp Leu Glu Ala Gln Met Glu Ser Leu Lys Glu Glu Leu
 130 135 140
 Leu Ser Leu Lys Gln Asn His Glu Gln Glu Val Asn Thr Leu Arg Cys
 145 150 155 160
 45 Gln Leu Gly Asp Arg Leu Asn Val Glu Val Asp Ala Ala Pro Ala Val
 165 170 175
 Asp Leu Asn Gln Val Leu Asn Glu Thr Arg Asn Gln Tyr Glu Ala Leu
 180 185 190
 Val Glu Thr Asn Arg Arg Glu Val Glu Gln Trp Phe Ala Thr Gln Thr
 195 200 205
 50 Glu Glu Leu Asn Lys Gln Val Val Ser Ser Ser Glu Gln Leu Gln Ser
 210 215 220
 Tyr Gln Ala Glu Ile Ile Glu Leu Arg Arg Thr Val Asn Ala Leu Glu
 225 230 235 240
 Ile Glu Leu Gln Ala Gln His Asn Leu Arg Tyr Ser Leu Glu Asn Thr
 245 250 255
 55

Leu Thr Glu Ser Glu Ala Arg Tyr Ser Ser Gln Leu Ser Gln Val Gln
 260 265 270
 Ser Leu Ile Thr Asn Val Glu Ser Gln Leu Ala Glu Ile Arg Ser Asp
 275 280 285
 Leu Glu Arg Gln Asn Gln Glu Tyr Gln Val Leu Leu Asp Val Arg Ala
 290 295 300
 Arg Leu Glu Cys Glu Ile Asn Thr Tyr Arg Ser Leu Leu Glu Ser Glu
 305 310 315 320
 Asp Cys Lys Leu Pro Ser Asn Pro Cys Ala Thr Thr Asn Ala Cys Glu
 325 330 335
 Lys Pro Ile Gly Ser Cys Val Thr Asn Pro Cys Gly Pro Arg Ser Arg
 340 345 350
 Cys Gly Pro Cys Asn Thr Phe Gly Tyr
 355 360

<210> 53

<211> 3282

<212> DNA

<213> Homo sapiens

<400> 53

atgaaggaga	tggtaggagg	ctgctgcgta	tgttoggacg	agaggggctg	ggccgagaac	60
ccgctgggtct	actgcgatgg	gcacgcgtgc	agcgtggcog	tccaccaagc	ttgctatggc	120
atcgttcagg	tgccaacggg	accctgggtc	tgccggaaat	gtgaatctca	ggagcgagca	180
gccagggtga	ggtgtgagct	gtgcccacac	aaagacgggg	cattgaagag	gactgataat	240
ggaggctggg	cacacgtggt	gtgtgcccct	tacatcccog	aggtgcaatt	tgccaacgtg	300
ctcaccatgg	agcccatcgt	gctgcagtac	gtgcctcatg	atcgtttcaa	caagacctgt	360
tacatctcgg	aggagacggg	cggggagagc	aaggcggcct	cgggagcctg	catgacctgt	420
aaccgcacatg	gatgtcgaca	agctttccac	gtcaacctgtg	cccaaattggc	aggcttgctg	480
tgtgagggaag	aagtgcctgga	ggtggacaac	gtcaagtact	gcggctactg	caaataccac	540
ttcagcaaga	tgaagacatc	coggoacagc	agcggggggag	gcggaggagg	cgctggaggga	600
ggagggtggca	gcatgggggg	aggtggcagt	ggtttcatct	ctgggaggag	aagccgggtca	660
gcctcaccat	ccacgcagca	ggagaagcac	cccaccccacc	acgagagggg	ccagaagaag	720
agtcgaaagg	acaaagaacg	ccttaagcag	aagcacaaga	agcggcctga	gtcgcccccc	780
agcatcctca	ccccgcccgt	ggtccccact	gctgacaagg	tctcctcctc	ggcttccctc	840
tctctccacc	acgagggccag	caogcaggag	acctctgaga	gcagcaggga	gtcaaagggg	900
aaaaagtctt	ccagccatag	cctgagtcac	aaagggaaga	aactgagcag	tgggaaaggt	960
gtgagcagtt	ttacctccgc	ctcctctctc	tctcctcctc	cttctctctc	ctctgggggg	1020
cccttccagc	ctgcagtcct	gtccctgcag	agctccccctg	acttctctgc	attcccccaag	1080
ctggagcagc	cagaggagga	caagtactcc	aagcccacag	cccccgcccc	ttcagccctc	1140
ccttctccct	cagctcccga	gccccccaag	gctgaccttt	ttgagcagaa	ggtggtcttc	1200
tctggctttg	ggcccatcat	gcgcttctcc	accaccaact	ccagctcagg	ccgggcccgg	1260
gcgccctccc	ctggggacta	taagtctccc	cacgtcacgg	ggtctggggc	ctcggcaggc	1320
acccacaaac	ggatgcccgc	actgagtgcc	accctctgtc	ctgctgatga	gacccttag	1380
acaggcctga	aggagaagaa	gcacaaagcc	agcaagagga	gccgccatgg	gccaggccgt	1440
cccaagggca	gccggaacaa	ggagggcact	ggggggccag	ctgccccatc	cttgcccagt	1500
gccagctgg	ctggctttac	cgccactgct	gcctcaccct	tctctggagg	ttccctggtc	1560
agctccggcc	tgggaggtct	gtccctccga	acctttgggc	cttctgggag	cttgcccagc	1620
ttgagcctgg	agtccccctt	actaggggca	ggcatctaca	ccagtaataa	ggaccccatc	1680
tcccacagtg	gcgggctgct	gcgggctgtc	tgagcaaccc	ctctctcctc	cagcctcctg	1740
gggccccag	ggacctgggc	cctgccccgc	ctcagccgct	ccccgttcac	cagcaccctc	1800
ccctctctct	ctgcttctat	ctccaccact	caggtgtttt	ctctggctgg	ctctaccttt	1860
agcctccctt	ctaccacat	ctttggaacc	cccatgggtg	ccgttaatcc	cctcctctcc	1920
caagctgaga	gcagccacac	agagccagac	ctggaggact	gcagcttccg	gtgtcggggg	1980

5 acctccccc accagagagtgt gtcttccatg tcccccatca gcagcctccc cgcactcttc 2040
 gaccagacag cctctgcacc ctgtgggggc ggccagtttag acccgggggc cccagggagc 2100
 actaacatgg agcagcttct ggagaagcag ggagcggggc aggcggggc caacatcggtg 2160
 gatagctga agggcgtgca cgcgtgcag aaggagaacc agcggctgca agagcagatc 2220
 ctgagcctga cggccaaaaa ggagcggctg cagattctca acgtgcagct ctctgtgccc 2280
 ttccctgccc tgcctgctgc cctgcctgcc gccaacggcc ctgtccctgg gccctatggc 2340
 ctgcctcccc aagccgggag cagcgactcc ttgagcacca gcaagagccc tccgggaaag 2400
 agcagcctcg gcctggacaa ctgctgttcc acttcttctg aggacccaca ctcaggctgc 2460
 ccgagccgca gcagctcgtc gctgtccttc cacagcacgc ccccaaccgt gccctcctc 2520
 10 cagcagagcc ctgccactct gccctgggcc ctgcctgggg cccctgcccc actcccgccc 2580
 cagccgcaga acgggttggg ccgggcaccc ggggcagcgg ggctgggggc catgccatg 2640
 gctgaggggc tgttgggggg gctggcaggc agtggggggc tgccctcaa tgggtcctt 2700
 ggggggttga atggggccgc tgccccaac ccgcgaagc tgagccaggc tggcggggcc 2760
 cccacgctgc agctgccagg ctgtctcaac agccttacag agcagcagag acatctcctt 2820
 15 cagcagcaag agcagcagct ccagcaactc cagcagctcc tggcctcccc gcagctgacc 2880
 ccggaacacc agactgttgt ctaccagatg atccagcaga tccagcagaa accgggagctg 2940
 cagcgtctgc agatggctgg gggctcccg ctgccatgg ccagcctgct ggcaggaagc 3000
 tccacccccc tgctgtctgc gggctacccc ggctgtctgc ccacagcgtc tgctccacc 3060
 ctgtgcccgc ctggagccct agtggtctcc tgccttgcca acaacacaag tctcatggcc 3120
 20 gcagcagctg cagctgcagc agtagcagca gcaggcggac ctccagtcct cactgcccag 3180
 accaaccctt tctcagcct gtccgggagca gagggcagtg gcgggtggcc caaaggaggg 3240
 accgctgaca aaggagcctc agccaaccag gaaaaaggct aa 3282

<210> 54

25 <211> 2227

<212> DNA

<213> Homo sapiens

30

<400> 54

35 gagagcccca acaggaagag ggtacagctt tgtgcaggtc acatgcccac tgcagccctc 60
 cagcctctgg tccccagagc ggacttttga agctgaactg cttttgttgc tggagaactt 120
 atgttataat ttaccctggg tggaccaggg tcgtacaaaa gggcaacgct cccagctccc 180
 cccactcccg accccggaat catgcctcgg actacaogga tcaaaatcac agagctgaac 240
 cccacctca tgtgtgcct ctgcgggggg gtgogctacc tggagaccaa caaatactgc 300
 tgctgtcatt ccttctgcaa aacctgcac gtgogctacc tggagaccaa caaatactgc 360
 40 cccatgtgtg acgtgcaggc ccataaaacc cggcgcgtgc tgagcatcag gtctgacaaa 420
 acacttcaag acattgtcta caaattgggt cctgggcttt ttaaagatga gatgaaacgg 480
 cggcgggatt tctatgcagc gtaccccccgt acggagggtc ccaacgggtc caatgaggac 540
 cgcggcgagg tcttggagca ggagaagggg gctctgagtg atgatgagat tgtcagcctc 600
 tccatcgaat tctacgaagg tgccagggac cgggatgaga agaaggggcc cctggagaat 660
 ggggatgggg acaaagagaa aacagggggt cgttccctgc gatgcccagc agccatgacc 720
 45 gtcattgcac ttgccaagt tctccgcaac aagatggatg tgcccagcaa gtacaagggtg 780
 gaggttctgt acgaggacga gccactgaag gaatactaca ccctcatgga catcgctac 840
 atctacccct ggcggcgga cgggcctctc cccctcaagt accgtgtoca gccagcctgc 900
 aagcggctca ccctagccac ggtgcccacc ccctccgagg gcaccaaacac cagcggggcg 960
 tccgagtgtg agtcagtcag cgacaaggct cccagccctg ccaccctgcc agccacctcc 1020
 tctccctgc ccagcccagc caccatcc catggctctc ccagtccca tgggcctcca 1080
 50 gccaccacac ctacctcccc cactccccct tcgacagcca gtggggccac cacagctgcc 1140
 aacgggggta gcttgaactg cctgcagaca ccatoctcca ccagcagggg gcgcaagatg 1200
 actgtcaacg gcgtcccggt gccccctta acttgaggcc agggaccctc tcccttcttc 1260
 cagccaagcc tctccactcc ttccactttt tctgggccc tttttccact tcttctactt 1320
 tccccagctc tccccacctt gggggtgggg ggcggggttt ataaaaaat atatataat 1380
 55 atgtacatag gaaaaaccaa atatacatat ttattttcta tggaccaacc agattaattt 1440
 aatgccaca ggaaacaaac tttatgtgtg tgtgtatgtg tggaaaatgg tgttcatttt 1500

	ttttgggggg	ggtcttgtgt	aatttgctgt	ttttgggggt	gcctggagat	gaactggatg	1560
	ggccactgga	gtctcaataa	agctctgcac	catcctcgct	gtttcccaag	gcagggtggtg	1620
	tggtgggggc	cccttcagac	ccaaagcttt	aggcatgatt	ccaactggct	gcataatagga	1680
5	gtcagttaga	attgtttott	tctctccccg	tttctctccc	catcttggct	gctgtcctgc	1740
	ctctgaccag	tggccgcccc	ccgcgttgtt	gaatgtccag	aaattgctaa	gaacagtgcc	1800
	ttttacaaat	gcagtttatc	cctggttctg	aggagcaagt	gcagggtgga	ggtggcacct	1860
	gcatcacctc	ctcctcttgc	agtggaaact	ttgtgcaaag	aatagatagt	tctgcctctt	1920
	tttttttttt	ttcctgtgtg	tgtggccttt	gcatcattta	tottgtggaa	aagaagattc	1980
	aggccctgag	aggtctcagc	tcttgaggga	gggctaaggc	tttagcattg	tgaagcgctg	2040
10	cacccccacc	aaaccttacc	tcaccgggga	accctcacta	gcaggactgg	tggtggagtc	2100
	tcacotgggg	cctagagtgg	aagtgggggt	gggttaacct	cacacaagca	cagatccacg	2160
	actttgccag	aggcaaacag	ggaattccgc	cgatactgac	gggctccagg	agtcgtcgcc	2220
	acactcg						2227
15	<210> 55						
	<211> 4283						
	<212> DNA						
20	<213> Homo sapiens						
	<400> 55						
25	ttgcgggaaa	gagccaaaac	ctggcgtttg	ggggcccggg	cggggagccc	ctcccgcggt	60
	ccacagcgac	gcctgccacg	ccctcctccc	cttcgggctc	cggcacgggg	ccccgaggcg	120
	ttcggaggcc	aggcgggttt	ctgtcaggcc	cggggaggag	gggcggggcg	ggcggccgct	180
	gcctcccccg	gacgggcccgt	accacgcgga	cggggaggac	ggggccaggg	gactgcaggg	240
	cggtgcacc	gcccgggggc	ggggtgcgga	gcgggccggc	gggctccccg	gggcggggcg	300
	ggagggcggg	gcgtggggcg	gacggaaacca	ccggggcggg	gtgggaggta	acgggacggg	360
30	cgcgaccatg	gcgcggtgag	ggagcggggg	tggggatcgg	tcggggggag	gcctgaggcc	420
	gctggccttg	gcgtgtcttc	cgccgcccc	ctctttcgcc	gccgcgcgcg	ccgccccggg	480
	catgtcgtcc	aactgcacca	gcaccaoggc	ggtggcggtg	gcgcgcgtca	gcgccagcaa	540
	gaccaagacc	aagaagaagc	atttcgtgtg	ccagaaagtg	aagctattcc	ggccacgca	600
	gccgatccct	agcgtcctga	tgtggggggt	gaaccacacg	atcaatgagc	tgagcaatgt	660
35	tcctgttctc	gtcatgctaa	tgccagatga	cttcaaagcc	tacagcaaga	tcaaggtgga	720
	caatcatctc	ttcaataagg	agaacctgcc	cagccgcttt	aagtttaagg	agtattgccc	780
	catggtgttc	cgaaaccttc	gggagaggtt	tggaaattgat	gatcaggatt	accagaattc	840
	agtgacgggc	agcgccccca	tcaacagtga	cagccagggt	cggtgtggca	cgcgtttcct	900
	caccacctac	gaccggcgct	ttgtcatcaa	gactgtgtcc	agcgaggacg	tggcggagat	960
40	gcacaacatc	ttaaagaaat	accaccagtt	tatagtggag	tgtcatggca	acacgccttt	1020
	gccacagttc	ctgggcatgt	acgcctgtac	cgtggatggt	gtggaaacct	acatggtggt	1080
	taccaggaac	gtgttcagcc	atcggtcac	tgtgcacgcg	aagtatgacc	tcaagggttc	1140
	tacggttgcc	agagaagcga	gcgacaagga	gaaggccaag	gacttgccaa	cattcaaaga	1200
	caatgacttc	ctcaatgaag	ggcagaagct	gcattgtggg	gaggagagta	aaaagaactt	1260
	cctggagaaa	ctgaagcggg	acgttgagtt	cttggcacag	ctgaagatca	tggactacag	1320
45	cctgctggtg	ggcatccacg	acgtggacog	ggcagagcag	gaggagatgg	aggtggagga	1380
	gcgggcagag	gacgaggagt	gtgagaatga	tggggtgggt	ggcaacctac	tctgtcctta	1440
	tggcacacct	ccggacagcc	ctggcaacct	cctcagcttt	cctcggttct	ttggtccttg	1500
	ggaattcgac	cctctgttgg	acgtctatgc	catgaaaagc	catgaaaagt	cccccaagaa	1560
	ggaggtgtat	ttcatggcca	tcattgatata	cctcacgcca	tacgatacaa	agaagaaagc	1620
50	tgcacatgct	gccaaaacgg	tgaacacagg	ggcaggggcc	gagatctcga	ctgtgaaccc	1680
	tgagcagtac	tccaaacgct	tcaacgagtt	tatgtccaac	atcctgacgt	agttctcttc	1740
	taccttcagc	cagagccaga	gagctggata	tggggtcggg	gatcgggagt	tagggagaag	1800
	ggtgtatttg	ggctagatgg	gaggggtggg	gcagagtcgg	gtttgggagg	gctttagcaa	1860
	tgagactgca	gcctgtgaca	cogaaagaga	cttttagctga	agaggagggg	gatgtgtctg	1920
	gtgtgcacct	gctcacagga	tgtaaaccca	ccttctgctt	acccttgatt	ttttctcccc	1980
55	atthgacacc	caggttaaaa	aggggttccc	tttttggtag	cttgtaacct	tttaagatac	2040

	cttgggggcta	gagatgactt	cgtgggttta	tttgggtttt	gtttotgaaa	tttcattgct	2100
	ccaggtttgc	tatttataat	catatttcat	cagcctaccc	accctcccca	tctttgctga	2160
	gctctcagtt	cccttcaatt	aaagagatac	ccagtagacc	cagcacaagg	gtccttccag	2220
5	aaccaagtgc	tatggatgcc	agattggaga	ggtcagacac	ctcgccctgc	tgcatttgct	2280
	cttgtctgga	ttaactttgt	aatttatgga	gtattgtgca	caacttcttc	cacctttccc	2340
	ttggattcaa	gtgaaaactg	ttgcattatt	cctccatcct	gtctggaata	caccaggtca	2400
	acaccagaga	tctcagatca	gaatcagaga	tctcagaggg	gaataagttc	atcctcatgg	2460
	gatggtgagg	ggcaggaaag	oggctgggct	cttggacacc	tggttctcag	agaaccctgt	2520
10	gatgatcacc	caagccccag	gctgtcttag	ccctggaggt	tcagaagtcc	tctctgtaaa	2580
	gcctgcctcc	cactagggtca	agaggaacta	gagtaccttt	ggatttatca	ggaccctcat	2640
	gtttaaatgg	ttattttccct	ttgggaaaac	ttcagaaact	gatgtatcaa	atgaggccct	2700
	gtgccctoga	tctattttcct	tcttccttct	gacctcctcc	caggcaactct	tacttctagc	2760
	cgaactctta	gctotgggca	gatctccaag	cgcttggagt	gcttttttagc	agagacacct	2820
15	cgttaagctc	cgggatgacc	ttgtaggaga	tctgtctccc	tgtgtctgga	gagttacagc	2880
	cagcaagggtg	ccccatcttt	agagtgtggt	gtccaaacgt	gaggtggctt	cctagttaaca	2940
	tgaggatgtg	atccaggaaa	tccagtttgg	aggcttgatg	tgggttttga	cctggcctca	3000
	gccttggggc	tgtttttcct	tggtgccccg	ctctagactt	ttagcagatc	tgccagccac	3060
	aggctttttt	ggaaggagtg	gcttcctgca	ggtgttccac	ctgccttcgg	agcctggcac	3120
20	ccaggccctc	agaactgagc	cacaggctgc	tctggccagg	agagaaacag	ctctgttgtt	3180
	ctgcattggg	ggaggtacat	tctgcatctc	tctcaccctc	tcaaccagga	actggggatt	3240
	tgggatgaga	tatggtcaga	cttgtagata	accccaaaga	tgtgaagatc	gcttgtgaaa	3300
	ccattttgaa	tgaatagatt	ggtttcctgt	ggctccctcc	aaacctggcc	aagcccagct	3360
	tccgaagcag	gaaccagcac	tgtctctgtg	cctgactcac	agcatatagg	tcaggaaaga	3420
	atggagacgg	cattcttgga	cttcactggg	gctgctggat	tggatgggaa	accttctgga	3480
25	agaggcagat	gggggtcaaa	ccactgcctt	ggccccagga	agggggcata	ggtaggtctg	3540
	aacaactgcc	gcaagaccac	tacatgactt	agggaaactg	aaaccaactg	gctcatggag	3600
	aaaacaaatt	tgacttggga	aagggattat	gtaggaataa	tgtttggact	tgatttcccc	3660
	acgtcataat	gaagaatgga	agtttggatc	tgtcctcgtc	caggcgcagc	atctctgaag	3720
	cttggaagc	tgtcttccag	cctccaaacc	tggccaagcc	cagcttcoga	agcaggaaacc	3780
30	agcactgtct	ctgtgcctga	ctcacagcat	ataggtcagg	aaagaatgga	gacggcattc	3840
	ttggacttca	ctggggctgc	tggattggat	gggaaacott	ctggaagagg	cagatggggg	3900
	tcaaaccact	gccttggccc	caggaagggg	ccataggtag	gtctgaacaa	ctgccgcaag	3960
	accactacat	gacttaggga	acttgaaacc	aaactggctca	tggagaaaac	aaatttgact	4020
	tgggaaaggg	attatgtagg	aataatgttt	ggacttgatt	tccccacgtc	ataatgaaga	4080
	atggaagttt	ggatctgctc	ctogtcaggc	gcagcatctc	tgaagcttgg	aaagctgtct	4140
35	tccagcagcc	tccgtggcct	cgggttccta	ccggcttctc	tgcatttggg	ctgctgatca	4200
	tgttgccata	atgtgtatgg	aaagtgtaac	acattcttac	tggttaaaga	cgactaccag	4260
	gtatctaact	tgtttaacat	tga				4283
	<210> 56						
40	<211> 6140						
	<212> DNA						
45	<213> Homo sapiens						
	<400> 56						
	gcggccgcag	cctgagccag	ggccccctcc	ctcgtcagga	ccggggcagc	aagcaggccg	60
50	ggggcaggtc	cgggcaccga	ccatgcgagg	cgagctctgg	ctcctggtgc	tgggtgctag	120
	ggaggtgccc	cgggcgctga	gccccagccc	cggagcaggt	cacgatgagg	gcccaggctc	180
	tggatgggct	gccaaaggga	ccgtgogggg	ctggaaccgg	agagcccag	agagccctgg	240
	gcatgtgtca	gagccggaca	ggaccagct	gagccaggac	ctgggtgggg	gcaccctggc	300
	catggacacg	ctgccagata	acaggaccag	ggtggtggag	gacaaccaca	gctattatgt	360
55	gtcccgctc	tatggcccca	gcgagcccca	cagccgggaa	ctgtgggtag	atgtggccga	420
	ggccaaccgg	agccaagtga	agatccacac	aatactctcc	aacacccacc	ggcaggcttc	480

	gagagtgggc	ttgtcctttg	atttcocctt	ctacggggcat	cctctgcggc	agatcaccat	540
	agcaactgga	ggcttcatct	tcatggggga	cgtgatccat	cggatgctca	cagctactca	600
	gtatgtggcg	ccctgatgg	ccaacttcaa	ccctggctac	tccgacaact	ccacagttgt	660
5	ttactttgac	aatgggacag	tctttgtggt	tcagtgggac	caogtttatc	tccaaggctg	720
	ggaagacaag	ggcagtttca	ccttcacggc	agctctgcac	catgacggcc	gcattgtctt	780
	tgccataaaa	gagatcccta	tgtctgtccc	ggaaatcagc	tctctccagc	atcctgtcaa	840
	aacoggcccta	tcggatgcct	tcatgattct	caatccatcc	ccggatgtgc	cagaatctcg	900
	gcgaaggagc	atctttgaat	accaccgcct	agagctggac	cccagcaagg	tcaccagcat	960
10	gtcggccgtg	gagttcaccc	cattgcccgc	ctgcctgcag	cataggagct	gtgacgcctg	1020
	catgtcctca	gacctgacct	tcaactgcag	ctgggtgccat	gtcctccaga	gatgctccag	1080
	tggctttgac	cgctatcgcc	aggagtggag	ggactatggc	tgtgcacagg	aggcagagg	1140
	caggatgtgc	gaggacctcc	aggatgagga	ccacgactca	gcctcccctg	acacttctct	1200
	cagccctcat	gatggagacc	tcaccactac	ctcctcctcc	ctcttcatcg	acagcctcac	1260
	cacagaagat	gacaccaagt	tgaatcccta	tgcaggagga	gaoggccttc	agaacaacct	1320
15	gtcccccaag	acaaagggca	ctcctgtgca	cctgggcacc	atcgtgggca	tcgtgctggc	1380
	agtcctctcc	gtggcggcca	tcatcctggc	tggaaattac	atcaatggcc	accccacatc	1440
	caatgctgog	ctcttcttca	togagcgtag	acctcaccac	tggccagcca	tgaagtctcg	1500
	cagccaccct	gaccattcca	cctatgcgga	gggtggagccc	tcgggccatg	agaaggagg	1560
	cttcatggag	gctgagcagt	gctgagaaca	ccaagtctcc	cctttgaaga	ctttgaggcc	1620
20	acagaaaaga	cagttaaagc	aaagaagaga	agtgaacttt	cctggcctct	cccagcatgc	1680
	cctgggctga	gatgagatgg	tggtttatgg	ctccagagct	gctgttgcct	tcgtcagcac	1740
	acccogaata	ttgaagaggg	ggccaaaaaa	caaccacatg	gattttttat	aggaacaaca	1800
	acctaatactc	atcctgtttt	gatgcaaggg	ttctcttctg	tgtcttgtaa	ccatgaaaca	1860
	gcagaagaac	taacataact	aactccatct	ttgtttaagg	ggcctttaac	tattcctgca	1920
25	cctaggctag	gataacttta	gagcactgac	ataaaacgca	aaaacaggaa	tcattgcctt	1980
	tgcaaaaacta	actctgggat	taaaggggaa	gcatgtaaac	agctaactgt	ttttgttaaa	2040
	gatttatagg	aatgaggagg	tttggctatt	gtcacatgac	agactgttag	ccaaggacaa	2100
	agaagtctcg	caaacctccc	ctggaccctt	gctgggtgct	agatgtctgc	ggttgtaagc	2160
	cccttccctt	ccccgcacct	aaacataaaa	gacaaggcaa	agcccgcata	attttaagac	2220
	ggttcttttag	gacattagtc	caccatcttc	ttggtttgct	ggctctccga	aataaagtc	2280
30	ctttccttgc	tccaactcct	tgtctctcaa	cgtattggct	atgacgcagc	aagcagaatg	2340
	aatttggact	cagttacagg	ctgtcaatgg	tctgctctgt	agcagctcca	gagcctccc	2400
	gaccactac	ctggagatag	ccagatagcc	agatgccctg	ctcctggcca	cctttaagc	2460
	ccctgcataat	gacacagggt	aactaaagtc	aagattgggg	ctgctgcatt	ccagggtccc	2520
	tagactcaca	agctggctct	tggccagggt	cagtggctca	cgctgtaat	cccagcactt	2580
35	tgggaggtcg	aggcaggcgg	atcacctgaa	gtcagaagtt	tgagaccago	ctggccaaca	2640
	taattaaaaat	gtctctacta	aaaatacaaa	aaattagctg	ggtgtggtga	cgcttgccctg	2700
	tatcccagct	actcaggaag	ctgagacacg	agaatcaact	gaacctggga	ggcagagggt	2760
	gcagtgaagt	cagatagtcg	cactgcactc	cagctgggt	gacagagcga	gactccgtct	2820
	caaaaaaaa	aaaagaaagc	agaacctcat	ggctaagag	ttggcatttt	agccccagct	2880
	tctgtagctc	tgaagccta	aagaaggtat	tctctccatc	tgtaaacac	agtatagtgg	2940
40	ctctcagccc	ttggggcatg	ttatcatggg	agggaagtca	aataagagga	gagaaaaagaa	3000
	ctcaaggggg	aaactgcatt	tttaggcttt	gctctcttac	cttgcccttt	ctactcagaa	3060
	ccaaataactt	ctgcatcaaa	acatgttaca	gcctgcatac	agggttttac	cccaacctgc	3120
	agccagcct	tcctgggtg	agcttgctat	gcgcagccac	atttaccatg	tggggctccc	3180
	tattctgatg	gcctgttcgg	tgcgggttt	actcaactgc	ctgttctgat	gtcagtgcct	3240
45	gtacatacct	ccaaaggcag	gacttgccctg	ataaatattt	ttcctcctct	gaactggatt	3300
	ttataggcat	taaagacaag	tcgggtggct	agagggctcc	ttgagacata	cctagcaggg	3360
	aactgcagggt	ggattctggt	gagaggcaaa	gcacctgagt	ggttgggaca	caggcagctg	3420
	gcatgggagg	gacttttttt	gagacagggt	ctcactgtgt	cgccagggc	aaggatgccc	3480
	aaagacacca	ggttggagag	gcacctgcc	actactgtct	ttccctggag	cctgcatgtg	3540
	cctgtgggggt	ggggaggcgt	aggggtctac	ggctgcctga	gatgggtgtg	cacagtgtgt	3600
50	gaagtaccta	cctcctggcc	ttgctggact	gtcagccagt	cgcagggccg	gccacaagac	3660
	ccatgtctcc	atctggtcat	actccatagc	taocaaagtt	acctgctcta	aactttggag	3720
	aactggatct	gtccaataaa	cgcttatttg	gccaaagcctg	atggctcgtg	cctgtactcc	3780
	cagcactttg	ggaggctgag	gtgggagggt	tgcttgagcc	caggggtttg	agaccagctt	3840
	gggcaacaac	aacaaaaatg	ccagggtgtg	tgggggtgcac	ctgtagtccc	agctactagg	3900
55	gaggctgagc	caggaggatc	acttgagccc	gggaggttga	ggctgcagtg	gggggtcata	3960
	atcatgccac	tgtactccag	cctgggtgac	agagtgcagc	cctgtctccg	aaaaaaaaaa	4020

	aaaaaaaaga	acggaaaaag	aaatgcttac	attgtcaggg	atcctgtaga	caatcattaa	4080
	ctctatgaga	tgcttgggtc	tatttttttg	ggagactttg	tccaagtgtt	ttggcottaag	4140
	aaatccatag	gcctctcttg	gtgacacatc	tctagtactt	tttgtcataa	acaaacaggc	4200
5	catctgcgcg	caaatacatc	cactccccat	gccactgaca	tcctatgggt	cagccaggct	4260
	tgctttgaat	gaggccgagg	catctggaac	tttctctgcc	tgccaggggt	agcagcagag	4320
	gcttcacacg	atcacaccac	cttccctcac	toctgacatt	ctttcccttc	agggatccaa	4380
	aatgggttgg	cgagctccca	gtgggaaaac	gtgtgctaga	gttggggagt	gagatgagt	4440
	gtgctgtcca	tggaatcagg	ccacagcagg	aactgcccc	ctggccattt	gagacacaca	4500
10	caggtggtaa	atgctctgct	ggtgggctgt	gcttccctca	ttcagagago	tctgttacag	4560
	ccactgtgt	cctttagaag	cttgaaagga	acccaactct	ttgtgtgact	gtcctttttc	4620
	ttcotcaaat	tcagaccctc	cttccaccgg	caccccctca	ctccaccctc	agctcttcct	4680
	tgcttgggtt	atcaagcaga	gctgaggccc	caagtttcca	actctgattg	tcacttgcat	4740
	cttcacaaag	gataaacacc	ggagcaactg	gaaaaccatc	agccaagcgt	tcggatgagt	4800
	ctggttattg	gtccaccccc	gaccagattc	ccttacactt	aactcaactc	tttctttggc	4860
15	aatgacccct	atgacatgta	taaattgggt	tgactaagaa	gaggctgtga	tctaactttt	4920
	atttgcgtcc	attttttact	ctggggagaa	gcagccccaa	ctcatcactg	ggaaagaact	4980
	ccccctgcaa	accagctaaa	tttgataatt	taaaccctct	gccccataaa	cttctcacag	5040
	agctggggag	ttggtggcaa	ctttccaagt	caaggtcttg	cttagaaagt	ccttcactac	5100
	atggccagggt	gcagtggctc	acgcctgtag	tcccagggtac	ttggggagcct	gaggcaggag	5160
20	gattgcttga	gctcaggagt	tcaaggctgc	agagagctat	gatcatocca	ctgcattttg	5220
	ttaaaaataa	atttttataa	tttgtgtgtt	ttatcagggg	tctcotgtac	agtgtatctg	5280
	tgatgttttg	tggtgtgtgt	tgtatacagc	cttgtttaat	gttttgagca	ataagatatg	5340
	cacacacagg	tattttgttg	ctaaagagat	tggaacaagg	tgtagctgtg	ctcaggcttc	5400
	agcttgggtt	gttaaatgta	gagataaaca	atgacaagag	ctgccagcca	accacactat	5460
	tcaaaaagca	aagtgttcac	cactaaagct	aaccattcat	ctggttgagc	gcaaggctaa	5520
25	ggctctctct	cctctagttc	ctggaacaga	ctcacagatt	ggcatgaagc	actgatcagg	5580
	ggctgcactc	agactccctg	gccaagcaaa	cctacaccag	aagagtccag	gtcacagata	5640
	tgatgcggcc	aatctctgtc	tccaaaaacc	tacctgaact	taatggtaga	attcaaagat	5700
	ctggggactg	agggcaccca	gccttctaaa	acacaatgta	ttcatgtgtt	tagtgtaaac	5760
	tctctgcatg	gattctcagt	gttaataata	aaaggaagca	ttctttttaca	actcotgctg	5820
30	tgtgcaaaaag	aaagtgcaaa	ggatttgagg	tggcattccg	aagatcacca	cacatacctt	5880
	ggttctgatg	gctgctgaac	tccgactctt	tcgctgagac	atgactgtgg	gaacagcctc	5940
	cagctatctg	ctcatcagag	gtgctttctt	caacctctct	caccacctcc	aagagaaaca	6000
	gootaaaaag	aaaccccagc	tgtttactta	tattggtctg	taaatccctg	gaagtaaacc	6060
	ccatgcattt	ttatctactg	tctgaggaca	tacaataaat	ctgagaaagt	ctatgctgtc	6120
	aaaaaaaaaa	aaaaaaaaaa					6140
35	<210> 57						
	<211> 2098						
40	<212> DNA						
	<213> Homo sapiens						
45	<400> 57						
	gcaggagcac	gtggagaggg	cgggtagcca	cagcggcagc	tccagcccg	cccggcagcg	60
	acatggaaga	tatacaaaaca	aatgcggaac	tgaaaagcac	tcaggagcag	tctgtgcccc	120
	cagaaagtgc	agcggttttg	aatgactaca	gtttaaccaa	atctcatgaa	atggaaaatg	180
	tggacagtgg	agaaggccca	gccaatgaag	atgaagacat	aggagatgat	tcaatgaaag	240
50	tgaagatgta	atacagtga	agagatgaga	atgtttttaa	gtcagaaccc	atgggaaatg	300
	cagaagagcc	tgaatccct	tacagctatt	caagagaata	taatgaatat	gaaaacatta	360
	agttggagag	acatgtttgc	tcattcgata	gtagcaggcc	aaccagtgga	aagatgaact	420
	gcgatgtgtg	tggattatoc	tgcatcagct	tcaatgtctt	aatggttcat	aagcgaagcc	480
	atactggtga	acgcccattc	cagtgtaatc	agtgtggggc	atcttttact	cagaaaggta	540
55	acctcctccg	ccacattaaa	ctgcacacag	gggaaaaacc	ttttaagtgt	cacctctgca	600
	actatgcatg	ccaaagaaga	gatgcgctca	cggggcatct	taggacacat	tctgtggaga	660

	aaccctacaa	atgtgagttt	tgtggaagga	gttacaagca	gagaagttcc	cttgaggagc	720
	acaaggagcg	ctgccgtaca	ttttctcaga	gcactgaccc	aggggacact	gcaagtgcgg	780
	aggcaagaca	catcaaagca	gagatgggaa	gtgaaagagc	tctcgtactg	gacagattag	840
5	caagcaatgt	ggcaaaacga	aaaagctcaa	tgcctcagaa	attcattggt	gagaagcgcc	900
	actgctttga	tgtcaactat	aattcaagtt	acatgtatga	gaaagagagt	gagctcatat	960
	agaccogcat	gatggacca	gccatcaata	acgccatcag	ctatcttggc	gccgaagccc	1020
	tgtgccccct	ggtccagaca	cgcctgctc	ccacctcgga	gatggttcca	gttatcagca	1080
	gcatgtatcc	catagccctc	acccgggctg	agatgtcaaa	cggtgccctc	caagagctgg	1140
10	aaaggaaaag	catcctcctt	ccagagaaga	gcgtgccttc	tgagagaggc	ctctctccca	1200
	acaatagtgg	ccacgactcc	acggacactg	acagcaacca	tgaagaacgc	cagaatcaca	1260
	tctatcagca	aaatcacatg	gtcctgtctc	gggcccgcga	tgggatgcca	cttctgaagg	1320
	aggttccccg	ctottacgaa	ctcctcaagc	ccccgcccat	ctgcccaga	gactctgtca	1380
	aagtgatcga	caaggaagg	gaggtgatgg	atgtgtatcg	gtgtgaccac	tgcgcgctcc	1440
	tcttcttgga	ctatgtgatg	ttcacgattc	acatgggctg	ccacggcttc	cgtgaccctt	1500
15	togagtgtaa	catgtgtgga	gatcgaagcc	atgatcggtg	tgaattctcg	tctcacatag	1560
	ccagaggaga	acacagaagc	ctgctgaagt	gaatatctgg	tctcagggat	tgctcctatg	1620
	tattcagcat	cgtttctaaa	aacagttgac	ctcgcctaac	agattgctct	caaaacatac	1680
	tcagttocaa	acttcttttc	ataccatttt	tagctgtggt	cacaggggta	gccagagaaa	1740
	cactgtcttc	cttcagaaat	tattcgcagg	tctagcatat	tattactttt	gtgaaacctt	1800
20	tgttttccca	tcagggactt	gaattttatg	gaatttaaaa	gcaaaaaagg	tatttgggtca	1860
	ttatcttcta	cagcagtgga	atgagtggtc	ccggagatgt	gctatatgaa	acattctttc	1920
	tgagatata	caaccacacg	tggaaaagcc	tttcagtcac	acatgcaaat	ccacaaagag	1980
	gaagagotga	ccagctgacc	ttgctgggaa	gcctcaccct	tctgcccttc	acaggctgaa	2040
	gggttaagat	ctaactctcc	taatctaaat	gacagtctaa	gagtaagtaa	aagaacag	2098
25	<210> 58						
	<211> 2947						
	<212> DNA						
30	<213> Homo sapiens						
	<400> 58						
35	atgccaatcc	ctcctccccc	gccaccccca	cctggtcctc	ctccacctcc	cacatttcat	60
	caggcaaaaca	cagagcagcc	caagctgagt	agagatgagc	agcgggggtcg	agggcgcctc	120
	ttacaggaca	tttgcaagg	gaccaagctg	aagaaggatga	ccaacattaa	tgatcggagt	180
	gtccccatcc	togagaagcc	gaaaggaagc	agtgggtggc	atggctctgg	aggagctgcc	240
	ctgcagccca	agggaggctc	cttccaagga	ggagtgtctg	agcttcgacc	tgtgggagcc	300
40	aaggatgggt	cagagaacct	agctggtaag	ccagccctgc	aaatccccag	ttctogagct	360
	gctgccccaa	ggcctccagt	atctgccgcc	agcgggctgc	ctcaggatga	tacagacagc	420
	agcggggcct	cactcccaga	actgccccgg	atgcagagac	cctctttaac	ggacctctct	480
	cggcctaata	ccaccagcag	taaggggcag	aagcacagct	cctctgcccc	tccccacca	540
	ccccaggggc	ggcgtgccaa	cgcacccccc	acacctctgc	ctatgcacag	cagcaaagcc	600
45	ccgcctaca	acagagagaa	acccttgcca	ccgacgcctg	gacaaaggct	tcacctgggt	660
	cagaggggac	ctcctgctcc	acccccagtc	aaacacctc	cttccccctg	gaatatcaga	720
	acaggaccaa	gtggccagtc	tctggctcct	cctcctccgc	cttaccgcca	gcctcctggg	780
	gtccccaatg	gacctcttag	ccccactaat	gagtcagccc	ctgagctgcc	acagagacac	840
	aattcttttc	ataggaagac	accagggcct	gtcagaggcc	tagcacctcc	tccaccacc	900
	tcggcctccc	catctttact	gagtaatagg	ccacctcccc	cagcccagaga	ccctcccagt	960
50	cggggagcag	ctcctccacc	cccaccacct	gtgatccgaa	atgggtgccag	ggatgctccc	1020
	cctccccccac	caccataccg	aatgcatggg	tcagaacccc	cgaagccagg	aaagccccc	1080
	cctccaccct	caaggacgcc	agctgggcca	ccccctcctc	ctccaccgcc	cctgagggaat	1140
	ggccacagag	attctatcac	caactgtccg	tctttcttgg	atgattttga	gtcaaagtat	1200
	tccttccatc	cagtagaaga	ctttcctgct	ccagaagaat	ataaacactt	tcagaggata	1260
55	tatcccagca	aaacaaaccg	agctgcccgt	ggagccccac	ctctgccacc	cattctcagg	1320
	tgaagcctga	cttggctccg	ttcctcagga	aaaggatgga	ccttctcttc	ttctcagatg	1380

	gtcccttcca	ttccctgaa	acotgcatga	gagctoctaa	catgtttctc	caatgcaatc	1440
	aagccctaga	ctccaaatgt	cctcccagct	cacctccatc	tatgcatctc	atctctggat	1500
	ttggtgatca	gactctatat	tgacagtagg	atctcaaacc	ctgcatccat	ccttctctcca	1560
5	gcaagccctg	ctagccacat	gaggaacaag	tttccgtgtc	ttctgccttc	ctcttgggga	1620
	aagggtgoot	gttgtgatga	attaactcac	tgtaggggca	gggtggagaa	tggtactcct	1680
	tccttctcct	gtccactgtg	ggggaagcct	ggcaggtata	ttatatttca	tcatttagga	1740
	ggctggcatg	accaggactt	atgggtggga	ggggagcatt	tttagtgaag	caagaaagga	1800
	gtttgccaag	aagtgatctg	ttttaaaggt	catatttgga	gaaagggcaa	ggaattgggt	1860
	ctgctttatt	tttgggggta	ttttgttttt	gttctcacct	gctgcccccc	caccccacca	1920
10	ccccagggat	aaattggata	taaacactaa	atactaatac	gttgaactta	acatttaata	1980
	aaaagaaagg	gtgaaataaa	ctgaagacca	ttttagaact	agtcagttct	ctgcagcaaa	2040
	gggaacagga	gccatttgaa	ccctctggga	ccctccacc	cactgcttca	gggtgctagg	2100
	ctgagggatg	tttttctctc	cccttaccgc	ccatgccctt	gaaagaaaag	tcactttttg	2160
	tggaaggcat	catttcattcc	tgattcacia	acccccaaaa	cctctggtgg	gagataggaa	2220
15	gatagggcgt	gggcctgggc	ottaacctca	atcttgtgtc	tgctcagtc	ttttctgact	2280
	ggccctgaag	ttgtcagtgg	ctctttctgt	ccttcagccc	ctggaagggtg	ctccaggata	2340
	acaaagaagg	gcaggttgaa	gcccctcatg	gaaggagctg	gctttgtggg	gctgcaaagg	2400
	acttttaagt	cctgcctgta	ctgaagttca	cagcccacct	gactgagcag	actcttctctg	2460
	ttcctttctc	taccaccctt	gccttcccag	gactgcacgg	tttaacacag	cagagtacag	2520
20	aagggtgaag	aagtgagcag	aggcttatga	agatattcag	atactcttct	atgccaggaa	2580
	gcacaaagac	tttgttgaga	tttgccctcag	ttcagtagat	cttccctggc	agccagccat	2640
	agggtgttct	tttgtctctc	gggtcctaaa	gagcacagag	aaaatggagg	tccccagctc	2700
	aggtaggaag	ctgattggat	gaggacttct	ttttttccga	cagcaggatg	gggtctcttg	2760
	gctccacaca	ccagatgctt	tggttttcta	caactgttgc	tatgtgtaga	gggtgctcag	2820
	agcgtggcat	gagagcaagg	agaccatggc	tactctttga	aatggatggg	gaaaatttagc	2880
25	ttaaaaattt	aatcacgaga	ttgcgccact	gcactccagc	ctgggcgaca	gagccagact	2940
	ccgtctct						2947
	<210> 59						
30	<211> 784						
	<212> DNA						
	<213> Homo sapiens						
35	<400> 59						
	gagcggttgc	gcagtgaagg	ctagaccocg	tttactggaa	ttgctctggc	gatcgagggg	60
	toctagtaca	ccgcaatcat	gtctattatg	toctataacg	gaggggcoct	catggccatg	120
40	aagggggaaga	actgtgtggc	oatogctgca	gacaggcgct	tcgggatcca	ggcccagatg	180
	gtgaccacgg	acttccagaa	gatctttccc	atgggtgacc	ggctgtacat	cggtctggcc	240
	gggctogcca	ctgacgtcca	gacagttgcc	cagcgcccca	agttccggct	gaacctgtat	300
	gagttgaagg	aaggtcggca	gatcaaacct	tataccctca	tgagcatggt	ggccaacctc	360
	ttgtatgaga	aacggtttgg	cccttactac	actgagccag	tcattgccgg	gttggaccocg	420
45	aagaccttta	agcccttcat	ttgctctcta	gaacctcatg	gctgccccat	gggtactgat	480
	gactttgtgg	tcagtggcac	ctgcgcgcaa	caaattgtacg	gaatgtgtga	gtccctctgg	540
	gagcccaaca	tggtatccgga	tcacctgttt	gaaacctctc	cccaagccat	gctgaatgct	600
	gtggaccggg	atgcagtgtc	aggcatggga	gtcattgtcc	acatcatoga	gaaggacaaa	660
	atcaccacca	ggacactgaa	ggcccgaatg	gactaaccct	gttcccagag	cccacttttt	720
	tttctttttt	tgaaataaaa	tagcctgtct	ttcaaaaaaa	aaaaaaaaaa	aaaaaaaaaa	780
50	aaaa						784
55							

<210> 60

<211> 3033

<212> DNA

<213> Homo sapiens

<400> 60

atactoctaa	gctcctcccc	cggcggcgag	ccaggagaa	aggatggccg	gcctggcgcc	60
gcggttggtc	ctgctagctg	gggcagcgcc	gctggcgagc	ggctcccagg	gcgaccgtga	120
gocggtgtac	cgcgactcgc	tactgcagtg	cgaagagcag	aactgctctg	ggggogctct	180
gaatcacttc	cgctccccgc	agocaaatota	catgagtota	gcaggctgga	cctgtcggga	240
cgactgtaag	tatgagtgtg	tgtgggtcac	cgttgggctc	tacctocagg	aaggtcacaa	300
agtgcctcag	ttocatggca	agtggccctt	ctcccggttc	ctgttctttc	aagagccggc	360
atcgcccggtg	gcctcgtttc	tcaatggcct	ggccagcctg	gtgatgctct	gccgtaccg	420
caccttcgtg	ccagcctcct	cccccatgta	ccacacctgt	gtggccttcg	cctgggtgtc	480
cctcaatgca	tggttctggt	ccacagtctt	ccacaccagg	gacactgacc	tcacagagaa	540
aatggactac	ttctgtgcct	ccactgtcat	cctacactca	atctacctgt	gotgogtcag	600
gtgagcctgc	ctgggtggct	gcaggggcaa	aatcgaaacc	tgggggcaga	aaggggtcac	660
ccagccttcc	cctggggggc	ttcttcaacta	gtctcccaac	acctacgccc	cccaaccccc	720
aacacatcag	ctgtcctggg	tgaggactct	ggggtaggac	tgggggccct	ggctcctgac	780
aaggagctgt	agcacttgct	gcccagctgt	ggcctgtttg	gtggggagag	gggtagtac	840
ttcagggggc	atgcaccaat	gttgggggga	ggagatgctt	cagggaatgc	tgctctgggg	900
atgggcccacc	tgcctctgta	gcaaccctgg	acgggtggggc	aggaccgtgg	ggctgcagca	960
cccagctgtg	gtcagtgcct	tccgggctct	cctgctgctc	atgctgaocg	tgacgtctc	1020
ctacctgagc	ctcatccgct	togactatgg	ctacaacctg	gtggccaacg	tggctattgg	1080
cctggtcaac	gtggtgtggt	ggctggcctg	gtgcctgtgg	aaccagcggc	ggctgcctca	1140
cgtgcgcaag	tgcgtggtgg	tgggtcttgc	gctgcagggg	ctgtccctgc	tcgagctgct	1200
tgacttccca	ccgctcttct	gggtcctgga	tgcccatgac	atctggcaca	tcagcaccat	1260
ccctgtccac	gtcctctttt	tcagctttct	ggaagatgac	agcctgtacc	tgctgaagga	1320
atcagaggac	aagttcaagc	tggactgaag	accttgagac	gagtctgccc	cagtggggat	1380
cctgcccccg	ccctgctggc	ctcccttctc	ccctcaaccc	ttgagatgat	tttctctttt	1440
caactctctg	aacttgagca	tgaaggatgt	gggccagaa	tcagtgtggc	agccaccccc	1500
ctgttgcccc	tcaccagcct	tggagtctgt	tcagggaag	gcctcccagc	atctgggact	1560
cgagagtggg	cagccctct	acctcctgga	gctgaactgg	ggtggaactg	agtgtgttct	1620
tagctctacc	gggaggacag	ctgcctgttt	ctcccccacc	agcctcctcc	ccacatcccc	1680
agctgcctgg	ctgggtcctg	aagccctctg	tctacctggg	agaccagggg	ccacaggcct	1740
tagggataca	gggggtcccc	ttctgttacc	acccccacc	ctcctccagg	acaccactag	1800
gtggtgctgg	atgcttgttc	tttggccagc	caagggtcac	ggcgattctc	cccatgggat	1860
cttgagggac	caagctgctg	ggattgggaa	ggagtttcac	cctgaccgtt	gccctagcca	1920
ggttcccagg	aggcctcacc	atactccctt	tcaggggccag	ggctccagca	agcccagggc	1980
aaggatcctg	tgctgctgtc	tggttgagag	cctgccaccg	tgtgtcggga	gtgtgggcca	2040
ggctgagtgc	ataggtgaca	gggccttgag	catgggcctg	ggtgtgtgtg	agctcaggcc	2100
taggtgcgca	gtgtggagac	gggtgttgtc	ggggaagagg	tgtggcttca	aagtgtgtgt	2160
gtgcaggggg	tgggtgtgtt	agcgtgggtt	aggggaaogt	gtgtgcgcgt	gctgggtggc	2220
atgtgagatg	agtgaactgc	ggtgaatgtg	tccacagtgt	agaggttgga	gcaggatgag	2280
ggaatcctgt	caccatcaat	aatcaactgt	ggagcgccag	ctctgcccac	gacgccacct	2340
gggcggacag	ccaggagctc	tccatggcca	gctgcctgt	gtgcatgttc	cctgtctggt	2400
gccccctttg	ccgcctcctg	caaacctcac	agggtcccca	cacaacagtg	ccctccagaa	2460
gcagccccctc	ggaggcagag	gaaggaaaat	ggggatggct	ggggctctct	ccatcctcct	2520
tttctccttg	ccttcgcatg	gctggccttc	ccctccaaaa	cctccattcc	cctgctgcca	2580
gccccctttg	catagcctga	ttttggggag	gagggaaggg	cgatttgagg	gagaagggga	2640
gaaagcttat	ggctgggtct	ggtttcttcc	cttcccagag	ggtcttactg	ttccaggggtg	2700
gccccagggc	aggcaggggc	cacactatgc	ctgcgcctg	gtaaaagtga	cccctgccat	2760
ttaccagcag	ccctggcatg	ttcctgcccc	acagggaatag	aatggaggga	gctccagaaa	2820
ctttccatcc	caaaggcagt	ctccgtgggt	gaagcagact	ggatttttgc	tctgccctgt	2880

accccttgtc	cctctttgag	ggaggggagc	tatgctagga	ctccaacctc	agggactcgg	2940
gtggcctgcg	ctagcttctt	ttgatactga	aaacttttaa	ggtgggaggg	tggcaaggga	3000
tgtgcttaat	aaatcaattc	caagcctcac	ctg			3033

5 <210> 61

<211> 1174

10 <212> DNA

<213> Homo sapiens

15 <400> 61

aagctcctcc	cccggcggcg	agccagggag	aaaggatggc	cggcctggcg	gogcggttgg	60
tcctgctagc	tggggcagcg	gcgctggcga	gcggctccca	gggcgaccgt	gagccggtgt	120
accgcgactg	cgtactgcag	tgcgaagagc	agaactgctc	tgggggcgct	ctgaatcact	180
tccgctcccg	ccagccaatc	tacatgagtc	tagcaggctg	gacctgtcgg	gacgactgta	240
agtatgagtg	tatgtgggtc	accgttgggc	tctaacctcca	ggaaggtcac	aaagtgcctc	300
agttccatgg	caagtggccc	ttctcccggt	tcctgttctt	tcaagagccg	gcacggtccg	360
tggcctcggt	tcctcaatggc	ctggccagcc	tggtgatgct	ctgccgctac	cgcaccttcg	420
tgccagcctc	ctcccccatg	taccacacct	gtgtggcctt	cgcctgggtg	tcctcfaatg	480
catggttctg	gtccacagtc	ttccacacca	gggacactga	cctcacagag	aaaatggact	540
acttctgtgc	ctccactgtc	atcctacact	caatctaacct	gtgctgcgtc	aggaccgtgg	600
ggctgcagca	cccagctgtg	gtcagtgcct	tcggggtctc	cctgctgctc	atgctgaccg	660
tgacagctctc	ctacctgagc	ctcatccgct	tgcactatgg	ctacaacctg	gtggccaacg	720
tggtctattg	cctggtcaac	gtggtgtggt	ggctggcctg	gtgcctgtgg	aaccagcggc	780
ggctgcctca	cgtgcgcaag	tgogtgggtg	tggtcttgcct	gctgcagggg	ctgtccctgc	840
tcgagctgct	tgacttccca	ccgctcttct	gggtccctgga	tgcccatgoc	atctggcaca	900
tcagcaccat	ccctgtccac	gtcctctttt	tcagctttct	ggaagatgac	agcctgtacc	960
tgctgaagga	atcagaggac	aagttcaagc	tggttgaagc	agactggatt	tttgctctgc	1020
ccctgacccc	ttgtccctct	ttgagggagg	ggagctatgc	taggactcca	acctcagggg	1080
ctcgggtggc	ctgcgctagc	ttcttttgat	actgaaaact	tttaagggtg	gaggggtggca	1140
agggatgtgc	ttaataaatc	aattccaagc	ctca			1174

35 <210> 62

<211> 3167

40 <212> DNA

<213> Homo sapiens

45 <400> 62

aagctcctcc	cccggcggcg	agccagggag	aaaggatggc	cggcctggcg	gogcggttgg	60
tcctgctagc	tggggcagcg	gcgctggcga	gcggctccca	gggcgaccgt	gagccggtgt	120
accgcgactg	cgtactgcag	tgcgaagagc	agaactgctc	tgggggcgct	ctgaatcact	180
tccgctcccg	ccagccaatc	tacatgagtc	tagcaggctg	gacctgtcgg	gacgactgta	240
agtatgagtg	tatgtgggtc	accgttgggc	tctaacctcca	ggaaggtcac	aaagtgcctc	300
agttccatgg	caagtggccc	ttctcccggt	tcctgttctt	tcaagagccg	gcacggtccg	360
tggcctcggt	tcctcaatggc	ctggccagcc	tggtgatgct	ctgccgctac	cgcaccttcg	420
tgccagcctc	ctcccccatg	taccacacct	gtgtggcctt	cgcctgggtg	agaaaactga	480
ggcacagcaa	ggctaaataa	cttgcccaag	gacacacagg	aaatgcagag	ccaggaactg	540
aaccctggca	gtctggctgt	agggcttgca	ttcttaataa	taccactacc	tcctcaaatct	600
gaggaaaggg	tgctccctcaa	tgcctggttc	tggtccacag	tcttccacac	cagggacact	660

	gacctcacag	agaaaatgga	ctaattctgt	gcctccactg	tcatcctaca	ctcaatctac	720
	ctgtgctgcg	tcaggtgagc	ctgcoctgggt	ggctgcaggg	gcaaaatcga	accctggggg	780
	cagaaagggg	tcaccacagc	ttcccctggg	ggccttcttc	actagtctcc	caacacctac	840
5	gccccccaac	cccccaacaca	tcagctgtcc	tgggtgagga	ctctggggta	ggactggggg	900
	ccctggctcc	tgacaaggag	ctgtagcact	tgttgocccag	ctgtggcctg	tttgggtggg	960
	agaggggtag	tgacttcagg	ggccatgcac	caatgttggg	gggaggagat	gottcaggga	1020
	atgctgctct	ggggatgggc	caoctgcoct	ctgagcaacc	ctggacgggtg	gggcaggacc	1080
	gtggggctgc	agcaccacgc	tgtggtcagt	gccttcocggg	ctctcctgot	gctcatgctg	1140
	accgtgcaog	tctcctacct	gagcctcatc	cgcttcogact	atggctacaa	cctgggtggc	1200
10	aacgtggcta	ttggcctggt	caacgtgggt	tgttggtgctg	cctgggtgco	gtggaaccag	1260
	cgggcgctgc	ctcacgtgcg	caagtgcgtg	tgttggtgct	tgtgctgca	ggggctgtcc	1320
	ctgctcgagc	tgcttgactt	cccaccgctc	ttctgggtcc	tggatgccc	tgccatctgg	1380
	cacatcagca	ccatccctgt	ccacgtcctc	tttttcagct	ttctggaaga	tgacagcctg	1440
	tacctgctga	aggaatcaga	ggacaagtcc	aagctggact	gaagaccttg	gagcgagtct	1500
15	gccccagtgg	ggatcctgcc	cccgccctgc	tggootccct	tctccctcca	acccttgaga	1560
	tgattttctc	ttttcaactt	cttgaacttg	gacatgaagg	atgtggggcc	agaatcatgt	1620
	ggccagccca	ccccctgttg	gcoctcacca	gccttgaggt	ctgttctagg	gaaggcctcc	1680
	cagcatctgg	gactcgagag	tgggcagccc	ctctacctcc	tggagctgaa	ctggggtgga	1740
	actgagtgtg	ttcttagctc	taocgggagg	acagctgcct	gtttcctccc	caccagcctc	1800
20	ctcccacat	ccccagctgc	ctggctgggt	cctgaagccc	tctgtctacc	tgggagacca	1860
	gggaccacag	gccttaggga	tacaggggtg	ccccttctgt	taccaccccc	caccctcctc	1920
	caggacacca	ctaggtggtg	ctggatgctt	gttctttggc	cagccaaggt	tcacggcgat	1980
	tctcccctag	ggatcttgag	ggaccaagct	gctgggattg	ggaaggaggt	tcaccctgac	2040
	cgttgcctca	gccaggttcc	caggaggcct	caccatactc	cctttcaggg	ccagggtctcc	2100
	agcaagccca	gggcaaggat	cctgtgctgc	tgtctgggtg	agagcctgoc	accgtgtgtc	2160
25	gggagtgtgg	gccaggctga	gtgcataggt	gacagggccg	tgagcatggg	cctgggtgtg	2220
	tgtgagctca	ggcctaggtg	cgcagtgtgg	agaoggggtg	tgtcggggaa	gaggtgtggc	2280
	ttcaaaagtgt	gtgtgtgcag	gggggtgggtg	tgttagcgtg	ggttagggga	acgtgtgtgc	2340
	gcgtgctggt	gggcatgtga	gatgagtgc	tgccgggtgaa	tgtgtccaca	gttgagaggt	2400
	tggagcagga	tgaggggaatc	ctgtcaccat	caataatcac	ttgtggagcg	ccagctctgc	2460
30	ccaagacgcc	acctgggcgg	acagccagga	gctctccatg	gccaggctgc	ctgtgtgcat	2520
	gttccctgtc	tggtgcccct	ttgcccgcct	cctgcaaacc	tcacaggggc	cccacacac	2580
	agtgcctctc	agaagcagcc	cctcgagggc	agaggaagga	aaatggggat	ggctggggct	2640
	ctctccatcc	tccttttctc	cttgccctcg	catggctggc	cttcccctcc	aaaacctcca	2700
	ttcccctgct	gccagcccct	ttgccatagc	ctgatttttg	ggaggaggaa	ggggcgattt	2760
	gagggagaag	gggagaaaagc	ttatggctgg	gtctgggttc	ttcccttccc	agagggtctt	2820
35	actgttccag	ggtggcccca	gggcaggcag	gggccacact	atgcoctgoc	cctggtaaaag	2880
	gtgaccocctg	ccattttacca	gcagccctgg	catgttccctg	ccccacagga	atagaatgga	2940
	gggagctcca	gaaactttcc	atcccaaagg	cagtctccgt	ggttgaagca	gactggattt	3000
	ttgctctgcc	cctgaccctc	tgtccctctt	tgaggagggg	gagctatgct	aggactccaa	3060
	cctcagggac	tcgggtggcc	tgcgctagct	tcttttgata	ctgaaaactt	ttaagggtggg	3120
40	aggggtggcaa	gggatgtgct	taataaatca	attccaagcc	tcacctg		3167

<210> 63

<211> 2733

45 <212> DNA

<213> Homo sapiens

50

55

<220>

<221> misc_feature

<222> (2694) .. (2694)

<223> n=a, c, g or t

<220>

<221> misc_feature

<222> (2724) .. (2724)

<223> n=a, c, g or t

<400> 63

aggagagaaag	gatggccggc	ctggcggcgc	gggtggctct	gctagctggg	gcagcggcgc	60
tggcgagcgg	ctcccagggc	gaccgtgagc	cggtgtaccg	cgactgcgta	ctgcagtgcg	120
aagagcagaa	ctgctctggg	ggcgctctga	atcacttccg	ctcccgccag	ccaatctaca	180
tgagtctagc	aggctggacc	tgtcgggacg	actgtaagta	tgagtgtatg	tgggtcaccg	240
ttgggtctota	cctccaggaa	ggtcacaaag	tgccctcagtt	ccatggcaag	tggcccttct	300
cccggttctt	gttctttcaa	gagccggcat	cggcogtggc	ctogttttct	aatggcctgg	360
ccagcctggg	gatgctctgc	cgctaccgca	ccttcgtgcc	agcctcctcc	cccatgtacc	420
acacctgtgt	ggccttcgcc	tgggtgtccc	tcaatgcatt	gttctggctc	acagtcttcc	480
acaccaggga	cactgacctc	cagagaaaat	ggactacttc	tgtgcctcct	gtatcctaca	540
ctcaatotac	ctgtgctgcg	tcaggaccgt	ggggctgcag	cacccagctg	tgggtcaagt	600
ccttcggggc	tctcctgctg	ctcatgctga	ccgtgcacgt	ctcctacctg	agcctcatcc	660
gcttcgacta	tggctacaac	ctgggtggcca	acgtggctat	tggcctgggc	aacgtgggtg	720
gggtggctgg	ctgggtgcct	tgggaaccag	ggcggctgcc	tcacgtgcgc	aagtgcgtgg	780
tgtgtgtctt	gctgctgcag	gggctgtccc	tgactgcagc	gcttgacttc	ccaccgctct	840
tctgggtctt	ggatgcccat	gccatctggc	acatcagcac	catccctgtc	cacgtcctct	900
ttttcagctt	tctggaagat	gacagcctgt	acctgctgaa	ggaatcagag	gacaagtcca	960
agctggactg	agaccttgga	gcgaagtctg	ccccagtggt	gatcctgccc	ccgccctgct	1020
ggcctccctt	ctccctccaa	cccttgagat	gattttctct	tttcaacttc	ttgaacttgg	1080
acatgaagga	tgtgggcccc	gaatcatgtg	gccagccccc	ccctggttgg	ccctcaccag	1140
ccttgaggct	tgtctagggg	aaggcctccc	agcatctggg	actcgagagt	gggcagcccc	1200
tctacctctt	ggactgaact	ggggtggaac	tgagtgtgtt	cttagctcta	ccgggaggac	1260
agctgcctgt	ttctctccca	ccagcctcct	ccccacatcc	ccagctgcct	ggctgggtcc	1320
tgaagccctc	tgtctacctg	ggagaccagg	gtaccacagg	ccttagggat	acaggggggc	1380
cccttctgtt	accacccccc	accctcctcc	aggacaccac	taggtgggtg	tggatgcttg	1440
ttctttggcc	agccaagggt	cacggcgatt	ctcccatagg	gatcttgagg	gaccaagctg	1500
ctgggattgg	gaaggagttt	caccctgacc	gttgccctag	ccagggtccc	aggaggcctc	1560
accatactcc	ctttcagggc	cagggtccca	gcaagccag	ggcaaggatc	ctgtgctgct	1620
gtctgggttg	gagcctgcca	ccgtgtgtcg	ggagtgtggg	ccaggctgag	tgcattaggt	1680
acagggccgt	gagcatgggc	ctgggtgtgt	gtgagctcag	gcctaggtgc	gcagtgtgga	1740
gacgggtgtt	gtcggggaag	aggtgtggct	tcaaagtgtg	tgtgtgcagg	gggtgggtgt	1800
gttagcgtgg	gttaggggaa	cgtgtgtgcg	cgtgctgggt	ggcatgtgag	atgagtgact	1860
gccggtgaat	gtgtccacag	ttgagaggtt	ggagcaggat	gagggaaatc	tgtcaccatc	1920
aataatcaat	tgtggagcgc	cagctctgcc	caagacgcca	cctgggcgga	cagccaggag	1980
ctctccatgg	ccaggtgcgc	tgtgtgcatt	ttccctgtct	ggtgcccctt	tgcccgcctc	2040
ctgcaaacct	ccacagggtc	ccacacaaca	gtgccctcca	gaagcagccc	ctcgagggca	2100
gaggaaggaa	aatggggatg	gctggggctc	tctccatcct	ccttttctcc	ttgccttcgc	2160
atggctggcc	ttccctccca	aaacctccat	tccctgctgc	ccagcccctt	tgccatagcc	2220
tgattttggg	gaggaggaag	gggcgatttg	agggagaagg	ggagaaagct	tatggctggg	2280
tctgggtttct	tcccttccca	gagggtctta	ctgttccagg	gtggccccag	gcagcagggc	2340
cacactatgc	ctgcgcctcg	gtaaaggtga	ccccctgcat	ttaccagcag	ccctggcatg	2400

ttcctgcccc	acaggaatag	aatggagggg	gctccagaaa	ctttccatcc	caaaggcagt	2460
ctccgtggtt	gaagcagact	ggattttttg	tctgcccctg	accccttgct	cctcttttag	2520
ggaggggagc	tatgctagga	ctccaacctc	agggaactgg	gtggcctgcg	ctagcttctt	2580
ttgatactga	aaacttttaa	ggtgggaggg	tggcaagggg	tgtgcttaag	oggccgcgaa	2640
ttcaaaaagc	ttctcgagag	tactttctaga	gcgccgcggg	gcccatcgat	tttnccaccc	2700
gggtggggta	cccaggtaag	tgtnccccat	atc			2733

<210> 64

<211> 2546

<212> DNA

<213> Homo sapiens

<400> 64

aagctccctcc	cccggcggcg	agccagggag	aaaggatggc	cgccctggcg	gcgcgggttg	60
tcctgctagc	tggggcagcg	gcgctggcga	gcggctccca	gggcgaccgt	gagccgggtgt	120
accgcgactg	cgactgagc	tgcgaagagc	agaactgctc	tggggcgctg	ctgaatcact	180
tccgctcccg	ccagccaatc	tacatgagtc	tagcaggctg	gacctgtcgg	gacgactgta	240
agtatgagtg	tatgtgggtc	accgttgggc	tctacctcca	ggaaggctac	aaagtgcctc	300
agttccatgg	caagtggccc	ttctcccggt	tcctgttctt	tcaagagccg	gcctcggccg	360
tggcctcggt	tctcaatggc	ctggccagcc	tggtagtgct	ctgccgtac	cgcaccttcg	420
tgccagcctc	ctcccccatg	taccacacct	gtgtggcctt	cgccctgggtg	tcctcaatg	480
catggtttctg	gtccacagtc	ttccacacca	gggacactga	cctcacagag	aaaatggact	540
acttctgtgc	ctccactgtc	atcctacact	caatctacct	gtgctgctgc	aggcctgggtc	600
aacgtgggtg	ggtggctggc	ctggtgcctg	tggaaaccagc	ggcggctgcc	tcacgtgcgc	660
aagtgcgtgg	tgggtggtctt	gctgctgcag	gggctgtccc	tgctcgagct	gcttgacttc	720
ccaccgctct	tctgggtcct	ggatgcccct	gccatctggc	acatcagcac	catccctgtc	780
cacgtccctc	ttttcagctt	tctggaagat	gacagctgtg	acctgctgaa	ggaatcagag	840
gacaagttca	agctggactg	aagacctggg	agcagtgctg	ccccagtggg	gacctgtccc	900
cgcctctgct	ggcctccctt	ctccctccaa	cccttgagat	gattttctct	tttcaacttc	960
ttgaacttgg	acatgaagga	tgtggggccc	gaatcatgtg	gccagcccac	cccctgttgg	1020
ccctcaccag	ccttgagtc	tgttctaggg	aaggcctccc	agcatctggg	actcgagagt	1080
gggcagcccc	tctacctcct	ggagctgaac	tggggtggaa	ctgagtgtgt	tcttagctct	1140
accgggagga	cagctgcctg	tttccctccc	accagcctcc	tccccacatc	cccagctgoc	1200
tggctgggtc	ctgaagccct	ctgtctacct	gggagaccag	ggaccacagg	ccttagggat	1260
acagggggtc	cccttctgtt	accaccccc	accctcctcc	aggacaccac	taggtgggtc	1320
tggatgcttg	ttctttggcc	agccaagggt	cacggcgatt	ctcccatagg	gatcttgagg	1380
gaccaagctg	ctgggattgg	gaaggagttt	caacctgacc	gttgccctag	ccaggttccc	1440
aggaggcctc	accatactcc	ctttcagggc	cagggctcca	gcaagcccag	ggcaaggatc	1500
ctgtgctgct	gtctggttga	gagcctgcc	ccgtgtgtcg	ggagtgtggg	ccaggctgag	1560
tgcataagtg	acagggccgt	gagcatgggc	ctgggtgtgt	gtgagctcag	gocctaggtc	1620
gcagtgtgga	gacgggtggt	gtoggggaag	aggtgtggct	tcaaagtgtg	tgtgtgcagg	1680
gggtgggtgt	gttagcgtgg	gttaggggaa	cgtgtgtgct	cgtgtgtgtg	ggcatgtgag	1740
atgagtgaact	gocggtgaat	gtgtccacag	ttgagaggtt	ggagcaggat	gagggaaatcc	1800
tgtcaccatc	aataatcact	tgtggagcgc	cagctctgcc	caagacgcca	cctgggcgga	1860
cagccaggag	ctctccatgg	ccaggctgcc	tgtgtgcatg	ttccctgtct	ggtgcccctt	1920
tgcccgcctc	ctgcaaacct	cacagggtcc	ccacacaaca	gtgccctcca	gaagcagccc	1980
ctcgagggca	gaggaaggaa	aatggggatg	gctggggctc	tctccatcct	ccttttctcc	2040
ttgccttogg	atggctggcc	ttccctccca	aaacctccat	tccctgtctg	ccagcccctt	2100
tgccatagcc	tgattttggg	gaggaggaag	gggcgatttg	agggagaagg	ggagaaagct	2160
tatggctggg	tctggtttct	tcccttccca	gagggtotta	ctgttccagg	gtggccccag	2220
ggcaggcagg	ggccacacta	tgcctgcgcc	ctggtaaagg	tgaccctgc	catttaccag	2280
cagccctggc	atgttctctg	cccacaggaa	tagaatggag	ggagctccag	aaactttcca	2340
tccc aaaggc	agtctccgtg	gttgaagcag	actggatttt	tgctctgccc	ctgaccctt	2400
gtccctcttt	gagggagggg	agctatgcta	ggactocaa	ctcagggaact	cggtgtggct	2460

gcgctagctt cttttgatac tgaaaacttt taagggtggga ggggtggcaag ggatgtgctt 2520
aataaatcaa ttccaagcct cacctg 2546

<210> 65

<211> 2683

<212> DNA

<213> Homo sapiens

<400> 65

aagctcctcc cccggcgggg agccagggag aaaggatggc cggcctggcg gcgcgggttg 60
tcctgctagc tggggcagcg gcgctggcga gcggctccca gggcgaccgt gagccgggtg 120
accgcgactg cgtactgcag tgcgaagagc agaactgctc tgggggcgct ctgaatcact 180
tcgcctcccg ccagccaatc tacatgagtc tagcaggctg gacctgtcgg gacgactgta 240
agtatgagtg tatgtgggtc accgttgggc tctacctcca ggaaggtcac aaagtgcctc 300
agttccatgg caagtggccc ttctcccggt tcctgttctt tcaagagccg gcatcgggcg 360
tggcctcggt tctcaatggc ctggccagcc tgggtgatgt ctgcgcgtac cgcaccttcg 420
tgccagcoto ctccccatg taccacacct gtgtggcctt cgcctgggtg tccctcaatg 480
catggttctg gtccacagtc ttccacacca gggacactga cctcacagag aaaatggact 540
acttctgtgc ctccactgtc atcctacact caatctacct gtgtgcgtc aggaccgtgg 600
ggctgcagca cccagctgtg gtcagtgcct tcggggctct cctgtgtctc atgctgaccg 660
tgacagtctc ctacctgagc ctcatccgct tcgactatgg ctacaacctg gtggccaacg 720
tggctatttg cctgggtaac gtggtgtggt ggctggcctg gtgcctgtgg aaccagcggc 780
ggctgcctca cgtgcgcaag tgogtgggtg tgggtttgtc gctgcagggg ctgtccctgc 840
tcgagctgct tgacttccca ccgctcttct gggctcctga tgcccatgcc atctggcaca 900
tcagcaccat ccctgtccac gtccctcttt tcagctttct ggaagatgac agcctgtacc 960
tgctgaagga atcagaggac aagttcaagc tggactgaag accttggagc gagtctgccc 1020
cagtggggat cctgcccccg cctctgtggc ctcccttctc cctcaaccc ttgagatgat 1080
ttctcttttt caacttcttg aacttggaca tgaaggatgt gggtccagaa tcatgtggcc 1140
agcccccccc ctgttggccc tcaccagcct tggagtctgt tctagggaag gcctcccagc 1200
atctgggact cgagagtggg cagccccctc acctcctgga gctgaactgg ggtggaactg 1260
agtgtgttct tagctctacc gggaggacag ctgcctgttt cctccccacc agcctcctcc 1320
ccacatcccc agctgcctgg ctgggtcctg aagccctctg tctacctggg agaccagggg 1380
ccacaggcct tagggatata ggggttcccc ttctgttacc accccccacc ctccctcagg 1440
acaccaactag gtggtgctgg atgcttgttc ttgtgccagc caaggttcac ggcgattctc 1500
ccatgggat cttgagggac caagctgctg ggattgggaa ggagtttcac cctgaccgtt 1560
gcoctagcca ggttccagc aggcctcacc ataactcctt tcagggccag ggctccagca 1620
agcccagggc aaggatcctg tgcctgtgtc tgggtgagag cctgccaccg tgtgtcggga 1680
gtgtgggcca ggctgagtgc ataggtgaca gggcctgtg catgggcctg ggtgtgtgtg 1740
agctcaggcc taggtgcgca gtgtggagac ggggtgtgtc ggggaagagg tgtggcttca 1800
aagtgtgtgt gtgcaggggg tgggtgtgtt agcgtgggtt aggggaacgt gtgtgcgctg 1860
gctgggtggg atgtgagatg agtgactgcc ggtgaatgtg tccacagttg agaggttggg 1920
gcaggatgag ggaatcctgt caccatcaat aatcacttgt ggagcgccag ctctgcccc 1980
gacgccacct gggcgagcag ccaggagctc tccatggcca ggctgcctgt gtgcatgttc 2040
cctgtctggt gcccttttgc ccgcctcctg caaacctcac aggttcccca cacaacagt 2100
cctccagaa gcagccctc ggaggcagag gaaggaaaat ggggatggct ggggctctct 2160
ccatcctcct tttctccttg ccttcgcatg gctggccttc cctccaaaa cctccattcc 2220
cctgtgcca gcccttttgc catagcctga ttttggggag gaggaagggg cgatttgagg 2280
gagaagggg gaaagcttat ggctgggtct ggtttcttcc cttcccagag ggtcttactg 2340
ttccagggtg gcccagggc aggcaggggc cacactatgc ctgcgcctg gtaaagggtg 2400
ccccgccat ttaccagcag cctggcatg ttccctgccc acaggaatag aatggaggga 2460
gtccagaaa ctttccatcc caaaggcagt ctccgtgggt gaagcagact ggatttttgc 2520
tetgcccctg accccttgtc cctcttttag ggaggggagc tatgctagga ctccaacctc 2580
agggactcgg gtggcctgcg ctagcttctt ttgatactga aaacttttaa ggtgggaggg 2640
tggcaaggga tgtgttaaat aaatcaattc caagcctcac ctg 2683

<210> 66

<211> 2341

<212> DNA

<213> Homo sapiens

<400> 66

aagctcctcc	cccggcggcg	agccaggag	aaaggatggc	cgccctggcg	gcgcgggttg	60
tcctgctagc	tggggcagcg	gcgctggcga	gcggctccca	gggcgaccgt	gagcgggtgt	120
accgcgactg	cgtactgcag	tgogaagagc	agaactgctc	tgggggcgct	ctgaatcact	180
tccgcctccg	ccagccaatc	tacatgagtc	tagcaggctg	gacctgtogg	gacgactgta	240
agtatgagtg	tatgtgggtc	acogttgggc	tctacctoca	ggaaggtcac	aaagtgcctc	300
agttccatgg	caagtggccc	ttotcccggg	tcotgttott	tcaagagccg	gcacgcggcg	360
tggcctcggt	totcaatggc	ctggccagcc	tggtgatgct	ctgccgctac	cgcaccttcg	420
tgccagcctc	ctcccccatg	taaccacact	gtgtggcctt	cgccctgggtg	tcctcaatg	480
catggtttctg	gtccacagtc	ttccacacca	gggacactga	cctcacagag	aaaatggact	540
acttctgtgc	ctccactgtc	atcctacact	caatctacct	gtgctgcgtc	agctttctgg	600
aagatgacag	cctgtacctg	ctgaaggaat	cagaggacaa	gttcaagctg	gactgaagac	660
cttggagcga	gtctgcccc	gtggggatcc	tgcccccgcc	ctgctggcct	cccttctccc	720
ctcaaccctt	gagatgattt	tctcttttca	acttcttgaa	cttggacatg	aaggatgtgg	780
gcccagaatc	atgtggccag	cccacccctt	gttggccctc	accagccttg	gagtctgttc	840
tagggaaggc	ctcccagcat	ctgggactcg	agagtgggca	gccccctac	ctcctggagc	900
tgaactgggg	tggaactgag	tgtgtttctt	gctctaccgg	gaggacagct	gcctgtttcc	960
tccccaccag	cctcctcccc	acatccccag	ctgcctggct	gggtcctgaa	gccctctgtc	1020
tacctgggag	accagggacc	acaggcctta	gggatacagg	gggtccccct	ctgttaccac	1080
ccccaccctt	cctccaggac	accactaggt	ggtgctggat	gcttgttctt	tggccagcca	1140
aggttcacgg	cgattctccc	catgggatct	tgaggggacca	agctgctggg	attgggaagg	1200
agtttcaccc	tgaccgttgc	cctagccagg	ttcccaggag	gcctcaccat	actccctttc	1260
agggccaggg	ctccagcaag	cccaggggcaa	ggatootgtg	ctgctgtctg	gttgagagcc	1320
tgccaccgtg	tgtcgggagt	gtggggccagg	ctgagtgcac	aggtgacagg	gccgtgagca	1380
tgggcctggg	tgtgtgtgag	ctcaggccca	ggtgcgcagt	gtggagacgg	gtgttgtcgg	1440
ggaagagggtg	tggcttcaaa	gtgtgtgtgt	gcaggggggtg	ggtgtgttag	cgtgggttag	1500
gggaacgtgt	gtgcgcgtgc	tggtgggcat	gtgagatgag	tgaactgtgtc	tgaatgtgtc	1560
cacagttagg	agggttgagg	aggatgaggg	aatcctgtca	ccatcaataa	tcacttgttg	1620
agcgcacagt	ctgcccaga	cgccacctgg	gcggacagcc	aggagctctc	catggccagg	1680
ctgcctgtgt	gcatgttccc	tgtotggtgc	ccotttgccc	gcctcctgca	aacctcacag	1740
ggtccccaca	caacagtgcc	ctccagaagc	agccoctcgg	aggcagagga	aggaaaatgg	1800
ggatggctgg	ggtctctctc	atcctccttt	tctccttgcc	ttcgcattgg	tggccttccc	1860
ctccaaaacc	tccattcccc	tgtgtccagc	ccctttgcca	tagcctgatt	ttggggagga	1920
ggaagggggg	atttgaggga	gaaggggaga	aagcttatgg	ctgggtctgg	tttcttccct	1980
tcccagaggg	tcttactgtt	ccagggtggc	cccaggggcag	gcagggggcca	cactatgcct	2040
gcgcctcggg	aaagggtgacc	cctgccatth	accagcagcc	ctggcatggt	cctgccccac	2100
aggaatagaa	tggagggagc	tccagaaact	ttocatccca	aaggcagtot	ccgtgggtga	2160
agcagactgg	atthttgtct	tgcccctgac	cccttgtccc	tctttgaggg	aggggagcta	2220
tgctaggact	ccaacctcag	ggactcgggt	ggcctgcgct	agcttctttt	gatactgaaa	2280
actthttaagg	tgggaggggtg	gcaagggatg	tgcttaataa	atcaattoca	agcctcacct	2340
g						2341

<210> 67

<211> 2109

<212> DNA

<213> Homo sapiens

<400> 67

gattcggccg	gagctgccag	cggggaggct	gcagccggcg	gttggttacag	ctgctggagc	60
agcagcggcc	cccgctccc	ggaaccgttc	cggggccgtt	gatcttcggc	cccacacgaa	120
cagcagagag	gggcagcagg	atgaatgtgg	gcacagcgca	cagcgagggtg	aaccccaaca	180
cgcggtgat	gaacagccgt	ggcatctggc	tctcctaegt	gctggccatc	ggtctcctcc	240
acatcgtgct	gctgagcatc	ccgtttgtga	gtgtccctgt	cgtctggacc	ctcaccaacc	300
tcattcacaa	catgggcatg	tatatottcc	tgcacacggg	gaaggggaca	ccctttgaga	360
ccccggacca	gggcaaggcg	aggotgctaa	cccactggga	gcagatggat	tatgggggtcc	420
agttcacggc	ctctcggaag	ttcttgacca	tcacacccat	cgtgctgtac	ttcctcacca	480
gcttctacac	taagtacgac	cagatccatt	ttgtgctcaa	caccgtgtcc	ctgatgagcg	540
tgcttatccc	caagctgccc	cagctccacg	gagtccggat	ttttggaatc	aataagtact	600
gagagtgcag	ccccttcccc	tgcccagggt	ggcaggggag	gggtagggtta	aaaggcatgt	660
gctgcaacac	tgaagacaga	aagaagaagc	ctctggacac	tgccagagat	gggggttgag	720
cctctggcct	aatttccccc	ctcgtctccc	ccagtagcca	acttgagta	gottgtagt	780
gggttggggg	aggccccctg	ggctctgacc	tttctggaat	ttttgatct	cttccttttg	840
ctttttgaat	agagactcca	tggagtgggt	catggaatgg	gctgggctcc	tgggctgaac	900
atggaccacg	cagttgcgac	aggaggccag	gggaaaaacc	cctgctcact	tgtttgccct	960
caggcagcca	aagcacttta	acccttgcac	agggagcaga	gggcggtaac	gcttctggat	1020
tgtttcaactg	tgattccctag	gttttttcga	tgccatgcag	tgtgtgcttt	tgtgtatgga	1080
agcaagtgtg	ggatgggtct	ttgoccttct	gggtaggggag	ctgtctaata	caagtcccag	1140
gcttttgcca	gcttctctgc	aaccacccgt	gggtcctggg	tgggagtggg	gagggtcagg	1200
ttggggaaag	atggggtaga	gtgtagatgg	cttggttcca	gaggtgaggg	ggccagggct	1260
gctgcatcc	tggcctgggtg	gaggttgggg	agctgtagga	gagctagtga	gtcgagactt	1320
agaagaatgg	ggccacatag	cagcagagga	ctggtgtaag	ggagggaggg	gtagggacag	1380
aagctagacc	caatctcctt	tgggatgtgg	gcagggaggg	aagcaggctt	ggagggttaa	1440
tttaccacaca	gaatgtgata	gtaatagggg	agggaggctg	ctgtgggttt	aaactcctggg	1500
ttggctgttg	ggtagacagg	tggggaaaag	gcccgtagt	cattgtaagc	acagggtccaa	1560
cttgccctg	actcctgcgg	gggtatgggg	aagctgtgac	agaaacgatg	ggtgctgtgg	1620
tcctctgcag	gccctcaccc	cttaacttcc	tcatgcagac	tggcactggg	cagggcctct	1680
catgtggcag	ccacatgtgg	cgttgtgagg	ccaccccatg	tggggtctgt	ggtgagagtc	1740
ctgtaggatc	cctgctcaag	cagcacagag	gaaggggcaa	gacgtggcct	gtaggcactg	1800
tctcagcctg	cagagaagaa	agtgaggccg	ggagcctgag	cctgggctgg	agccttctcc	1860
cctccccagt	tggactaggg	gcagtgttaa	ttttgaaaag	gtgtgggtcc	ctgtgtcctt	1920
ttccaggggt	ccaaggaac	aggagaggtc	actgggcctg	ttttctccct	cctgaccttg	1980
catctcccac	cctgtgtatc	ataggggaact	ttcaccttaa	aatctttcta	agcaaagtgt	2040
gaataggatt	tttactccct	ttgtacagta	ttctgaggaa	cgcaaataaa	agggcaacat	2100
gtttctggt						2109

<210> 68

<211> 2423

<212> DNA

<213> Homo sapiens

<400> 68

	gagagccgag	ctagcgacga	gcagtcgttg	cggccgcgcg	cgccgcggga	ggtggtggag	60
	gcctagccgg	agccgagagg	tctctgttgc	cogtcccacg	gtcccggcgt	cacccctccg	120
5	gcgcccagtc	cccgtcccgg	aactcccggg	cctgtcctgg	gcccccggtc	tgtgcactcc	180
	gctcgcgcga	gogcccgccc	cgggccgcac	ccgcccggcc	catgaggagg	gacgtgaacg	240
	gagtgacca	gagcagggtt	gagatgttct	caaatagtga	tgaagctgta	atcaataaaa	300
	aacttcccaa	agaactcctg	ttacggatat	tttcttttct	agatgttggt	accctgtgcc	360
	gctgtgctca	ggtctccagg	gcctggaatg	ttctggctct	ggatggcagt	aactggcagc	420
10	gaattgacct	atttgatttc	cagagggata	ttgagggccg	agtagtgag	aatatttcaa	480
	aacgatgtgg	gggcttttta	cgaaagttaa	gtcttcgtgg	atgtcttgga	gtgggagaca	540
	atgcattaag	aacctttgca	caaaactgca	ggaacattga	agtactgaat	ctaaatgggt	600
	gtacaaagac	aacagacgct	acatgtacta	gccttagcaa	gttctgttcc	aaactcaggc	660
	accttgactt	ggcttcctgt	acatcaataa	caaacatgto	tctaaaagct	ctgagtggag	720
	gatgtccact	ggtggagcag	ttgaacattt	cctggtgtga	ccaagtaacc	aaggatggca	780
15	ttcaagcact	agtgaggggc	tgtggggggtc	tcaaggcctt	attcttaaaa	ggctgcaogc	840
	agctagaaga	tgaagctctc	aagtacatag	gtgcacactg	ccctgaactg	gtgactttga	900
	acttgacgac	ttgcttgcaa	atcacagatg	aagggtctcat	tactatatgc	agagggtgcc	960
	ataagttaca	atccctttgt	gcctctggct	gtcccaacat	cacagatgoc	atcctgaatg	1020
	ctctaggtca	gaactgcccc	cggcttagaa	tattggaagt	ggcaagatgt	tctcaattaa	1080
20	cagatgtggg	ctttaccact	ctagccagga	attgoccatga	acttgaaaag	atggacctgg	1140
	aagagtgtgt	tcagataaca	gatagcacat	taatccaaact	ttctatacac	tgtcctcgac	1200
	ttcaagtatt	gagtcctgtc	cactgtgagc	tgatcacaga	tgatggaatt	cgtcacctgg	1260
	ggaatggggc	ctgcgcccac	gaccagctgg	aggtgattga	gctggacaac	tgcccactaa	1320
	tcacagatgc	atccctggag	cacttgaaag	gctgtcatag	ccttgagcgg	atagaactct	1380
	atgactgcca	gcaaatcaca	cgggctggaa	tcaagagact	caggacccat	ttacccaata	1440
25	ttaaagtcca	cgcctacttc	gcacctgtca	ctccaccccc	atcagtaggg	ggcagcagac	1500
	agcgcttctg	cagatgctgc	atcatcctat	gacaatggag	gtggtcaacc	ttggcgaact	1560
	gagtatttaa	tgacacttct	agagctaccg	tggagtctct	ccagtggaag	caaccccagt	1620
	gttctgagca	agggttacaa	agtgagggag	ggcagtgctc	agatccccag	agccacacat	1680
	acatacacat	acacaccctt	accccatcc	actctagctt	tgtgaccatg	ggactgaagt	1740
30	ttgtgatggc	ttttttatca	agtagattgg	taaaatttaa	ccattcctgt	tgaggtgccc	1800
	ataagaaaa	cataggccaa	gatagggagg	ggcattccag	caaaccccgt	gttaatgcta	1860
	ctgtgggttt	taaatttttg	tctaggggtt	tctttgggga	ttttagaaca	gcctctgctg	1920
	tcctccgggg	tcaagaaaag	catggaaaag	caatatatga	tgtaaccagg	gaccagaaag	1980
	aaaatttctt	tgcatcttag	aaatggtaga	cattcattgt	gactaaagag	cttctatgct	2040
	tccttggttc	catgccaaca	tgctgagcat	gtccacaaag	aaggctcgtc	cattcctcct	2100
35	gtgttttagt	atttgcccca	gaggtttcct	aaatggttgc	cttgaaatca	ctgtggtcca	2160
	aatgtaatto	ttacacactc	aaattatcac	tgtctgtagc	acacttgtag	acctgtctta	2220
	cattctctgt	tgctccccc	cacactcttg	ctcagttctg	caactgttca	gtctgtttac	2280
	tcactcaatt	gttacccttt	tgctgttgct	gtgtttacag	tttgcatatt	gaatgattag	2340
	ttgggattac	caaacatttt	ttaaaaagat	attatcaata	aatatTTTTT	taattctaaa	2400
40	ttttaaaaaa	aaaaaaaaaa	aaa				2423

<210> 69

<211> 1841

<212> DNA

<213> Homo sapiens

<400> 69

	agctgggacc	ggaggggtgag	cccgccagag	gcagagacac	acgcgggagag	gaggagagggc	60
	tgagggaggg	aggtggagaa	ggacgggaga	ggcagagaga	ggagacacgc	agagacactc	120
	aggaggggag	agacaccgag	acgcagagac	actcaggagg	ggagagacac	cgagacgcag	180
55	agacacccag	gccggggagc	gcgagggagc	gagggcacaga	cctgggtcag	cgagcgcggg	240
	gggcgagccc	cgagtcccga	gagcctgggg	gcgcgcccag	cccggggcgc	gacctcctc	300

5 cccgtcccgc gccctcccct cggcggggcac ggtattttta tccgtgogog aacagccctc 360
 ctctctctot cgcgcacag cccgcgcct gcgcggggga gccagcaca gaccgcgcgc 420
 gggaccccga gtcgcgacc ccagccccac cgcgccatg gaccccaagg 480
 accgcaagaa gatccagtcc tgggtgcccgc cgcgccctag ccagctcgac ccccgccagg 540
 tggagatgat cggcgcgagg agaccaacgc ctgccatgct gttccggctc tcagagcact 600
 cctcaccaga ggaggaagcc tccccccacc agagagcctc aggagagggg caccatctca 660
 agtcgaagag acccaacccc tgtgcctaca caccaccttc gctgaaagct gtgcagcgca 720
 ttgtctgagtc tcacctgcag totatcagca atttgaatga gaaccaggcc tcagaggagg 780
 aggatgagct gggggagctt cgggagctgg gttatccaag agaggaagat gaggaggaaag 840
 10 aggaggatga tgaagaagag gaagaagaag aggacagcca ggtgaaagtc ctgaagggtca 900
 tcaggcagtc tgctgggcaa aagacaacct gtggccaggg tctggaaggg cctggggagc 960
 gccacccccc tctggatgag tccgagagag atggaggctc tgaggaccaa gtggaagacc 1020
 cagcactaag tgagcctggg gaggaacctc accctctgag cctggcacat 1080
 aggcacccc cctgcctctc ccaggaggaa gtggagggga catcgctgtt cccagaaac 1140
 15 ccactctatc ctcacctgt tttgtgctct tccctcgcc tgctagggtc gcggtctctg 1200
 acttctagaa gactaaggct ggtctgtgtt tgcttgtttg cccacctttg gctgataccc 1260
 agagaacctg ggcaacttgc gctgatgcc caccctgcc agtcattcct ccattcacc 1320
 agcgggagggt gggatgtgag acagcccaca ttggaatac cagaaaacog ggaacaggga 1380
 tttgcccttc acaattctac tcccagatc ctctcccctg gacacaggag acccacaggg 1440
 20 caggacccta agatctgggg aaaggagtc ctgagaacct tgaggtaacc ttagatcctt 1500
 ttctacccac ttctctatgg aggatccaa gtcaccactt ctctcacogg cttctaccag 1560
 ggtccaggac taaggcgttt ttctccatag cctcaacatt ttgggaatot tcccttaac 1620
 acccttgctc ctctgggtg cctggaagat ggactggcag agacctottt gttgcgtttt 1680
 gtgctttgat gccaggaatg ccgcctagtt tatgtccccg gtggggcaca cagcgggggg 1740
 cgcaggttt tccttgctcc ccagctgctc tgcccccttc cccttcttcc ctgactccag 1800
 25 gctgaaccc ctcccgtgct gtaataaatc tttgtaata a 1841

<210> 70

<211> 748

30 <212> DNA

<213> Homo sapiens

35 <400> 70

40 ggcgcgcgat agcgggggagc cggggcagac gtccgtagcg cccctccccg aggaggtcga 60
 gccgggcagtc ggggtccgca tcgtggtgga gtactgtgaa ccctgcggct tcgaggcgac 120
 ctacctggag ctggccagtgc ctgtgaagga gcagtatccg ggcacgcaga tcgagtcgcg 180
 cctcgggggc acagggtgcct ttgagataga gataaatgga cagctggtgt tctccaagct 240
 ggagaatggg ggctttccct atgagaaaga tctcattgag gccatccgaa gagccagtaa 300
 tggagaaacc ctagaaaaga tcaccaacag ccgtccctcc tgcgtcatcc tgtgactgca 360
 caggactctg ggttctgct ctgttctggg gtccaaacct ttggtctcct 420
 tgggagctcc ccctgcctct ttccctact tagctcctta gcaaagagac cctggcctcc 480
 45 actttgcct ttgggtacaa agaaggaata gaagattccg tggccttggg ggcaggagag 540
 agacactctc catgaacact tctccagcca cctcatacc ccttcccagg gtaagtgc 600
 acgaaagccc agtccactct tcgcctcggg aatacctgtc tgatgccaca gattttattt 660
 attctccct aaccagggc aatgtcagct attggcagta aagtggcgct acaaacta 720
 aaaaaaaaa aaaaaaaaa aaaaaaaa 748

50

55

<210> 71

<211> 795

<212> DNA

<213> Homo sapiens

<400> 71

taaggctg	agaagacgac	agaagctaga	cccaatctcc	tttgggatgt	gggcagggag	60
ggaagcaggc	ttggaggggt	aatttaccca	cagaatgtga	tagtaatagg	ggagggaggc	120
tgctgcgggt	ttaactcctg	ggttggctgt	tgggtagaca	ggtggggaaa	aggcccgta	180
gtcattgtaa	gcacaggtcc	aacttggccc	tgactcctgc	gggggtatgg	ggaagctgtg	240
acagaaacga	tgggtgctgt	ggtcctctgc	aggccctcac	cccttaactt	cctcatacag	300
actggcactg	ggcagggcct	ctcatgtggc	agccacatgt	ggcgttgtga	ggccacccca	360
tgtgggtctc	gtggtgagag	tcctgtagga	tcctgtctca	agcagcacag	aggaaggggc	420
aagaogtggc	ctgtaggcac	tgtctcagcc	tgcagagaag	aaagtgaggc	cgggagcctg	480
agcctgggct	ggagccttct	ccctcccca	gttggactag	gggcagtgtt	aattttgaaa	540
aggtgtgggt	ccctgtgtcc	tcttcagggt	gtccaaggga	acaggagagg	tactggggcc	600
tgttttctcc	ctcctgaccc	tgcactctcc	accccggtga	tcatagggaa	ctttcacctt	660
aaaatctttc	taagcaaatg	gtgaatagga	tttttactcc	ctttgtacag	tattctgaga	720
aacgcaata	aaagggcaac	atgtttctgt	taaaaaaaa	aaaaagtacg	caaaaaaaaa	780
aaaaaaaaaa	aaaaa					795

<210> 72

<211> 2356

<212> DNA

<213> Homo sapiens

<400> 72

ggcacgaggg	cggaagtga	ctctagagcg	gtggtgaaac	tggcagttga	cggtcctctg	60
gactagatcc	cgcgaggtag	ccccgaact	atctctctac	gtttctctct	gacccctccg	120
aaatcttcca	gatccgcgta	gtgaggaatc	gtctccaccg	tcattggggg	cggagacctg	180
aatctgaaga	agagctggca	cccgcagacc	ctcaggaatg	tggagaaaat	gtggaaggcc	240
gagcagaagc	atgaggtgta	gcggaagaag	attgaggagc	ttcagcggga	gctgcgagaa	300
gagagagccc	gggaagagat	gcagcgtat	gcggaggatg	ttggggccgt	caagaaaaaa	360
gaagaaaagt	tggactggat	gtaccagggt	cctggtggga	tgggtgaaccg	tgacgagtac	420
ctgotggggc	gccccattga	caaatatgtt	tttgagaaga	tggaggagaa	ggaggcaggc	480
tgtcttctct	aaacaggact	totcccaggc	tcctctcttg	cccatcagg	tgccaattcc	540
cttcttgaca	tggccagcaa	gatccgggag	gacccactct	tcctcatcag	gaagaaggag	600
gaggagaaaa	aacgagaggt	attaaataat	ccagtgaata	tgaagaaaat	caaagaattg	660
ttgcaaatga	gtctggaaaa	aaaggagaag	aagaaaaaga	aggagaagaa	aaagaagcac	720
aagaaacata	agcacagaag	ctcgagtagt	gatcgttcca	gcagcgagga	tgagcacagt	780
gcaggagat	cacagaagaa	gatggcaaat	tcctccctcg	ttttgtccaa	agtccctgga	840
tatggcttac	aggtccggaa	ctctgaccgt	aaccagggtc	ttcagggtcc	tctgacagca	900
gagcaaaaaga	gagggcatgg	gatgaagaac	cattccagat	ccagaagctc	ctccactca	960
cccccaagac	atgccagcaa	gaagagcacc	agggaagcag	ggtcccggga	caggaggtct	1020
cgatccctgg	gcagaaggtc	acgggtcccca	agaccagca	aactgcacaa	ctctaagggtg	1080
aacaggagag	agacaggcca	aactaggagc	ccatcaccta	aaaaagaggt	ctaccaaaagg	1140
cgacatgctc	ccggatacac	cagaaaactc	totgcagagg	aattagagcg	aaaacggcaa	1200
gagatgatgg	aaaacgccaa	atggagggag	gaggagagac	tgaacatcct	caagaggcat	1260
gctaaggatg	aggaacggga	gcagaggcta	gagaagctgg	actcccggga	tgggaagttc	1320

	atocaccgca	tgaagctgga	gagtgcatct	acttccctcc	tggaggatcg	ggtgaagcgg	1380
	aatatctact	ctttacagag	aaacttcggt	gctctggaga	agaactttat	gaaaagatga	1440
5	aaactgtccc	ctctcttatt	ggttttccctg	cattttccag	ggaagctgct	gacctcttaa	1500
	ttctctttat	aagagttcaa	atgacttctt	tcacagatgt	caaaccacca	gtgttcaaag	1560
	tgacctgtct	tcattgagtc	ctgaaacagc	tcacttccct	tgagagctag	tgtgacttgc	1620
	tttgtgggac	actcagtaac	tttgggtttt	gactctttta	cgggtgggca	ctggaccatc	1680
	togggtggag	tgcttgtgcc	actctggaag	gctgttccct	gggttgtga	tgtttatcat	1740
	gccacttcc	tottacctgt	gccaacagac	ctatttcact	gootcagcgt	acaccagacc	1800
10	cttcagaaac	ctctctgggtg	tcaccagat	agattgtgct	tactgagaca	aatgaaogtt	1860
	tacttgattt	agaagataat	gtgacagaat	gatgtcaggt	taggtcaaag	ccaagggagt	1920
	gacagaatct	ggaaaatcaa	acaatacaaa	aagccctaaa	tgaactgtta	actatttgat	1980
	ctttggatgt	aaaattgtaa	tgcgatatgt	tacaaatgta	caatttttac	atgcttttaa	2040
	aaaaggttag	ctttgtgaaa	ataccttggt	tggtcaatga	ctttactggg	taatagaacc	2100
	acattgaacc	ttgatggcaa	gtaatacaat	aaggcaggcc	agctcgtttt	tctctctgaa	2160
15	tctggctgg	ttaggaggag	ctgggttta	togacgagat	ctggagtatc	tattcttttc	2220
	cactccttgc	agtctccaat	gtaggcagtg	taaaggtata	gtaaaatgat	tttaggagtc	2280
	agaaccaa	tgccaatatg	ctccatggct	cctaaaggaa	aataaaatgg	aagtttttaa	2340
	aaaaaaaa	aaaaaa					2356
20	<210> 73						
	<211> 1646						
	<212> DNA						
25	<213> Homo sapiens						
	<400> 73						
30	gtggaatgtc	atcagttaag	gctattttca	tttcttttgt	ggatcttcag	ttgcttcagg	60
	ccatctggat	gtatacatgc	aggtcacagg	gaatatgatg	gcttagcttg	ggttcagagg	120
	cctgacacct	caggctgcc	aatgtggaag	atttaaatac	ttgaaccaat	accctcctcc	180
	caaaaactga	aattggcttc	tgtttctgag	ttggtccagg	cgcaatgttc	agcgtatttg	240
	aggaaatcac	aagaattgta	gttaaggaga	tggtatgctg	aggggatatg	attgccgtta	300
	gaagccttgt	tgatgctgat	agattccgct	gcttccatct	ggtgggggag	aagagaactt	360
35	tctttggatg	ccggcactac	acaacaggcc	tcacctgat	ggacattctg	gacacacatg	420
	gggacaagt	gttagatgaa	ctggattctg	ggctccaagg	tcaaaaggct	gagtttcaaa	480
	ttctggataa	tgtagactca	acgggagagt	tgatagtga	attaccctaa	gaaataacaa	540
	tttcaggcag	ttccagggc	ttccaccatc	agaaaatcaa	gatatcggag	aaccggatat	600
	ccagcagta	tctggctacc	cttgaaaaca	ggaagctgaa	gagggaacta	cccttttcat	660
40	tcogatcaat	taatacgaga	gaaaacctgt	atctgggtgac	agaaactctg	gagacggtaa	720
	aggaggaaac	cctgaaaaagc	gaccggcaat	ataaattttg	gagccagatc	tctcagggcc	780
	atctcagcta	taaacacaag	ggccaaagg	aagtgaacct	ccccccaaat	ogggctcctga	840
	gctatcgagt	aaagcagctt	gtcttcccca	acaaggagac	gatgagaaag	tctttgggtt	900
	oggaggattc	cagaaacatg	aaggagaagt	tggaggacat	ggagagtgtc	ctcaaggacc	960
	tgacagagga	gaagagaaaa	gatgtgctaa	actccctcgc	taagtgcctc	ggcaaggagg	1020
45	atattcggca	ggatctagag	caaagagtat	ctgaggtcct	gatttccggg	gagctacaca	1080
	tggaggaccc	agacaagcct	ctcctaagca	gcctttttta	tgctgctggg	gtcttggttag	1140
	aagcgcgtgc	aaaagccatt	ctggacttcc	tggtatgccct	gctagagctg	tctgaagagc	1200
	agcagtttgt	ggctgaggcc	ctggagaagg	ggacccttcc	tctgttgaa	gaccaggtga	1260
	aatctgtcat	ggagcagaac	tgggatgagc	tggccagcag	tcctcctgac	atggactatg	1320
	accctgaggc	acgaattctc	tgtgcgctgt	atgttgttgt	ctctatcctg	ctggagctgg	1380
50	ctgagggggc	tacctctgtc	tcttcctaac	tacaaaagcc	ctttctcccc	acaagcctct	1440
	gggttttccc	tttaccagtc	tgctcctcact	gccatcgcca	ctaccatcct	gtcaccagtg	1500
	ggacctcttt	aaaacaagca	gccaaaccatt	ctttgatgta	tccatttcgc	tccatgttaa	1560
	catccaaaac	cagcctggat	ttcatacatg	gacttctgat	taaaagtggc	aggttgtgca	1620
55	tgttaaaaaa	aaaaaaaa	aaaaaa				1646

<210> 74

<211> 3340

<212> DNA

<213> Homo sapiens

<400> 74

cgggcgccca	gagacagcgc	cgccctcagat	atcctgctgg	atgacattgt	ccttaccat	60
tctctcttcc	tcccgacgga	gaaatttctg	caggagctac	accagtactt	tggtcgggca	120
ggaggcatgg	agggccctga	agggctgggc	oggaagcaag	cctgtctagc	catgcttctc	180
catttcttgg	acacctacca	ggggctgctt	caagaggaag	agggggccgg	ccacatcatc	240
aaggatctat	acctgctaata	tatgaaggac	gagtcccttt	accagggcct	cogagaggac	300
actctgaggc	tgcaccagct	ggtggagacg	gtggaactaa	agattccaga	ggagaaccag	360
ccacccagca	agcaggtgaa	gccactcttc	cgccacttcc	gcggataga	ctcctgtctg	420
cagaccggg	tgcccttccg	gggtctctgat	gagatcttct	gcgtgtata	catgcctgac	480
cactottatg	tgaccatacg	cagccgcctt	tcagcatctg	tcaggacat	totgggctct	540
gtgacggaga	aacttcaata	ttcagaggag	ccgcggggc	gtgaggattc	cctcatcctg	600
gtagctgtgt	cctcctctgg	agagaaggtc	cttctccagc	ccactgagga	ctgtgttttc	660
accgcactgg	gcataacag	ccacctgttt	gcctgtactc	gggacagcta	tgaggctctg	720
gtgcccctcc	cogaggagat	ccaggctctcc	cctggagaca	cagagatcca	ccgagtggag	780
cctgaggacg	ttgccaacca	cctaactgcc	ttccactggg	agctgttccg	atgtgtgcat	840
gagctggagt	tcgtggacta	cgtgttccac	ggggagcgcg	gccgccggga	gacggccaac	900
ttgtgagctg	tgctgcagcg	ctgcagcgag	gtcacgcact	gggtggccac	cgaagtgtctg	960
ctctgcgagg	cccogggcaa	gcgcgcgcag	ctgctcaaga	agttcatcaa	gatcgcggcc	1020
ctctgcaagc	agaaccagga	cctgctgtct	ttctacgcgc	tggtcatggg	gctggacaac	1080
gccgctgtca	gccgccttcg	actcacctgg	gagaagctgc	cagggaaatt	caagaacttg	1140
tttgcgaaat	ttgagaacct	gacggacccc	tcagggaacc	acaaaagcta	cogagaagtg	1200
atctccaaaa	tgaagccccc	tgtgattccc	ttcgtgcctc	tgatcctcaa	agacctgact	1260
ttcctgcacg	aagggaagtaa	gacccttgta	gatggtttgg	tgaacatcga	gaagctgcat	1320
tcagtggccg	aaaaagttag	gacaatccgc	aaataccgga	gocggccctt	ttgcctggac	1380
atggaggcat	cccccaatca	cctgcagacc	aaggcctatg	tgccgcagtt	tcaggctcatc	1440
gacaaccaga	acctcctctt	cgagctctcc	tacaagctgg	aggcaaacag	tcagttagag	1500
tgaggagctcc	agtcagaccc	gccagatcct	tgggcacctg	gcactcaagc	actttgcacg	1560
atgtotcaac	caacatctga	catctttccc	gtggagcaac	ttcctgctcc	acgggaaaga	1620
ggctgcctga	tttacccttg	gaccataagc	ctgtttcatc	ctgctgaagt	cccctcccca	1680
ttgtcctctc	aagccaaaac	tacactttgc	tggttctctg	cccctctgag	aaaggggata	1740
gaaagctcct	tcctctatgt	cctcccatcg	agatctgttc	tggggatgga	gcttccaact	1800
tcctotttga	gcaggaaaga	atgctgctca	cccttctgtc	ttgcagagtg	ggattgtggg	1860
agggattggc	agccttcttc	tcacccacct	gtccagcttc	ttcctggtca	gggctgggac	1920
ccccaggaat	attatgttgc	cgtgtgtgtg	tgtgtgtgtg	tgtgtgtgtg	tgtgtgtgtg	1980
tgtgtcttct	tttagggagc	aggagtgcac	ctggtaattg	agggtgatg	ttgtgtgtgc	2040
tggggagggg	tccttctgtt	tggtgctacc	cttgtctact	ctgcccctgg	atgggtgcggg	2100
gtgctttctc	cacccccaca	ctccctgtct	agctcctcgt	gctgccctgc	atgccaggc	2160
ttgtgagcca	aggtgctttt	tggggcaggg	agtagcagca	ggtagggagg	gttaccatc	2220
agcccttgca	agtcccccac	tcaggcctct	ggaaggcca	gggatgggct	ctgatgagag	2280
ggtaaaagat	gctcagggga	acacaggcct	cagctgccta	gaggaccctc	cccctgcctt	2340
gcagtgggct	cgggtagagc	agtatcagga	gctagggttg	totgctgccc	acactcctgc	2400
tttttgggat	atctaactgc	taaggaggga	gttgacatcc	cccttctggc	tcattgtgtct	2460
gacaccaaca	acatgggtctc	cgtccctctc	tcttagactc	tcctttgttc	ctccccatag	2520
agctgggggtg	gggtggatcc	ctatactggg	gcaggcagcc	ccaaagtggg	ggagggggat	2580
ggcagagact	gtaaaggcgc	cactggactc	tggaaggcc	tttattacct	ttactccctc	2640
cctctcccat	caccagcctc	aaggcctgag	gggtgcaggg	gctcctggca	gctactgggt	2700
gaggtttctc	ggcagagact	caccttctt	ctctgcacca	ctcttccct	tttgaagaga	2760
cagcaacagc	cgtagcaaaa	gcagctgctg	ctcctgctat	gagggtgtat	atatTTTTTTA	2820
cccaaagctc	tgggaattgta	catttatTTTt	ttaaaactca	aagagggaaa	gagccttgta	2880

	tcatatgtga	acattgtatc	ataggtaatg	ttgtacagac	ccttttatac	agtgatctgt	2940
	cttgttccctg	cagcaaaaat	ccttatatgga	cataggaggt	gctgtgtccc	atgccttctt	3000
	gccttgacag	tgtcccatgg	gcccccttct	gctccctgcc	ccctccctgc	tactgtctgat	3060
5	gcactgtcct	ctccctgcag	cccctggctt	cccagccttc	ctcctgaccc	cttccaacag	3120
	ccttggaaact	ccagctgcc	ccaccctctg	ggtcggacac	tgggacccac	tggcccagtc	3180
	ttggctgctg	cttaccctta	goccttgatgc	ctgcccaggg	acccccagcc	ccctcccgtt	3240
	gccttgacgc	tttaacagag	tgaacatgt	gtattgtaca	ggcggggttg	tcattgcaga	3300
	aacogotggg	tggagaagaa	gccgataaag	tctatgaatc			3340
10	<210> 75						
	<211> 4005						
	<212> DNA						
15	<213> Homo sapiens						
	<400> 75						
20	gggcaacagt	ctgcccacct	gtggacacca	gatoctggga	gctcctgggt	agcaagttag	60
	atctctggga	tgtcagtgag	gctggttgaa	gaccagaggt	aaactgcaga	ggtcaccacc	120
	cccaocatgt	cccaggtgat	gtccagccca	ctgctggcag	gaggccatgc	tgtcagcttg	180
	gcgccttgtg	atgagcccag	gaggaccctg	cacccagcac	ccagccccag	cctgccaccc	240
	cagtgttctt	actacaccac	ggaaggctgg	ggagcccagg	ccctgatggc	cccogtgccc	300
25	tgcattgggc	cccctggccg	actccagcaa	gccccacagg	tggaggccaa	agccacctgc	360
	ttcctgcctg	cccctgggtg	gaaggccttg	gggaccccag	aggaccttga	ctcctacatt	420
	gactttctac	tggagagcct	caatcagatg	atcctggaac	tggaccccac	cttcagctg	480
	cttccccag	ggactggggg	ctcccaggct	gagctggccc	agagcaccat	gtcaatgaga	540
	aagaaggagg	aatctgaagc	cttggacata	aagtacatcg	aggtgacctc	cgccagatca	600
	aggtgccacg	attggcccca	gcactgctcc	agcccctctg	tcaccccggc	cttoggtccc	660
30	cctgcagtg	gtggcctcct	cctttccaga	gaggtccccc	gagagacacg	aagcagctg	720
	gagagcctca	tcttctctgg	gaaccagggc	agggggcacc	agcgccctct	gccccctca	780
	gaggtctctt	cccctcgacc	cccaaattcc	cccagcatct	caatcccttg	catggggagc	840
	aaggcctcga	gcccccatgg	tttgggctcc	ccgctgggtg	cttctccaag	actggagaag	900
	cggtctggag	gcctggcccc	acagcggggc	agcaggatct	ctgtgctgtc	agccagccca	960
35	gtgtctgatg	tcagctatat	gtttggaagc	agccagtccc	tcctgcactc	cagcaactcc	1020
	agccatcagt	catcttccag	atccttgga	agtcacagca	actcttctct	cagcctccac	1080
	agccttggct	cagtgtccct	gtgtacaaga	ccagtgact	tcaggctcc	cagaaacccc	1140
	accctaacca	tgggccaacc	cagaacaccc	caactctcac	caactggccaa	agaacatgcc	1200
	agcatctgcc	ccccatccat	caccaactcc	atggttgaca	taccatttgt	gctgatcaac	1260
40	ggctgcccag	aaccagggtc	ttctccaccc	cagcggaccc	caggacacca	gaactccgtt	1320
	caacctggag	ctgcttctcc	cagcaacccc	tgtccagcca	ccaggagcaa	cagccagacc	1380
	ctgtcagatg	ccccctttac	cacatgcccc	gaggttcccg	ccaggacat	gcagcccacc	1440
	atgaagtctg	tgatggacac	atctaaatac	tggtttaagc	caaacatcac	ccgagagcaa	1500
	gcaatcgagc	tgttgaggaa	ggaggagcca	ggggcttttg	tcataaggga	cagctcttca	1560
	taccgaggct	ccttcggcct	ggccctgaag	gtgcaggagg	ttcccgcgtc	tgtcagaat	1620
45	cgaccagggt	aggacagcaa	tgacctcatc	cgacacttcc	tcctcgagtc	gtctgcaaaa	1680
	ggagtgcac	tcaaaggagc	agatgaggag	ccctactttg	ggagcctctc	tgccttctgt	1740
	tgcagcatt	ccatcatggc	cctggccctg	ccctgcaaac	tcaccatccc	acagagagaa	1800
	ctgggagggt	cagatggggc	ctcggactct	acagacagcc	cagcctcctg	ccagaagaaa	1860
	tctgcgggct	gccacaccct	gtacctgagc	tcagttagcg	tggagaccct	gactggagcc	1920
	ctggccgtgc	agaaagccat	ctccaccacc	tttgagaggg	acatcctccc	cacgcccacc	1980
50	gtgggtccact	togaagtcac	agagcagggc	atcactctga	ctgatgtcca	gaggaagggt	2040
	tttttccggc	gccattaccc	actcaccacc	ctccgcttct	gtggatgtga	ccctgagcaa	2100
	cggaagtggc	agaagtactg	caaaccctcc	tggatctttg	ggtttgtggc	caagagccag	2160
	acagagcctc	aggagaacgt	atgccacctc	tttgcggagt	atgacatggg	ccagccagcc	2220
	tgcaggtca	tggcctgggt	gactgctctg	ctgcaggacg	cagaaaggat	gtagggggaga	2280
55	gactgcctgt	gcacctaac	aacacctcca	ggggctcgt	aaggagcccc	cctccacccc	2340

ctgaatgggt gtggccttggt gccatattga cagaccaatc tatgggacta gggggattgg 2400
 catcaagttg acacccttga acctgctatg gccttcagca gtcaccatca tccagacccc 2460
 cggggcctca gtttccctcaa tcatagaaga agaccaatag acaagatcag ctgttcttag 2520
 atgctgggtg gcatttgaaac atgctcctcc atgattctga agcatgcaca cctctgaaga 2580
 cccctgcatg aaaataacct ccaaggaccc tctgacccca tgcacctggg cccctgcccac 2640
 acaacagtct gagcaagaga cctgcagccc ctgtttcgtg gcagacagca ggtgcctggc 2700
 ggtgaccacac ggggctcctg gcttgacgtc ggtgatggtc aagaactgac tacaaaaacag 2760
 gaatggatag actctatttc cttccatata tgttcctctg ttccttttcc cactttctgg 2820
 gtggcctttt ggggtccaccc agccaggatg ctgcaggcca agctgggtgt ggtatttagg 2880
 gcagctcagc aggggggaact tgtcccatat gtcagaggag acccagctgt cctgcacccc 2940
 cttgcagatg agtatcaccc catcttttct tccacttggt tttttatttt tatttttttt 3000
 gagacagagt ctcactgtca cccaggctga actgcagtgg tgtgatctag gctcactgca 3060
 acctccacct cccaggttca agcaattatc ctgcctcagg ctcccagta gctgggatta 3120
 caggcatgtg caactcaccc agctaatttt gtatttttag tagagacagg gtttcacccat 3180
 gttggccagg ctggtcttga actcctgacc gcaggtaate cacctgcttc ggccctccaa 3240
 agtgctggga ttacaggcgc aagccaccca gccagcttc tttccattcc ttgataggcg 3300
 agtattccaa agctggtatc gtagctgccc taatgttgca tattaggcgg cgggggcaga 3360
 gataagggcc atctctctgt gattctgcct cagctcctgt cttgctgagc cctcccccaa 3420
 cccacgctcc aacacacaca cacacacaca cacacacaca cacacacaca cacacacaca 3480
 cagccccctc tactgctatg tggcttcaac cagcctcaca gccacacggg ggaagcagag 3540
 agtcaagaat gcaaagaggc cgcttcccta agaggcttgg aggagctggg ctctatocca 3600
 caccaccccc caccaccccc ccaccagcc tccagaagct ggaaccattt ctcccagag 3660
 cctgagttcc taaggaaacc accctacogg ggtggaaggg agggtcaggg aagaaaccca 3720
 ctcttgctct acgaggagca agtgccctgcc cctcccagc agccagccct gccaaagttg 3780
 cattatcttt ggccaagget gggcctgacg gttatgatit cagccctggg cctgcaggag 3840
 aggtgagat cagccccccc agccagtggg cgagcactgc cccgcgcgca aagtctgcag 3900
 aatgtgagat gaggttctca aggtcacagg ccccagtccc agcctggggg ctggcagagg 3960
 ccccatata ctctgtctaca gctcctatca tgaaaaataa aatgt 4005

<210> 76

<211> 1093

<212> PRT

<213> Homo sapiens

<400> 76

Met Lys Glu Met Val Gly Gly Cys Cys Val Cys Ser Asp Glu Arg Gly
 1 5 10 15
 Trp Ala Glu Asn Pro Leu Val Tyr Cys Asp Gly His Ala Cys Ser Val
 20 25 30
 Ala Val His Gln Ala Cys Tyr Gly Ile Val Gln Val Pro Thr Gly Pro
 35 40 45
 Trp Phe Cys Arg Lys Cys Glu Ser Gln Glu Arg Ala Ala Arg Val Arg
 50 55 60
 Cys Glu Leu Cys Pro His Lys Asp Gly Ala Leu Lys Arg Thr Asp Asn
 65 70 75 80
 Gly Gly Trp Ala His Val Val Cys Ala Leu Tyr Ile Pro Glu Val Gln
 85 90 95
 Phe Ala Asn Val Leu Thr Met Glu Pro Ile Val Leu Gln Tyr Val Pro
 100 105 110
 His Asp Arg Phe Asn Lys Thr Cys Tyr Ile Cys Glu Glu Thr Gly Arg
 115 120 125
 Glu Ser Lys Ala Ala Ser Gly Ala Cys Met Thr Cys Asn Arg His Gly
 130 135 140

	Cys	Arg	Gln	Ala	Phe	His	Val	Thr	Cys	Ala	Gln	Met	Ala	Gly	Leu	Leu
	145					150					155					160
5	Cys	Glu	Glu	Glu	Val	Leu	Glu	Val	Asp	Asn	Val	Lys	Tyr	Cys	Gly	Tyr
					165				170						175	
	Cys	Lys	Tyr	His	Phe	Ser	Lys	Met	Lys	Thr	Ser	Arg	His	Ser	Ser	Gly
				180					185					190		
	Gly	Gly	Gly	Gly	Gly	Ala	Gly	Gly	Gly	Gly	Ser	Met	Gly	Gly	Gly	
				195				200				205				
10	Gly	Ser	Gly	Phe	Ile	Ser	Gly	Arg	Arg	Ser	Arg	Ser	Ala	Ser	Pro	Ser
	210						215					220				
	Thr	Gln	Gln	Glu	Lys	His	Pro	Thr	His	His	Glu	Arg	Gly	Gln	Lys	Lys
	225					230					235					240
	Ser	Arg	Lys	Asp	Lys	Glu	Arg	Leu	Lys	Gln	Lys	His	Lys	Lys	Arg	Pro
					245					250					255	
15	Glu	Ser	Pro	Pro	Ser	Ile	Leu	Thr	Pro	Pro	Val	Val	Pro	Thr	Ala	Asp
				260					265					270		
	Lys	Val	Ser	Ser	Ser	Ala	Ser	Ser	Ser	Ser	His	His	Glu	Ala	Ser	Thr
			275					280					285			
	Gln	Glu	Thr	Ser	Glu	Ser	Ser	Arg	Glu	Ser	Lys	Gly	Lys	Lys	Ser	Ser
	290						295					300				
20	Ser	His	Ser	Leu	Ser	His	Lys	Gly	Lys	Lys	Leu	Ser	Ser	Gly	Lys	Gly
	305					310					315					320
	Val	Ser	Ser	Phe	Thr	Ser	Ala	Ser	Ser	Ser	Ser	Ser	Ser	Ser	Ser	Ser
				325					330							335
	Ser	Ser	Gly	Gly	Pro	Phe	Gln	Pro	Ala	Val	Ser	Ser	Leu	Gln	Ser	Ser
25				340					345					350		
	Pro	Asp	Phe	Ser	Ala	Phe	Pro	Lys	Leu	Glu	Gln	Pro	Glu	Glu	Asp	Lys
			355					360					365			
	Tyr	Ser	Lys	Pro	Thr	Ala	Pro	Ala	Pro	Ser	Ala	Pro	Pro	Ser	Pro	Ser
	370						375					380				
30	Ala	Pro	Glu	Pro	Pro	Lys	Ala	Asp	Leu	Phe	Glu	Gln	Lys	Val	Val	Phe
	385					390					395					400
	Ser	Gly	Phe	Gly	Pro	Ile	Met	Arg	Phe	Ser	Thr	Thr	Thr	Ser	Ser	Ser
					405					410					415	
	Gly	Arg	Ala	Arg	Ala	Pro	Ser	Pro	Gly	Asp	Tyr	Lys	Ser	Pro	His	Val
				420					425					430		
35	Thr	Gly	Ser	Gly	Ala	Ser	Ala	Gly	Thr	His	Lys	Arg	Met	Pro	Ala	Leu
				435				440					445			
	Ser	Ala	Thr	Pro	Val	Pro	Ala	Asp	Glu	Thr	Pro	Glu	Thr	Gly	Leu	Lys
				450			455					460				
	Glu	Lys	Lys	His	Lys	Ala	Ser	Lys	Arg	Ser	Arg	His	Gly	Pro	Gly	Arg
	465					470					475					480
40	Pro	Lys	Gly	Ser	Arg	Asn	Lys	Glu	Gly	Thr	Gly	Gly	Pro	Ala	Ala	Pro
					485					490					495	
	Ser	Leu	Pro	Ser	Ala	Gln	Leu	Ala	Gly	Phe	Thr	Ala	Thr	Ala	Ala	Ser
				500					505					510		
	Pro	Phe	Ser	Gly	Gly	Ser	Leu	Val	Ser	Ser	Gly	Leu	Gly	Gly	Leu	Ser
45				515				520					525			
	Ser	Arg	Thr	Phe	Gly	Pro	Ser	Gly	Ser	Leu	Pro	Ser	Leu	Ser	Leu	Glu
				530			535					540				
	Ser	Pro	Leu	Leu	Gly	Ala	Gly	Ile	Tyr	Thr	Ser	Asn	Lys	Asp	Pro	Ile
	545					550					555					560
50	Ser	His	Ser	Gly	Gly	Met	Leu	Arg	Ala	Val	Cys	Ser	Thr	Pro	Leu	Ser
					565					570					575	
	Ser	Ser	Leu	Leu	Gly	Pro	Pro	Gly	Thr	Ser	Ala	Leu	Pro	Arg	Leu	Ser
				580					585					590		
	Arg	Ser	Pro	Phe	Thr	Ser	Thr	Leu	Pro	Ser	Ser	Ser	Ala	Ser	Ile	Ser
				595				600						605		

Thr Thr Gln Val Phe Ser Leu Ala Gly Ser Thr Phe Ser Leu Pro Ser
 610 615 620
 Thr His Ile Phe Gly Thr Pro Met Gly Ala Val Asn Pro Leu Leu Ser
 5 625 630 635 640
 Gln Ala Glu Ser Ser His Thr Glu Pro Asp Leu Glu Asp Cys Ser Phe
 645 650 655
 Arg Cys Arg Gly Thr Ser Pro Gln Glu Ser Leu Ser Ser Met Ser Pro
 660 665 670
 Ile Ser Ser Leu Pro Ala Leu Phe Asp Gln Thr Ala Ser Ala Pro Cys
 10 675 680 685
 Gly Gly Gly Gln Leu Asp Pro Ala Ala Pro Gly Thr Thr Asn Met Glu
 690 695 700
 Gln Leu Leu Glu Lys Gln Gly Asp Gly Glu Ala Gly Val Asn Ile Val
 705 710 715 720
 15 Glu Met Leu Lys Ala Leu His Ala Leu Gln Lys Glu Asn Gln Arg Leu
 725 730 735
 Gln Glu Gln Ile Leu Ser Leu Thr Ala Lys Lys Glu Arg Leu Gln Ile
 740 745 750
 Leu Asn Val Gln Leu Ser Val Pro Phe Pro Ala Leu Pro Ala Ala Leu
 755 760 765
 20 Pro Ala Ala Asn Gly Pro Val Pro Gly Pro Tyr Gly Leu Pro Pro Gln
 770 775 780
 Ala Gly Ser Ser Asp Ser Leu Ser Thr Ser Lys Ser Pro Pro Gly Lys
 785 790 795 800
 Ser Ser Leu Gly Leu Asp Asn Ser Leu Ser Thr Ser Ser Glu Asp Pro
 25 805 810 815
 His Ser Gly Cys Pro Ser Arg Ser Ser Ser Ser Leu Ser Phe His Ser
 820 825 830
 Thr Pro Pro Pro Leu Pro Leu Leu Gln Gln Ser Pro Ala Thr Leu Pro
 835 840 845
 30 Leu Ala Leu Pro Gly Ala Pro Ala Pro Leu Pro Pro Gln Pro Gln Asn
 850 855 860
 Gly Leu Gly Arg Ala Pro Gly Ala Ala Gly Leu Gly Ala Met Pro Met
 865 870 875 880
 Ala Glu Gly Leu Leu Gly Gly Leu Ala Gly Ser Gly Gly Leu Pro Leu
 885 890 895
 35 Asn Gly Leu Leu Gly Gly Leu Asn Gly Ala Ala Ala Pro Asn Pro Ala
 900 905 910
 Ser Leu Ser Gln Ala Gly Gly Ala Pro Thr Leu Gln Leu Pro Gly Cys
 915 920 925
 Leu Asn Ser Leu Thr Glu Gln Gln Arg His Leu Leu Gln Gln Gln Glu
 930 935 940
 40 Gln Gln Leu Gln Gln Leu Gln Gln Leu Leu Ala Ser Pro Gln Leu Thr
 945 950 955 960
 Pro Glu His Gln Thr Val Val Tyr Gln Met Ile Gln Gln Ile Gln Gln
 965 970 975
 Lys Arg Glu Leu Gln Arg Leu Gln Met Ala Gly Gly Ser Gln Leu Pro
 980 985 990
 45 Met Ala Ser Leu Leu Ala Gly Ser Ser Thr Pro Leu Leu Ser Ala Gly
 995 1000 1005
 Thr Pro Gly Leu Leu Pro Thr Ala Ser Ala Pro Pro Leu Leu Pro
 1010 1015 1020
 50 Ala Gly Ala Leu Val Ala Pro Ser Leu Gly Asn Asn Thr Ser Leu
 1025 1030 1035
 Met Ala Ala Ala Ala Ala Ala Ala Ala Val Ala Ala Ala Gly Gly
 1040 1045 1050
 Pro Pro Val Leu Thr Ala Gln Thr Asn Pro Phe Leu Ser Leu Ser
 1055 1060 1065

55

Gly Ala Glu Gly Ser Gly Gly Gly Pro Lys Gly Gly Thr Ala Asp
 1070 1075 1080
 Lys Gly Ala Ser Ala Asn Gln Glu Lys Gly
 1085 1090

<210> 77

<211> 344

<212> PRT

<213> Homo sapiens

<400> 77

Met His Arg Thr Thr Arg Ile Lys Ile Thr Glu Leu Asn Pro His Leu
 1 5 10 15
 Met Cys Ala Leu Cys Gly Gly Tyr Phe Ile Asp Ala Thr Thr Ile Val
 20 25 30
 Glu Cys Leu His Ser Phe Cys Lys Thr Cys Ile Val Arg Tyr Leu Glu
 35 40 45
 Thr Asn Lys Tyr Cys Pro Met Cys Asp Val Gln Val His Lys Thr Arg
 50 55 60
 Pro Leu Leu Ser Ile Arg Ser Asp Lys Thr Leu Gln Asp Ile Val Tyr
 65 70 75 80
 Lys Leu Val Pro Gly Leu Phe Lys Asp Glu Met Lys Arg Arg Arg Asp
 85 90 95
 Phe Tyr Ala Ala Tyr Pro Leu Thr Glu Val Pro Asn Gly Ser Asn Glu
 100 105 110
 Asp Arg Gly Glu Val Leu Glu Gln Glu Lys Gly Ala Leu Ser Asp Asp
 115 120 125
 Glu Ile Val Ser Leu Ser Ile Glu Phe Tyr Glu Gly Ala Arg Asp Arg
 130 135 140
 Asp Glu Lys Lys Gly Pro Leu Glu Asn Gly Asp Gly Asp Lys Glu Lys
 145 150 155 160
 Thr Gly Val Arg Phe Leu Arg Cys Pro Ala Ala Met Thr Val Met His
 165 170 175
 Leu Ala Lys Phe Leu Arg Asn Lys Met Asp Val Pro Ser Lys Tyr Lys
 180 185 190
 Val Glu Val Leu Tyr Glu Asp Glu Pro Leu Lys Glu Tyr Tyr Thr Leu
 195 200 205
 Met Asp Ile Ala Tyr Ile Tyr Pro Trp Arg Arg Asn Gly Pro Leu Pro
 210 215 220
 Leu Lys Tyr Arg Val Gln Pro Ala Cys Lys Arg Leu Thr Leu Ala Thr
 225 230 235 240
 Val Pro Thr Pro Ser Glu Gly Thr Asn Thr Ser Gly Ala Ser Glu Cys
 245 250 255
 Glu Ser Val Ser Asp Lys Ala Pro Ser Pro Ala Thr Leu Pro Ala Thr
 260 265 270
 Ser Ser Ser Leu Pro Ser Pro Ala Thr Pro Ser His Gly Ser Pro Ser
 275 280 285
 Ser His Gly Pro Pro Ala Thr His Pro Thr Ser Pro Thr Pro Pro Ser
 290 295 300
 Thr Ala Ser Gly Ala Thr Thr Ala Ala Asn Gly Gly Ser Leu Asn Cys
 305 310 315 320
 Leu Gln Thr Pro Ser Ser Thr Ser Arg Gly Arg Lys Met Thr Val Asn
 325 330 335

Gly Ala Pro Val Pro Pro Leu Thr
340

<210> 78

<211> 416

<212> PRT

<213> Homo sapiens

<400> 78

Met	Ser	Ser	Asn	Cys	Thr	Ser	Thr	Thr	Ala	Val	Ala	Val	Ala	Pro	Leu
1			5						10					15	
Ser	Ala	Ser	Lys	Thr	Lys	Thr	Lys	Lys	Lys	His	Phe	Val	Cys	Gln	Lys
			20					25					30		
Val	Lys	Leu	Phe	Arg	Ala	Ser	Glu	Pro	Ile	Leu	Ser	Val	Leu	Met	Trp
		35					40					45			
Gly	Val	Asn	His	Thr	Ile	Asn	Glu	Leu	Ser	Asn	Val	Pro	Val	Pro	Val
	50					55					60				
Met	Leu	Met	Pro	Asp	Asp	Phe	Lys	Ala	Tyr	Ser	Lys	Ile	Lys	Val	Asp
	65				70					75					80
Asn	His	Leu	Phe	Asn	Lys	Glu	Asn	Leu	Pro	Ser	Arg	Phe	Lys	Phe	Lys
				85					90					95	
Glu	Tyr	Cys	Pro	Met	Val	Phe	Arg	Asn	Leu	Arg	Glu	Arg	Phe	Gly	Ile
			100					105					110		
Asp	Asp	Gln	Asp	Tyr	Gln	Asn	Ser	Val	Thr	Arg	Ser	Ala	Pro	Ile	Asn
		115					120					125			
Ser	Asp	Ser	Gln	Gly	Arg	Cys	Gly	Thr	Arg	Phe	Leu	Thr	Thr	Tyr	Asp
	130					135					140				
Arg	Arg	Phe	Val	Ile	Lys	Thr	Val	Ser	Ser	Glu	Asp	Val	Ala	Glu	Met
	145				150					155					160
His	Asn	Ile	Leu	Lys	Lys	Tyr	His	Gln	Phe	Ile	Val	Glu	Cys	His	Gly
				165					170					175	
Asn	Thr	Leu	Leu	Pro	Gln	Phe	Leu	Gly	Met	Tyr	Arg	Leu	Thr	Val	Asp
		180						185					190		
Gly	Val	Glu	Thr	Tyr	Met	Val	Val	Thr	Arg	Asn	Val	Phe	Ser	His	Arg
	195						200					205			
Leu	Thr	Val	His	Arg	Lys	Tyr	Asp	Leu	Lys	Gly	Ser	Thr	Val	Ala	Arg
	210					215					220				
Glu	Ala	Ser	Asp	Lys	Glu	Lys	Ala	Lys	Asp	Leu	Pro	Thr	Phe	Lys	Asp
	225				230					235					240
Asn	Asp	Phe	Leu	Asn	Glu	Gly	Gln	Lys	Leu	His	Val	Gly	Glu	Glu	Ser
			245						250					255	
Lys	Lys	Asn	Phe	Leu	Glu	Lys	Leu	Lys	Arg	Asp	Val	Glu	Phe	Leu	Ala
			260						265				270		
Gln	Leu	Lys	Ile	Met	Asp	Tyr	Ser	Leu	Leu	Val	Gly	Ile	His	Asp	Val
	275						280					285			
Asp	Arg	Ala	Glu	Gln	Glu	Glu	Met	Glu	Val	Glu	Glu	Arg	Ala	Glu	Asp
	290					295					300				
Glu	Glu	Cys	Glu	Asn	Asp	Gly	Val	Gly	Gly	Asn	Leu	Leu	Cys	Ser	Tyr
	305				310					315					320
Gly	Thr	Pro	Pro	Asp	Ser	Pro	Gly	Asn	Leu	Leu	Ser	Phe	Pro	Arg	Phe
				325					330					335	
Phe	Gly	Pro	Gly	Glu	Phe	Asp	Pro	Ser	Val	Asp	Val	Tyr	Ala	Met	Lys
			340					345					350		

Ser His Glu Ser Ser Pro Lys Lys Glu Val Tyr Phe Met Ala Ile Ile
 355 360 365
 Asp Ile Leu Thr Pro Tyr Asp Thr Lys Lys Lys Ala Ala His Ala Ala
 370 375 380
 Lys Thr Val Lys His Gly Ala Gly Ala Glu Ile Ser Thr Val Asn Pro
 385 390 395 400
 Glu Gln Tyr Ser Lys Arg Phe Asn Glu Phe Met Ser Asn Ile Leu Thr
 405 410 415

<210> 79

<211> 500

<212> PRT

<213> Homo sapiens

<400> 79

Met Arg Gly Glu Leu Trp Leu Leu Val Leu Val Leu Arg Glu Ala Ala
 1 5 10 15
 Arg Ala Leu Ser Pro Gln Pro Gly Ala Gly His Asp Glu Gly Pro Gly
 20 25 30
 Ser Gly Trp Ala Ala Lys Gly Thr Val Arg Gly Trp Asn Arg Arg Ala
 35 40 45
 Arg Glu Ser Pro Gly His Val Ser Glu Pro Asp Arg Thr Gln Leu Ser
 50 55 60
 Gln Asp Leu Gly Gly Gly Thr Leu Ala Met Asp Thr Leu Pro Asp Asn
 65 70 75 80
 Arg Thr Arg Val Val Glu Asp Asn His Ser Tyr Tyr Val Ser Arg Leu
 85 90 95
 Tyr Gly Pro Ser Glu Pro His Ser Arg Glu Leu Trp Val Asp Val Ala
 100 105 110
 Glu Ala Asn Arg Ser Gln Val Lys Ile His Thr Ile Leu Ser Asn Thr
 115 120 125
 His Arg Gln Ala Ser Arg Val Val Leu Ser Phe Asp Phe Pro Phe Tyr
 130 135 140
 Gly His Pro Leu Arg Gln Ile Thr Ile Ala Thr Gly Gly Phe Ile Phe
 145 150 155 160
 Met Gly Asp Val Ile His Arg Met Leu Thr Ala Thr Gln Tyr Val Ala
 165 170 175
 Pro Leu Met Ala Asn Phe Asn Pro Gly Tyr Ser Asp Asn Ser Thr Val
 180 185 190
 Val Tyr Phe Asp Asn Gly Thr Val Phe Val Val Gln Trp Asp His Val
 195 200 205
 Tyr Leu Gln Gly Trp Glu Asp Lys Gly Ser Phe Thr Phe Gln Ala Ala
 210 215 220
 Leu His His Asp Gly Arg Ile Val Phe Ala Tyr Lys Glu Ile Pro Met
 225 230 235 240
 Ser Val Pro Glu Ile Ser Ser Ser Gln His Pro Val Lys Thr Gly Leu
 245 250 255
 Ser Asp Ala Phe Met Ile Leu Asn Pro Ser Pro Asp Val Pro Glu Ser
 260 265 270
 Arg Arg Arg Ser Ile Phe Glu Tyr His Arg Ile Glu Leu Asp Pro Ser
 275 280 285
 Lys Val Thr Ser Met Ser Ala Val Glu Phe Thr Pro Leu Pro Thr Cys
 290 295 300

5 Leu Gln His Arg Ser Cys Asp Ala Cys Met Ser Ser Asp Leu Thr Phe
 305 310 315 320
 Asn Cys Ser Trp Cys His Val Leu Gln Arg Cys Ser Ser Gly Phe Asp
 325 330 335
 Arg Tyr Arg Gln Glu Trp Met Asp Tyr Gly Cys Ala Gln Glu Ala Glu
 340 345 350
 Gly Arg Met Cys Glu Asp Phe Gln Asp Glu Asp His Asp Ser Ala Ser
 355 360 365
 10 Pro Asp Thr Ser Phe Ser Pro Tyr Asp Gly Asp Leu Thr Thr Thr Ser
 370 375 380
 Ser Ser Leu Phe Ile Asp Ser Leu Thr Thr Glu Asp Asp Thr Lys Leu
 385 390 395 400
 Asn Pro Tyr Ala Gly Gly Asp Gly Leu Gln Asn Asn Leu Ser Pro Lys
 405 410 415
 15 Thr Lys Gly Thr Pro Val His Leu Gly Thr Ile Val Gly Ile Val Leu
 420 425 430
 Ala Val Leu Leu Val Ala Ala Ile Ile Leu Ala Gly Ile Tyr Ile Asn
 435 440 445
 Gly His Pro Thr Ser Asn Ala Ala Leu Phe Phe Ile Glu Arg Arg Pro
 450 455 460
 20 His His Trp Pro Ala Met Lys Phe Arg Ser His Pro Asp His Ser Thr
 465 470 475 480
 Tyr Ala Glu Val Glu Pro Ser Gly His Glu Lys Glu Gly Phe Met Glu
 485 490 495
 Ala Glu Gln Cys
 500
 25
 <210> 80
 <211> 509
 30 <212> PRT
 <213> Homo sapiens
 35
 <400> 80
 Met Glu Asp Ile Gln Thr Asn Ala Glu Leu Lys Ser Thr Gln Glu Gln
 1 5 10 15
 Ser Val Pro Ala Glu Ser Ala Ala Val Leu Asn Asp Tyr Ser Leu Thr
 20 25 30
 40 Lys Ser His Glu Met Glu Asn Val Asp Ser Gly Glu Gly Pro Ala Asn
 35 40 45
 Glu Asp Glu Asp Ile Gly Asp Asp Ser Met Lys Val Lys Asp Glu Tyr
 50 55 60
 Ser Glu Arg Asp Glu Asn Val Leu Lys Ser Glu Pro Met Gly Asn Ala
 65 70 75 80
 45 Glu Glu Pro Glu Ile Pro Tyr Ser Tyr Ser Arg Glu Tyr Asn Glu Tyr
 85 90 95
 Glu Asn Ile Lys Leu Glu Arg His Val Val Ser Phe Asp Ser Ser Arg
 100 105 110
 Pro Thr Ser Gly Lys Met Asn Cys Asp Val Cys Gly Leu Ser Cys Ile
 115 120 125
 50 Ser Phe Asn Val Leu Met Val His Lys Arg Ser His Thr Gly Glu Arg
 130 135 140
 Pro Phe Gln Cys Asn Gln Cys Gly Ala Ser Phe Thr Gln Lys Gly Asn
 145 150 155 160
 55

	Leu	Leu	Arg	His	Ile	Lys	Leu	His	Thr	Gly	Glu	Lys	Pro	Phe	Lys	Cys
					165					170					175	
5	His	Leu	Cys	Asn	Tyr	Ala	Cys	Gln	Arg	Arg	Asp	Ala	Leu	Thr	Gly	His
				180					185					190		
	Leu	Arg	Thr	His	Ser	Val	Glu	Lys	Pro	Tyr	Lys	Cys	Glu	Phe	Cys	Gly
			195					200					205			
	Arg	Ser	Tyr	Lys	Gln	Arg	Ser	Ser	Leu	Glu	Glu	His	Lys	Glu	Arg	Cys
		210					215					220				
10	Arg	Thr	Phe	Leu	Gln	Ser	Thr	Asp	Pro	Gly	Asp	Thr	Ala	Ser	Ala	Glu
	225				230						235					240
	Ala	Arg	His	Ile	Lys	Ala	Glu	Met	Gly	Ser	Glu	Arg	Ala	Leu	Val	Leu
				245						250					255	
	Asp	Arg	Leu	Ala	Ser	Asn	Val	Ala	Lys	Arg	Lys	Ser	Ser	Met	Pro	Gln
				260					265					270		
15	Lys	Phe	Ile	Gly	Glu	Lys	Arg	His	Cys	Phe	Asp	Val	Asn	Tyr	Asn	Ser
			275					280					285			
	Ser	Tyr	Met	Tyr	Glu	Lys	Glu	Ser	Glu	Leu	Ile	Gln	Thr	Arg	Met	Met
		290					295					300				
20	Asp	Gln	Ala	Ile	Asn	Asn	Ala	Ile	Ser	Tyr	Leu	Gly	Ala	Glu	Ala	Leu
	305				310						315					320
	Cys	Pro	Leu	Val	Gln	Thr	Pro	Pro	Ala	Pro	Thr	Ser	Glu	Met	Val	Pro
				325						330					335	
	Val	Ile	Ser	Ser	Met	Tyr	Pro	Ile	Ala	Leu	Thr	Arg	Ala	Glu	Met	Ser
				340					345					350		
25	Asn	Gly	Ala	Pro	Gln	Glu	Leu	Glu	Arg	Lys	Ser	Ile	Leu	Leu	Pro	Glu
			355					360					365			
	Lys	Ser	Val	Pro	Ser	Glu	Arg	Gly	Leu	Ser	Pro	Asn	Asn	Ser	Gly	His
			370				375					380				
	Asp	Ser	Thr	Asp	Thr	Asp	Ser	Asn	His	Glu	Glu	Arg	Gln	Asn	His	Ile
	385				390						395					400
30	Tyr	Gln	Gln	Asn	His	Met	Val	Leu	Ser	Arg	Ala	Arg	Asn	Gly	Met	Pro
				405						410					415	
	Leu	Leu	Lys	Glu	Val	Pro	Arg	Ser	Tyr	Glu	Leu	Leu	Lys	Pro	Pro	Pro
				420					425					430		
	Ile	Cys	Pro	Arg	Asp	Ser	Val	Lys	Val	Ile	Asp	Lys	Glu	Gly	Glu	Val
			435					440					445			
35	Met	Asp	Val	Tyr	Arg	Cys	Asp	His	Cys	Arg	Val	Leu	Phe	Leu	Asp	Tyr
		450				455						460				
	Val	Met	Phe	Thr	Ile	His	Met	Gly	Cys	His	Gly	Phe	Arg	Asp	Pro	Phe
	465				470						475					480
	Glu	Cys	Asn	Met	Cys	Gly	Asp	Arg	Ser							

<210> 81

<211> 440

<212> PRT

<213> Homo sapiens

<400> 81

5 Met Pro Ile Pro Pro Pro Pro Pro Pro Pro Gly Pro Pro Pro Pro
 1 5 10 15
 Pro Thr Phe His Gln Ala Asn Thr Glu Gln Pro Lys Leu Ser Arg Asp
 20 25 30
 Glu Gln Arg Gly Arg Gly Ala Leu Leu Gln Asp Ile Cys Lys Gly Thr
 35 40 45
 Lys Leu Lys Lys Val Thr Asn Ile Asn Asp Arg Ser Ala Pro Ile Leu
 50 55 60
 10 Glu Lys Pro Lys Gly Ser Ser Gly Gly Tyr Gly Ser Gly Gly Ala Ala
 65 70 75 80
 Leu Gln Pro Lys Gly Gly Leu Phe Gln Gly Gly Val Leu Lys Leu Arg
 85 90 95
 Pro Val Gly Ala Lys Asp Gly Ser Glu Asn Leu Ala Gly Lys Pro Ala
 100 105 110
 15 Leu Gln Ile Pro Ser Ser Arg Ala Ala Ala Pro Arg Pro Pro Val Ser
 115 120 125
 Ala Ala Ser Gly Arg Pro Gln Asp Asp Thr Asp Ser Ser Arg Ala Ser
 130 135 140
 20 Leu Pro Glu Leu Pro Arg Met Gln Arg Pro Ser Leu Pro Asp Leu Ser
 145 150 155 160
 Arg Pro Asn Thr Thr Ser Ser Thr Gly Met Lys His Ser Ser Ser Ala
 165 170 175
 Pro Pro Pro Pro Pro Gly Arg Arg Ala Asn Ala Pro Pro Thr Pro
 180 185 190
 25 Leu Pro Met His Ser Ser Lys Ala Pro Ala Tyr Asn Arg Glu Lys Pro
 195 200 205
 Leu Pro Pro Thr Pro Gly Gln Arg Leu His Pro Gly Arg Glu Gly Pro
 210 215 220
 Pro Ala Pro Pro Pro Val Lys Pro Pro Pro Ser Pro Val Asn Ile Arg
 225 230 235 240
 30 Thr Gly Pro Ser Gly Gln Ser Leu Ala Pro Pro Pro Pro Tyr Arg
 245 250 255
 Gln Pro Pro Gly Val Pro Asn Gly Pro Ser Ser Pro Thr Asn Glu Ser
 260 265 270
 35 Ala Pro Glu Leu Pro Gln Arg His Asn Ser Leu His Arg Lys Thr Pro
 275 280 285
 Gly Pro Val Arg Gly Leu Ala Pro Pro Pro Thr Ser Ala Ser Pro
 290 295 300
 Ser Leu Leu Ser Asn Arg Pro Pro Pro Pro Ala Arg Asp Pro Pro Ser
 305 310 315 320
 40 Arg Gly Ala Ala Pro Pro Pro Pro Pro Pro Val Ile Arg Asn Gly Ala
 325 330 335
 Arg Asp Ala Pro Pro Pro Pro Pro Tyr Arg Met His Gly Ser Glu
 340 345 350
 Pro Pro Ser Arg Gly Lys Pro Pro Pro Pro Ser Arg Thr Pro Ala
 355 360 365
 45 Gly Pro Pro Pro Pro Pro Pro Pro Pro Leu Arg Asn Gly His Arg Asp
 370 375 380
 Ser Ile Thr Thr Val Arg Ser Phe Leu Asp Asp Phe Glu Ser Lys Tyr
 385 390 395 400
 Ser Phe His Pro Val Glu Asp Phe Pro Ala Pro Glu Glu Tyr Lys His
 405 410 415
 50 Phe Gln Arg Ile Tyr Pro Ser Lys Thr Asn Arg Ala Ala Arg Gly Ala
 420 425 430
 Pro Pro Leu Pro Pro Ile Leu Arg
 435 440

55

<210> 82

<211> 205

<212> PRT

<213> Homo sapiens

<400> 82

```

Met Ser Ile Met Ser Tyr Asn Gly Gly Ala Val Met Ala Met Lys Gly
1          5          10          15
Lys Asn Cys Val Ala Ile Ala Ala Asp Arg Arg Phe Gly Ile Gln Ala
20          25          30
Gln Met Val Thr Thr Asp Phe Gln Lys Ile Phe Pro Met Gly Asp Arg
35          40          45
Leu Tyr Ile Gly Leu Ala Gly Leu Ala Thr Asp Val Gln Thr Val Ala
50          55          60
Gln Arg Leu Lys Phe Arg Leu Asn Leu Tyr Glu Leu Lys Glu Gly Arg
65          70          75          80
Gln Ile Lys Pro Tyr Thr Leu Met Ser Met Val Ala Asn Leu Leu Tyr
85          90          95
Glu Lys Arg Phe Gly Pro Tyr Tyr Thr Glu Pro Val Ile Ala Gly Leu
100          105          110
Asp Pro Lys Thr Phe Lys Pro Phe Ile Cys Ser Leu Asp Leu Ile Gly
115          120          125
Cys Pro Met Val Thr Asp Asp Phe Val Val Ser Gly Thr Cys Ala Glu
130          135          140
Gln Met Tyr Gly Met Cys Glu Ser Leu Trp Glu Pro Asn Met Asp Pro
145          150          155          160
Asp His Leu Phe Glu Thr Ile Ser Gln Ala Met Leu Asn Ala Val Asp
165          170          175
Arg Asp Ala Val Ser Gly Met Gly Val Ile Val His Ile Ile Glu Lys
180          185          190
Asp Lys Ile Thr Thr Arg Thr Leu Lys Ala Arg Met Asp
195          200          205

```

<210> 83

<211> 190

<212> PRT

<213> Homo sapiens

<400> 83

```

Leu Thr Arg Ser Cys Ser Thr Cys Cys Pro Ala Val Ala Cys Leu Val
1          5          10          15
Gly Arg Gly Val Val Thr Ser Gly Ala Met His Gln Cys Trp Gly Glu
20          25          30
Glu Met Leu Gln Gly Met Leu Leu Trp Gly Trp Ala Thr Cys Pro Leu
35          40          45
Ser Asn Pro Gly Arg Trp Gly Arg Thr Val Gly Leu Gln His Pro Ala
50          55          60
Val Val Ser Ala Phe Arg Ala Leu Leu Leu Leu Met Leu Thr Val His
65          70          75          80

```

Val Ser Tyr Leu Ser Leu Ile Arg Phe Asp Tyr Gly Tyr Asn Leu Val
 85 90 95
 5 Ala Asn Val Ala Ile Gly Leu Val Asn Val Val Trp Trp Leu Ala Trp
 100 105 110
 Cys Leu Trp Asn Gln Arg Arg Leu Pro His Val Arg Lys Cys Val Val
 115 120 125
 Val Val Leu Leu Leu Gln Gly Leu Ser Leu Leu Glu Leu Leu Asp Phe
 130 135 140
 10 Pro Pro Leu Phe Trp Val Leu Asp Ala His Ala Ile Trp His Ile Ser
 145 150 155 160
 Thr Ile Pro Val His Val Leu Phe Phe Ser Phe Leu Glu Asp Asp Ser
 165 170 175
 Leu Tyr Leu Leu Lys Glu Ser Glu Asp Lys Phe Lys Leu Asp
 180 185 190
 15
 <210> 84
 <211> 368
 20 <212> PRT
 <213> Homo sapiens
 25
 <400> 84
 Ala Pro Pro Pro Ala Ala Ser Gln Gly Glu Arg Met Ala Gly Leu Ala
 1 5 10 15
 Ala Arg Leu Val Leu Leu Ala Gly Ala Ala Ala Leu Ala Ser Gly Ser
 20 25 30
 30 Gln Gly Asp Arg Glu Pro Val Tyr Arg Asp Cys Val Leu Gln Cys Glu
 35 40 45
 Glu Gln Asn Cys Ser Gly Gly Ala Leu Asn His Phe Arg Ser Arg Gln
 50 55 60
 Pro Ile Tyr Met Ser Leu Ala Gly Trp Thr Cys Arg Asp Asp Cys Lys
 65 70 75 80
 35 Tyr Glu Cys Met Trp Val Thr Val Gly Leu Tyr Leu Gln Glu Gly His
 85 90 95
 Lys Val Pro Gln Phe His Gly Lys Trp Pro Phe Ser Arg Phe Leu Phe
 100 105 110
 Phe Gln Glu Pro Ala Ser Ala Val Ala Ser Phe Leu Asn Gly Leu Ala
 115 120 125
 40 Ser Leu Val Met Leu Cys Arg Tyr Arg Thr Phe Val Pro Ala Ser Ser
 130 135 140
 Pro Met Tyr His Thr Cys Val Ala Phe Ala Trp Val Ser Leu Asn Ala
 145 150 155 160
 Trp Phe Trp Ser Thr Val Phe His Thr Arg Asp Thr Asp Leu Thr Glu
 165 170 175
 45 Lys Met Asp Tyr Phe Cys Ala Ser Thr Val Ile Leu His Ser Ile Tyr
 180 185 190
 Leu Cys Cys Val Arg Thr Val Gly Leu Gln His Pro Ala Val Val Ser
 195 200 205
 50 Ala Phe Arg Ala Leu Leu Leu Leu Met Leu Thr Val His Val Ser Tyr
 210 215 220
 Leu Ser Leu Ile Arg Phe Asp Tyr Gly Tyr Asn Leu Val Ala Asn Val
 225 230 235 240
 Ala Ile Gly Leu Val Asn Val Val Trp Trp Leu Ala Trp Cys Leu Trp
 245 250 255
 55

Asn Gln Arg Arg Leu Pro His Val Arg Lys Cys Val Val Val Val Leu
 260 265 270
 Leu Leu Gln Gly Leu Ser Leu Leu Glu Leu Leu Asp Phe Pro Pro Leu
 275 280 285
 Phe Trp Val Leu Asp Ala His Ala Ile Trp His Ile Ser Thr Ile Pro
 290 295 300
 Val His Val Leu Phe Phe Ser Phe Leu Glu Asp Asp Ser Leu Tyr Leu
 305 310 315 320
 Leu Lys Glu Ser Glu Asp Lys Phe Lys Leu Val Glu Ala Asp Trp Ile
 325 330 335
 Phe Ala Leu Pro Leu Thr Pro Cys Pro Ser Leu Arg Glu Gly Ser Tyr
 340 345 350
 Ala Arg Thr Pro Thr Ser Gly Thr Arg Val Ala Cys Ala Ser Phe Phe
 355 360 365

<210> 85

<211> 190

<212> PRT

<213> Homo sapiens

<400> 85

Leu Thr Arg Ser Cys Ser Thr Cys Cys Pro Ala Val Ala Cys Leu Val
 1 5 10 15
 Gly Arg Gly Val Val Thr Ser Gly Ala Met His Gln Cys Trp Gly Glu
 20 25 30
 Glu Met Leu Gln Gly Met Leu Leu Trp Gly Trp Ala Thr Cys Pro Leu
 35 40 45
 Ser Asn Pro Gly Arg Trp Gly Arg Thr Val Gly Leu Gln His Pro Ala
 50 55 60
 Val Val Ser Ala Phe Arg Ala Leu Leu Leu Leu Met Leu Thr Val His
 65 70 75 80
 Val Ser Tyr Leu Ser Leu Ile Arg Phe Asp Tyr Gly Tyr Asn Leu Val
 85 90 95
 Ala Asn Val Ala Ile Gly Leu Val Asn Val Val Trp Trp Leu Ala Trp
 100 105 110
 Cys Leu Trp Asn Gln Arg Arg Leu Pro His Val Arg Lys Cys Val Val
 115 120 125
 Val Val Leu Leu Leu Gln Gly Leu Ser Leu Leu Glu Leu Leu Asp Phe
 130 135 140
 Pro Pro Leu Phe Trp Val Leu Asp Ala His Ala Ile Trp His Ile Ser
 145 150 155 160
 Thr Ile Pro Val His Val Leu Phe Phe Ser Phe Leu Glu Asp Asp Ser
 165 170 175
 Leu Tyr Leu Leu Lys Glu Ser Glu Asp Lys Phe Lys Leu Asp
 180 185 190

<210> 86

<211> 318

<212> PRT

<213> Homo sapiens

5

<400> 86

10

15

20

25

30

35

40

Met Ala Gly Leu Ala Ala Arg Leu Val Leu Leu Ala Gly Ala Ala Ala
 1 5 10 15
 Leu Ala Ser Gly Ser Gln Gly Asp Arg Glu Pro Val Tyr Arg Asp Cys
 20 25 30
 Val Leu Gln Cys Glu Glu Gln Asn Cys Ser Gly Gly Ala Leu Asn His
 35 40 45
 Phe Arg Ser Arg Gln Pro Ile Tyr Met Ser Leu Ala Gly Trp Thr Cys
 50 55 60
 Arg Asp Asp Cys Lys Tyr Glu Cys Met Trp Val Thr Val Gly Leu Tyr
 65 70 75 80
 Leu Gln Glu Gly His Lys Val Pro Gln Phe His Gly Lys Trp Pro Phe
 85 90 95
 Ser Arg Phe Leu Phe Phe Gln Glu Pro Ala Ser Ala Val Ala Ser Phe
 100 105 110
 Leu Asn Gly Leu Ala Ser Leu Val Met Leu Cys Arg Tyr Arg Thr Phe
 115 120 125
 Val Pro Ala Ser Ser Pro Met Tyr His Thr Cys Val Ala Phe Ala Trp
 130 135 140
 Val Ser Leu Asn Ala Trp Phe Trp Ser Thr Val Phe His Thr Arg Asp
 145 150 155 160
 Thr Asp Leu Gln Arg Lys Trp Thr Thr Ser Val Pro Pro Val Ser Tyr
 165 170 175
 Thr Gln Ser Thr Cys Ala Ala Ser Gly Pro Trp Gly Cys Ser Thr Gln
 180 185 190
 Leu Trp Ser Ser Ala Phe Arg Ala Leu Leu Leu Leu Met Leu Thr Val
 195 200 205
 His Val Ser Tyr Leu Ser Leu Ile Arg Phe Asp Tyr Gly Tyr Asn Leu
 210 215 220
 Val Ala Asn Val Ala Ile Gly Leu Val Asn Val Val Trp Trp Leu Ala
 225 230 235 240
 Trp Cys Leu Trp Asn Gln Arg Arg Leu Pro His Val Arg Lys Cys Val
 245 250 255
 Val Val Val Leu Leu Leu Gln Gly Leu Ser Leu Leu Glu Leu Leu Asp
 260 265 270
 Phe Pro Pro Leu Phe Trp Val Leu Asp Ala His Ala Ile Trp His Ile
 275 280 285
 Ser Thr Ile Pro Val His Val Leu Phe Phe Ser Phe Leu Glu Asp Asp
 290 295 300
 Ser Leu Tyr Leu Leu Lys Glu Ser Glu Asp Lys Phe Lys Leu
 305 310 315

45

<210> 87

<211> 226

<212> PRT

50

<213> Homo sapiens

55

<400> 87

Met Ala Gly Leu Ala Ala Arg Leu Val Leu Leu Ala Gly Ala Ala Ala
 1 5 10 15
 Leu Ala Ser Gly Ser Gln Gly Asp Arg Glu Pro Val Tyr Arg Asp Cys
 20 25 30
 Val Leu Gln Cys Glu Glu Gln Asn Cys Ser Gly Gly Ala Leu Asn His
 35 40 45
 Phe Arg Ser Arg Gln Pro Ile Tyr Met Ser Leu Ala Gly Trp Thr Cys
 50 55 60
 Arg Asp Asp Cys Lys Tyr Glu Cys Met Trp Val Thr Val Gly Leu Tyr
 65 70 75 80
 Leu Gln Glu Gly His Lys Val Pro Gln Phe His Gly Lys Trp Pro Phe
 85 90 95
 Ser Arg Phe Leu Phe Phe Gln Glu Pro Ala Ser Ala Val Ala Ser Phe
 100 105 110
 Leu Asn Gly Leu Ala Ser Leu Val Met Leu Cys Arg Tyr Arg Thr Phe
 115 120 125
 Val Pro Ala Ser Ser Pro Met Tyr His Thr Cys Val Ala Phe Ala Trp
 130 135 140
 Val Ser Leu Asn Ala Trp Phe Trp Ser Thr Val Phe His Thr Arg Asp
 145 150 155 160
 Thr Asp Leu Thr Glu Lys Met Asp Tyr Phe Cys Ala Ser Thr Val Ile
 165 170 175
 Leu His Ser Ile Tyr Leu Cys Cys Val Arg Pro Gly Gln Arg Gly Val
 180 185 190
 Val Ala Gly Leu Val Pro Val Glu Pro Ala Ala Ala Ser Arg Ala
 195 200 205
 Gln Val Arg Gly Gly Gly Leu Ala Ala Ala Gly Ala Val Pro Ala Arg
 210 215 220
 Ala Ala
 225

<210> 88

<211> 320

<212> PRT

<213> Homo sapiens

<400> 88

Met Ala Gly Leu Ala Ala Arg Leu Val Leu Leu Ala Gly Ala Ala Ala
 1 5 10 15
 Leu Ala Ser Gly Ser Gln Gly Asp Arg Glu Pro Val Tyr Arg Asp Cys
 20 25 30
 Val Leu Gln Cys Glu Glu Gln Asn Cys Ser Gly Gly Ala Leu Asn His
 35 40 45
 Phe Arg Ser Arg Gln Pro Ile Tyr Met Ser Leu Ala Gly Trp Thr Cys
 50 55 60
 Arg Asp Asp Cys Lys Tyr Glu Cys Met Trp Val Thr Val Gly Leu Tyr
 65 70 75 80
 Leu Gln Glu Gly His Lys Val Pro Gln Phe His Gly Lys Trp Pro Phe
 85 90 95
 Ser Arg Phe Leu Phe Phe Gln Glu Pro Ala Ser Ala Val Ala Ser Phe
 100 105 110
 Leu Asn Gly Leu Ala Ser Leu Val Met Leu Cys Arg Tyr Arg Thr Phe
 115 120 125

	Val	Pro	Ala	Ser	Ser	Pro	Met	Tyr	His	Thr	Cys	Val	Ala	Phe	Ala	Trp
	130						135					140				
5	Val	Ser	Leu	Asn	Ala	Trp	Phe	Trp	Ser	Thr	Val	Phe	His	Thr	Arg	Asp
	145					150					155					160
	Thr	Asp	Leu	Thr	Glu	Lys	Met	Asp	Tyr	Phe	Cys	Ala	Ser	Thr	Val	Ile
					165						170				175	
	Leu	His	Ser	Ile	Tyr	Leu	Cys	Cys	Val	Arg	Thr	Val	Gly	Leu	Gln	His
				180					185					190		
10	Pro	Ala	Val	Val	Ser	Ala	Phe	Arg	Ala	Leu	Leu	Leu	Leu	Met	Leu	Thr
								200						205		
	Val	His	Val	Ser	Tyr	Leu	Ser	Leu	Ile	Arg	Phe	Asp	Tyr	Gly	Tyr	Asn
	210						215					220				
	Leu	Val	Ala	Asn	Val	Ala	Ile	Gly	Leu	Val	Asn	Val	Val	Trp	Trp	Leu
	225					230					235					240
15	Ala	Trp	Cys	Leu	Trp	Asn	Gln	Arg	Arg	Leu	Pro	His	Val	Arg	Lys	Cys
					245					250					255	
	Val	Val	Val	Val	Leu	Leu	Leu	Gln	Gly	Leu	Ser	Leu	Leu	Glu	Leu	Leu
					260				265					270		
	Asp	Phe	Pro	Pro	Leu	Phe	Trp	Val	Leu	Asp	Ala	His	Ala	Ile	Trp	His
			275					280					285			
20	Ile	Ser	Thr	Ile	Pro	Val	His	Val	Leu	Phe	Phe	Ser	Phe	Leu	Glu	Asp
	290						295					300				
	Asp	Ser	Leu	Tyr	Leu	Leu	Lys	Glu	Ser	Glu	Asp	Lys	Phe	Lys	Leu	Asp
	305					310					315					320
25	<210> 89															
	<211> 217															
	<212> PRT															
30	<213> Homo sapiens															
	<400> 89															
35	Ala	Pro	Pro	Pro	Ala	Ala	Ser	Gln	Gly	Glu	Arg	Met	Ala	Gly	Leu	Ala
	1				5					10					15	
	Ala	Arg	Leu	Val	Leu	Leu	Ala	Gly	Ala	Ala	Ala	Leu	Ala	Ser	Gly	Ser
				20					25					30		
	Gln	Gly	Asp	Arg	Glu	Pro	Val	Tyr	Arg	Asp	Cys	Val	Leu	Gln	Cys	Glu
			35					40					45			
40	Glu	Gln	Asn	Cys	Ser	Gly	Gly	Ala	Leu	Asn	His	Phe	Arg	Ser	Arg	Gln
	50					55						60				
	Pro	Ile	Tyr	Met	Ser	Leu	Ala									

Lys Met Asp Tyr Phe Cys Ala Ser Thr Val Ile Leu His Ser Ile Tyr
 180 185 190
 Leu Cys Cys Val Ser Phe Leu Glu Asp Asp Ser Leu Tyr Leu Leu Lys
 195 200 205
 Glu Ser Glu Asp Lys Phe Lys Leu Asp
 210 215

<210> 90

<211> 153

<212> PRT

<213> Homo sapiens

<400> 90

Met Asn Val Gly Thr Ala His Ser Glu Val Asn Pro Asn Thr Arg Val
 1 5 10 15
 Met Asn Ser Arg Gly Ile Trp Leu Ser Tyr Val Leu Ala Ile Gly Leu
 20 25 30
 Leu His Ile Val Leu Leu Ser Ile Pro Phe Val Ser Val Pro Val Val
 35 40 45
 Trp Thr Leu Thr Asn Leu Ile His Asn Met Gly Met Tyr Ile Phe Leu
 50 55 60
 His Thr Val Lys Gly Thr Pro Phe Glu Thr Pro Asp Gln Gly Lys Ala
 65 70 75 80
 Arg Leu Leu Thr His Trp Glu Gln Met Asp Tyr Gly Val Gln Phe Thr
 85 90 95
 Ala Ser Arg Lys Phe Leu Thr Ile Thr Pro Ile Val Leu Tyr Phe Leu
 100 105 110
 Thr Ser Phe Tyr Thr Lys Tyr Asp Gln Ile His Phe Val Leu Asn Thr
 115 120 125
 Val Ser Leu Met Ser Val Leu Ile Pro Lys Leu Pro Gln Leu His Gly
 130 135 140
 Val Arg Ile Phe Gly Ile Asn Lys Tyr
 145 150

<210> 91

<211> 436

<212> PRT

<213> Homo sapiens

<400> 91

Met Arg Arg Asp Val Asn Gly Val Thr Lys Ser Arg Phe Glu Met Phe
 1 5 10 15
 Ser Asn Ser Asp Glu Ala Val Ile Asn Lys Lys Leu Pro Lys Glu Leu
 20 25 30
 Leu Leu Arg Ile Phe Ser Phe Leu Asp Val Val Thr Leu Cys Arg Cys
 35 40 45
 Ala Gln Val Ser Arg Ala Trp Asn Val Leu Ala Leu Asp Gly Ser Asn
 50 55 60

5 Trp Gln Arg Ile Asp Leu Phe Asp Phe Gln Arg Asp Ile Glu Gly Arg
 65 70 75 80
 Val Val Glu Asn Ile Ser Lys Arg Cys Gly Gly Phe Leu Arg Lys Leu
 85 90 95
 Ser Leu Arg Gly Cys Leu Gly Val Gly Asp Asn Ala Leu Arg Thr Phe
 100 105 110
 Ala Gln Asn Cys Arg Asn Ile Glu Val Leu Asn Leu Asn Gly Cys Thr
 115 120 125
 10 Lys Thr Thr Asp Ala Thr Cys Thr Ser Leu Ser Lys Phe Cys Ser Lys
 130 135 140
 Leu Arg His Leu Asp Leu Ala Ser Cys Thr Ser Ile Thr Asn Met Ser
 145 150 155 160
 Leu Lys Ala Leu Ser Glu Gly Cys Pro Leu Leu Glu Gln Leu Asn Ile
 165 170 175
 15 Ser Trp Cys Asp Gln Val Thr Lys Asp Gly Ile Gln Ala Leu Val Arg
 180 185 190
 Gly Cys Gly Gly Leu Lys Ala Leu Phe Leu Lys Gly Cys Thr Gln Leu
 195 200 205
 Glu Asp Glu Ala Leu Lys Tyr Ile Gly Ala His Cys Pro Glu Leu Val
 210 215 220
 20 Thr Leu Asn Leu Gln Thr Cys Leu Gln Ile Thr Asp Glu Gly Leu Ile
 225 230 235 240
 Thr Ile Cys Arg Gly Cys His Lys Leu Gln Ser Leu Cys Ala Ser Gly
 245 250 255
 Cys Ser Asn Ile Thr Asp Ala Ile Leu Asn Ala Leu Gly Gln Asn Cys
 260 265 270
 25 Pro Arg Leu Arg Ile Leu Glu Val Ala Arg Cys Ser Gln Leu Thr Asp
 275 280 285
 Val Gly Phe Thr Thr Leu Ala Arg Asn Cys His Glu Leu Glu Lys Met
 290 295 300
 30 Asp Leu Glu Glu Cys Val Gln Ile Thr Asp Ser Thr Leu Ile Gln Leu
 305 310 315 320
 Ser Ile His Cys Pro Arg Leu Gln Val Leu Ser Leu Ser His Cys Glu
 325 330 335
 Leu Ile Thr Asp Asp Gly Ile Arg His Leu Gly Asn Gly Ala Cys Ala
 340 345 350
 35 His Asp Gln Leu Glu Val Ile Glu Leu Asp Asn Cys Pro Leu Ile Thr
 355 360 365
 Asp Ala Ser Leu Glu His Leu Lys Ser Cys His Ser Leu Glu Arg Ile
 370 375 380
 Glu Leu Tyr Asp Cys Gln Gln Ile Thr Arg Ala Gly Ile Lys Arg Leu
 385 390 395 400
 40 Arg Thr His Leu Pro Asn Ile Lys Val His Ala Tyr Phe Ala Pro Val
 405 410 415
 Thr Pro Pro Pro Ser Val Gly Gly Ser Arg Gln Arg Phe Cys Arg Cys
 420 425 430
 45 Cys Ile Ile Leu
 435

 <210> 92
 <211> 204
 50 <212> PRT
 <213> Homo sapiens

<400> 92

Met Asp Pro Lys Asp Arg Lys Lys Ile Gln Phe Ser Val Pro Ala Pro
 1 5 10 15
 Pro Ser Gln Leu Asp Pro Arg Gln Val Glu Met Ile Arg Arg Arg Arg
 20 25 30
 Pro Thr Pro Ala Met Leu Phe Arg Leu Ser Glu His Ser Ser Pro Glu
 35 40 45
 Glu Glu Ala Ser Pro His Gln Arg Ala Ser Gly Glu Gly His His Leu
 50 55 60
 Lys Ser Lys Arg Pro Asn Pro Cys Ala Tyr Thr Pro Pro Ser Leu Lys
 65 70 75 80
 Ala Val Gln Arg Ile Ala Glu Ser His Leu Gln Ser Ile Ser Asn Leu
 85 90 95
 Asn Glu Asn Gln Ala Ser Glu Glu Glu Asp Glu Leu Gly Glu Leu Arg
 100 105 110
 Glu Leu Gly Tyr Pro Arg Glu Glu Asp Glu Glu Glu Glu Asp Asp
 115 120 125
 Glu Glu Glu Glu Glu Glu Glu Asp Ser Gln Ala Glu Val Leu Lys Val
 130 135 140
 Ile Arg Gln Ser Ala Gly Gln Lys Thr Thr Cys Gly Gln Gly Leu Glu
 145 150 155 160
 Gly Pro Trp Glu Arg Pro Pro Pro Leu Asp Glu Ser Glu Arg Asp Gly
 165 170 175
 Gly Ser Glu Asp Gln Val Glu Asp Pro Ala Leu Ser Glu Pro Gly Glu
 180 185 190
 Glu Pro Gln Arg Pro Ser Pro Ser Glu Pro Gly Thr
 195 200

<210> 93

<211> 115

<212> PRT

<213> Homo sapiens

<400> 93

Met Ser Gly Glu Pro Gly Gln Thr Ser Val Ala Pro Pro Pro Glu Glu
 1 5 10 15
 Val Glu Pro Gly Ser Gly Val Arg Ile Val Val Glu Tyr Cys Glu Pro
 20 25 30
 Cys Gly Phe Glu Ala Thr Tyr Leu Glu Leu Ala Ser Ala Val Lys Glu
 35 40 45
 Gln Tyr Pro Gly Ile Glu Ile Glu Ser Arg Leu Gly Gly Thr Gly Ala
 50 55 60
 Phe Glu Ile Glu Ile Asn Gly Gln Leu Val Phe Ser Lys Leu Glu Asn
 65 70 75 80
 Gly Gly Phe Pro Tyr Glu Lys Asp Leu Ile Glu Ala Ile Arg Arg Ala
 85 90 95
 Ser Asn Gly Glu Thr Leu Glu Lys Ile Thr Asn Ser Arg Pro Pro Cys
 100 105 110
 Val Ile Leu
 115

<210> 94

<211> 144

<212> PRT

<213> Homo sapiens

<400> 94

Met Gly Ala Val Val Leu Cys Arg Pro Ser Pro Leu Asn Phe Leu Ile
 1 5 10 15
 Gln Thr Gly Thr Gly Gln Gly Leu Ser Cys Gly Ser His Met Trp Arg
 20 25 30
 Cys Glu Ala Thr Pro Cys Gly Val Cys Gly Glu Ser Pro Val Gly Ser
 35 40 45
 Leu Leu Lys Gln His Arg Gly Arg Gly Lys Thr Trp Pro Val Gly Thr
 50 55 60
 Val Ser Ala Cys Arg Glu Glu Ser Glu Ala Gly Ser Leu Ser Leu Gly
 65 70 75 80
 Trp Ser Leu Leu Pro Ser Pro Val Gly Leu Gly Ala Val Leu Ile Leu
 85 90 95
 Lys Arg Cys Gly Ser Leu Cys Pro Leu Pro Gly Val Gln Gly Asn Arg
 100 105 110
 Arg Gly His Trp Ala Cys Phe Leu Pro Pro Asp Pro Ala Ser Pro Thr
 115 120 125
 Pro Cys Ile Ile Gly Asn Phe His Leu Lys Ile Phe Leu Ser Lys Val
 130 135 140

<210> 95

<211> 425

<212> PRT

<213> Homo sapiens

<400> 95

Met Gly Gly Gly Asp Leu Asn Leu Lys Lys Ser Trp His Pro Gln Thr
 1 5 10 15
 Leu Arg Asn Val Glu Lys Val Trp Lys Ala Glu Gln Lys His Glu Ala
 20 25 30
 Glu Arg Lys Lys Ile Glu Glu Leu Gln Arg Glu Leu Arg Glu Glu Arg
 35 40 45
 Ala Arg Glu Glu Met Gln Arg Tyr Ala Glu Asp Val Gly Ala Val Lys
 50 55 60
 Lys Lys Glu Glu Lys Leu Asp Trp Met Tyr Gln Gly Pro Gly Gly Met
 65 70 75 80
 Val Asn Arg Asp Glu Tyr Leu Leu Gly Arg Pro Ile Asp Lys Tyr Val
 85 90 95
 Phe Glu Lys Met Glu Glu Lys Glu Ala Gly Cys Ser Ser Glu Thr Gly
 100 105 110
 Leu Leu Pro Gly Ser Ile Phe Ala Pro Ser Gly Ala Asn Ser Leu Leu
 115 120 125
 Asp Met Ala Ser Lys Ile Arg Glu Asp Pro Leu Phe Ile Ile Arg Lys
 130 135 140
 Lys Glu Glu Glu Lys Lys Arg Glu Val Leu Asn Asn Pro Val Lys Met
 145 150 155 160

	Lys	Lys	Ile	Lys	Glu	Leu	Leu	Gln	Met	Ser	Leu	Glu	Lys	Lys	Glu	Lys
					165					170					175	
5	Lys	Lys	Lys	Lys	Glu	Lys	Lys	Lys	Lys	His	Lys	Lys	His	Lys	His	Arg
				180					185					190		
	Ser	Ser	Ser	Ser	Asp	Arg	Ser	Ser	Ser	Glu	Asp	Glu	His	Ser	Ala	Gly
				195					200				205			
	Arg	Ser	Gln	Lys	Lys	Met	Ala	Asn	Ser	Ser	Pro	Val	Leu	Ser	Lys	Val
		210					215					220				
10	Pro	Gly	Tyr	Gly	Leu	Gln	Val	Arg	Asn	Ser	Asp	Arg	Asn	Gln	Gly	Leu
	225					230					235					240
	Gln	Gly	Pro	Leu	Thr	Ala	Glu	Gln	Lys	Arg	Gly	His	Gly	Met	Lys	Asn
				245						250					255	
	His	Ser	Arg	Ser	Arg	Ser	Ser	Ser	His	Ser	Pro	Pro	Arg	His	Ala	Ser
				260					265					270		
15	Lys	Lys	Ser	Thr	Arg	Glu	Ala	Gly	Ser	Arg	Asp	Arg	Arg	Ser	Arg	Ser
			275					280					285			
	Leu	Gly	Arg	Arg	Ser	Arg	Ser	Pro	Arg	Pro	Ser	Lys	Leu	His	Asn	Ser
		290					295					300				
	Lys	Val	Asn	Arg	Arg	Glu	Thr	Gly	Gln	Thr	Arg	Ser	Pro	Ser	Pro	Lys
	305					310					315					320
20	Lys	Glu	Val	Tyr	Gln	Arg	Arg	His	Ala	Pro	Gly	Tyr	Thr	Arg	Lys	Leu
					325					330					335	
	Ser	Ala	Glu	Glu	Leu	Glu	Arg	Lys	Arg	Gln	Glu	Met	Met	Glu	Asn	Ala
				340					345					350		
25	Lys	Trp	Arg	Glu	Glu	Glu	Arg	Leu	Asn	Ile	Leu	Lys	Arg	His	Ala	Lys
			355					360					365			
	Asp	Glu	Glu	Arg	Glu	Gln	Arg	Leu	Glu	Lys	Leu	Asp	Ser	Arg	Asp	Gly
		370					375					380				
	Lys	Phe	Ile	His	Arg	Met	Lys	Leu	Glu	Ser	Ala	Ser	Thr	Ser	Ser	Leu
	385					390					395					400
30	Glu	Asp	Arg	Val	Lys	Arg	Asn	Ile	Tyr	Ser	Leu	Gln	Arg	Thr	Ser	Val
					405					410					415	
	Ala	Leu	Glu	Lys	Asn	Phe	Met	Lys	Arg							
				420					425							

<210> 96

<211> 394

<212> PRT

<213> Homo sapiens

<400> 96

Met Phe Ser Val Phe Glu Glu Ile Thr Arg Ile Val Val Lys Glu Met
1 5 10 15
Asp Ala Gly Gly Asp Met Ile Ala Val Arg Ser Leu Val Asp Ala Asp
20 25 30

Arg	Phe	Arg	Cys	Phe	His	Leu	Val	Gly	Glu	Lys	Arg	Thr	Phe	Phe	Gly
		35					40					45			
Cys	Arg	His	Tyr	Thr	Thr	Gly	Leu	Thr	Leu	Met	Asp	Ile	Leu	Asp	Thr
		50				55					60				
His	Gly	Asp	Lys	Trp	Leu	Asp	Glu	Leu	Asp	Ser	Gly	Leu	Gln	Gly	Gln
65					70					75					80
Lys	Ala	Glu	Phe	Gln	Ile	Leu	Asp	Asn	Val	Asp	Ser	Thr	Gly	Glu	Leu
				85					90					95	

Ile Val Arg Leu Pro Lys Glu Ile Thr Ile Ser Gly Ser Phe Gln Gly
 100 105 110
 Phe His His Gln Lys Ile Lys Ile Ser Glu Asn Arg Ile Ser Gln Gln
 115 120 125
 Tyr Leu Ala Thr Leu Glu Asn Arg Lys Leu Lys Arg Glu Leu Pro Phe
 130 135 140
 Ser Phe Arg Ser Ile Asn Thr Arg Glu Asn Leu Tyr Leu Val Thr Glu
 145 150 155 160
 Thr Leu Glu Thr Val Lys Glu Glu Thr Leu Lys Ser Asp Arg Gln Tyr
 165 170 175
 Lys Phe Trp Ser Gln Ile Ser Gln Gly His Leu Ser Tyr Lys His Lys
 180 185 190
 Gly Gln Arg Glu Val Thr Ile Pro Pro Asn Arg Val Leu Ser Tyr Arg
 195 200 205
 Val Lys Gln Leu Val Phe Pro Asn Lys Glu Thr Met Arg Lys Ser Leu
 210 215 220
 Gly Ser Glu Asp Ser Arg Asn Met Lys Glu Lys Leu Glu Asp Met Glu
 225 230 235 240
 Ser Val Leu Lys Asp Leu Thr Glu Glu Lys Arg Lys Asp Val Leu Asn
 245 250 255
 Ser Leu Ala Lys Cys Leu Gly Lys Glu Asp Ile Arg Gln Asp Leu Glu
 260 265 270
 Gln Arg Val Ser Glu Val Leu Ile Ser Gly Glu Leu His Met Glu Asp
 275 280 285
 Pro Asp Lys Pro Leu Leu Ser Ser Leu Phe Asn Ala Ala Gly Val Leu
 290 295 300
 Val Glu Ala Arg Ala Lys Ala Ile Leu Asp Phe Leu Asp Ala Leu Leu
 305 310 315 320
 Glu Leu Ser Glu Glu Gln Gln Phe Val Ala Glu Ala Leu Glu Lys Gly
 325 330 335
 Thr Leu Pro Leu Leu Lys Asp Gln Val Lys Ser Val Met Glu Gln Asn
 340 345 350
 Trp Asp Glu Leu Ala Ser Ser Pro Pro Asp Met Asp Tyr Asp Pro Glu
 355 360 365
 Ala Arg Ile Leu Cys Ala Leu Tyr Val Val Val Ser Ile Leu Leu Glu
 370 375 380
 Leu Ala Glu Gly Pro Thr Ser Val Ser Ser
 385 390

<210> 97

<211> 456

<212> PRT

<213> Homo sapiens

<400> 97

Met Glu Gly Pro Glu Gly Leu Gly Arg Lys Gln Ala Cys Leu Ala Met
 1 5 10 15
 Leu Leu His Phe Leu Asp Thr Tyr Gln Gly Leu Leu Gln Glu Glu Glu
 20 25 30
 Gly Ala Gly His Ile Ile Lys Asp Leu Tyr Leu Leu Ile Met Lys Asp
 35 40 45
 Glu Ser Leu Tyr Gln Gly Leu Arg Glu Asp Thr Leu Arg Leu His Gln
 50 55 60

5 Leu Val Glu Thr Val Glu Leu Lys Ile Pro Glu Glu Asn Gln Pro Pro
 65 70 75 80
 Ser Lys Gln Val Lys Pro Leu Phe Arg His Phe Arg Arg Ile Asp Ser
 85 90 95
 10 Cys Leu Gln Thr Arg Val Ala Phe Arg Gly Ser Asp Glu Ile Phe Cys
 100 105 110
 Arg Val Tyr Met Pro Asp His Ser Tyr Val Thr Ile Arg Ser Arg Leu
 115 120 125
 15 Ser Ala Ser Val Gln Asp Ile Leu Gly Ser Val Thr Glu Lys Leu Gln
 130 135 140
 Tyr Ser Glu Glu Pro Ala Gly Arg Glu Asp Ser Leu Ile Leu Val Ala
 145 150 155 160
 Val Ser Ser Ser Gly Glu Lys Val Leu Leu Gln Pro Thr Glu Asp Cys
 165 170 175
 20 Val Phe Thr Ala Leu Gly Ile Asn Ser His Leu Phe Ala Cys Thr Arg
 180 185 190
 Asp Ser Tyr Glu Ala Leu Val Pro Leu Pro Glu Glu Ile Gln Val Ser
 195 200 205
 Pro Gly Asp Thr Glu Ile His Arg Val Glu Pro Glu Asp Val Ala Asn
 210 215 220
 25 His Leu Thr Ala Phe His Trp Glu Leu Phe Arg Cys Val His Glu Leu
 225 230 235 240
 Glu Phe Val Asp Tyr Val Phe His Gly Glu Arg Gly Arg Arg Glu Thr
 245 250 255
 Ala Asn Leu Glu Leu Leu Leu Gln Arg Cys Ser Glu Val Thr His Trp
 260 265 270
 Val Ala Thr Glu Val Leu Leu Cys Glu Ala Pro Gly Lys Arg Ala Gln
 275 280 285
 Leu Leu Lys Lys Phe Ile Lys Ile Ala Ala Leu Cys Lys Gln Asn Gln
 290 295 300
 30 Asp Leu Leu Ser Phe Tyr Ala Val Val Met Gly Leu Asp Asn Ala Ala
 305 310 315 320
 Val Ser Arg Leu Arg Leu Thr Trp Glu Lys Leu Pro Gly Lys Phe Lys
 325 330 335
 Asn Leu Phe Arg Lys Phe Glu Asn Leu Thr Asp Pro Cys Arg Asn His
 340 345 350
 35 Lys Ser Tyr Arg Glu Val Ile Ser Lys Met Lys Pro Pro Val Ile Pro
 355 360 365
 Phe Val Pro Leu Ile Leu Lys Asp Leu Thr Phe Leu His Glu Gly Ser
 370 375 380
 Lys Thr Leu Val Asp Gly Leu Val Asn Ile Glu Lys Leu His Ser Val
 385 390 395 400
 40 Ala Glu Lys Val Arg Thr Ile Arg Lys Tyr Arg Ser Arg Pro Leu Cys
 405 410 415
 Leu Asp Met Glu Ala Ser Pro Asn His Leu Gln Thr Lys Ala Tyr Val
 420 425 430
 Arg Gln Phe Gln Val Ile Asp Asn Gln Asn Leu Leu Phe Glu Leu Ser
 435 440 445
 45 Tyr Lys Leu Glu Ala Asn Ser Gln
 450 455

 50 <210> 98

 <211> 715

 <212> PRT

 <213> Homo sapiens

<400> 98
 5 Met Ser Gln Val Met Ser Ser Pro Leu Leu Ala Gly Gly His Ala Val
 1 5 10 15
 Ser Leu Ala Pro Cys Asp Glu Pro Arg Arg Thr Leu His Pro Ala Pro
 20 25 30
 Ser Pro Ser Leu Pro Pro Gln Cys Ser Tyr Tyr Thr Thr Glu Gly Trp
 35 40 45
 10 Gly Ala Gln Ala Leu Met Ala Pro Val Pro Cys Met Gly Pro Pro Gly
 50 55 60
 Arg Leu Gln Gln Ala Pro Gln Val Glu Ala Lys Ala Thr Cys Phe Leu
 65 70 75 80
 Pro Ser Pro Gly Glu Lys Ala Leu Gly Thr Pro Glu Asp Leu Asp Ser
 85 90 95
 15 Tyr Ile Asp Phe Ser Leu Glu Ser Leu Asn Gln Met Ile Leu Glu Leu
 100 105 110
 Asp Pro Thr Phe Gln Leu Leu Pro Pro Gly Thr Gly Gly Ser Gln Ala
 115 120 125
 20 Glu Leu Ala Gln Ser Thr Met Ser Met Arg Lys Lys Glu Glu Ser Glu
 130 135 140
 Ala Leu Asp Ile Lys Tyr Ile Glu Val Thr Ser Ala Arg Ser Arg Cys
 145 150 155 160
 His Asp Trp Pro Gln His Cys Ser Ser Pro Ser Val Thr Pro Pro Phe
 165 170 175
 25 Gly Ser Pro Arg Ser Gly Gly Leu Leu Ser Arg Asp Val Pro Arg
 180 185 190
 Glu Thr Arg Ser Ser Ser Glu Ser Leu Ile Phe Ser Gly Asn Gln Gly
 195 200 205
 Arg Gly His Gln Arg Pro Leu Pro Pro Ser Glu Gly Leu Ser Pro Arg
 210 215 220
 30 Pro Pro Asn Ser Pro Ser Ile Ser Ile Pro Cys Met Gly Ser Lys Ala
 225 230 235 240
 Ser Ser Pro His Gly Leu Gly Ser Pro Leu Val Ala Ser Pro Arg Leu
 245 250 255
 Glu Lys Arg Leu Gly Gly Leu Ala Pro Gln Arg Gly Ser Arg Ile Ser
 260 265 270
 35 Val Leu Ser Ala Ser Pro Val Ser Asp Val Ser Tyr Met Phe Gly Ser
 275 280 285
 Ser Gln Ser Leu Leu His Ser Ser Asn Ser Ser His Gln Ser Ser Ser
 290 295 300
 40 Arg Ser Leu Glu Ser Pro Ala Asn Ser Ser Ser Ser Leu His Ser Leu
 305 310 315 320
 Gly Ser Val Ser Leu Cys Thr Arg Pro Ser Asp Phe Gln Ala Pro Arg
 325 330 335
 Asn Pro Thr Leu Thr Met Gly Gln Pro Arg Thr Pro His Ser Pro Pro
 340 345 350
 45 Leu Ala Lys Glu His Ala Ser Ile Cys Pro Pro Ser Ile Thr Asn Ser
 355 360 365
 Met Val Asp Ile Pro Ile Val Leu Ile Asn Gly Cys Pro Glu Pro Gly
 370 375 380
 Ser Ser Pro Pro Gln Arg Thr Pro Gly His Gln Asn Ser Val Gln Pro
 385 390 395 400
 50 Gly Ala Ala Ser Pro Ser Asn Pro Cys Pro Ala Thr Arg Ser Asn Ser
 405 410 415
 Gln Thr Leu Ser Asp Ala Pro Phe Thr Thr Cys Pro Glu Gly Pro Ala
 420 425 430
 Arg Asp Met Gln Pro Thr Met Lys Phe Val Met Asp Thr Ser Lys Tyr
 435 440 445

Trp Phe Lys Pro Asn Ile Thr Arg Glu Gln Ala Ile Glu Leu Leu Arg
 450 455 460
 Lys Glu Glu Pro Gly Ala Phe Val Ile Arg Asp Ser Ser Ser Tyr Arg
 465 470 475 480
 Gly Ser Phe Gly Leu Ala Leu Lys Val Gln Glu Val Pro Ala Ser Ala
 485 490 495
 Gln Asn Arg Pro Gly Glu Asp Ser Asn Asp Leu Ile Arg His Phe Leu
 500 505 510
 Ile Glu Ser Ser Ala Lys Gly Val His Leu Lys Gly Ala Asp Glu Glu
 515 520 525
 Pro Tyr Phe Gly Ser Leu Ser Ala Phe Val Cys Gln His Ser Ile Met
 530 535 540
 Ala Leu Ala Leu Pro Cys Lys Leu Thr Ile Pro Gln Arg Glu Leu Gly
 545 550 555 560
 Gly Ala Asp Gly Ala Ser Asp Ser Thr Asp Ser Pro Ala Ser Cys Gln
 565 570 575
 Lys Lys Ser Ala Gly Cys His Thr Leu Tyr Leu Ser Ser Val Ser Val
 580 585 590
 Glu Thr Leu Thr Gly Ala Leu Ala Val Gln Lys Ala Ile Ser Thr Thr
 595 600 605
 Phe Glu Arg Asp Ile Leu Pro Thr Pro Thr Val Val His Phe Glu Val
 610 615 620
 Thr Glu Gln Gly Ile Thr Leu Thr Asp Val Gln Arg Lys Val Phe Phe
 625 630 635 640
 Arg Arg His Tyr Pro Leu Thr Thr Leu Arg Phe Cys Gly Met Asp Pro
 645 650 655
 Glu Gln Arg Lys Trp Gln Lys Tyr Cys Lys Pro Ser Trp Ile Phe Gly
 660 665 670
 Phe Val Ala Lys Ser Gln Thr Glu Pro Gln Glu Asn Val Cys His Leu
 675 680 685
 Phe Ala Glu Tyr Asp Met Val Gln Pro Ala Ser Gln Val Ile Gly Leu
 690 695 700
 Val Thr Ala Leu Leu Gln Asp Ala Glu Arg Met
 705 710 715

35 <210> 99

<211> 35

<212> DNA

40

<213> Artificial sequence

45 <220>

<223> PCR primer

<400> 99

ccatatataa aaccactgtc ctgtcctttg tggct

35

50 <210> 100

<211> 26

<212> DNA

55

<213> Artificial sequence

5 <220>

 <223> PCR primer

 <400> 100

 cccccacatctg totgtctata ttgtc 26

10 <210> 101

 <211> 22

15 <212> DNA

 <213> Artificial sequence

20 <220>

 <223> PCR primer

 <400> 101

 tgcctacgct gacgactatg tg 22

25 <210> 102

 <211> 25

30 <212> DNA

 <213> Artificial sequence

35 <220>

 <223> PCR primer

 <400> 102

 tttggttttc tacaactggt gctat 25

40 <210> 103

 <211> 19

45 <212> DNA

 <213> Artificial sequence

50 <220>

 <223> PCR primer

 <400> 103

 gggctccaca caccagatg 19

55

<210> 104
 <211> 21
 5 <212> DNA
 <213> Artificial sequence
 10
 <220>
 <223> PCR primer
 15 <400> 104
 acgctctgag caccctctac a 21
 <210> 105
 <211> 31
 20 <212> DNA
 <213> Artificial sequence
 25
 <220>
 <223> PCR primer
 30 <400> 105
 tgtcacaggg actgaaaacc tctcctcatg t 31
 <210> 106
 <211> 17
 35 <212> DNA
 <213> Artificial sequence
 40
 <220>
 <223> PCR primer
 45 <400> 106
 cccaaggcca cgagctt 17
 <210> 107
 <211> 24
 50 <212> DNA
 <213> Artificial sequence
 55

<220>
 <223> PCR primer
 5 <400> 107
 tgttgctctc ttaacgaatc gaaa 24
 <210> 108
 10 <211> 29
 <212> DNA
 <213> Artificial sequence
 15
 <220>
 <223> PCR primer
 20 <400> 108
 ctggtcaaac aaactctctg aacccctcc 29
 <210> 109
 <211> 20
 25 <212> DNA
 <213> Artificial sequence
 30
 <220>
 <223> PCR primer
 35 <400> 109
 tggtgaggaa aagcggacat 20
 <210> 110
 <211> 21
 40 <212> DNA
 <213> Artificial sequence
 45
 <220>
 <223> PCR primer
 50 <400> 110
 ctggcttgga ggacagtgaa g 21
 <210> 111
 <211> 24
 55

	<212> DNA	
5	<213> Artificial sequence	
	<220>	
10	<223> PCR primer	
	<400> 111	24
	ccaagccctc cccatcccat gtat	
	<210> 112	
15	<211> 21	
	<212> DNA	
	<213> Artificial sequence	
20	<220>	
	<223> PCR primer	
25	<400> 112	21
	gaggtgtcgt accgcgttct a	
	<210> 113	
30	<211> 21	
	<212> DNA	
	<213> Artificial sequence	
35	<220>	
	<223> PCR primer	
40	<400> 113	21
	ccgtttctget cttccctgtc t	
	<210> 114	
45	<211> 23	
	<212> DNA	
	<213> Artificial sequence	
50	<220>	
	<223> PCR primer	
55	<400> 114	23
	ccagaccggc ttcaactgacc tgc	

	<210> 115	
5	<211> 20	
	<212> DNA	
	<213> Artificial sequence	
10		
	<220>	
	<223> PCR primer	
15	<400> 115	
	cgccgtgtact tcagcatgga	20
	<210> 116	
20	<211> 18	
	<212> DNA	
	<213> Artificial sequence	
25		
	<220>	
	<223> PCR primer	
30	<400> 116	
	gcgggttcagc tgggtggaa	18
	<210> 117	
35	<211> 25	
	<212> DNA	
	<213> Artificial sequence	
40		
	<220>	
	<223> PCR primer	
45	<400> 117	
	accccgaggc atcaccacaa atcat	25
	<210> 118	
50	<211> 23	
	<212> DNA	
	<213> Artificial sequence	
55		

<220>
 <223> PCR primer
 5 <400> 118
 agttctgcct ctctgacaac cat 23
 <210> 119
 10 <211> 23
 <212> DNA
 <213> Artificial sequence
 15
 <220>
 <223> PCR primer
 20 <400> 119
 taggctcaga gtcagaccca aac 23
 <210> 120
 <211> 21
 25 <212> DNA
 <213> Artificial sequence
 30
 <220>
 <223> PCR primer
 35 <400> 120
 ccctcgtggg cttgtgctcg g 21
 <210> 121
 <211> 21
 40 <212> DNA
 <213> Artificial sequence
 45
 <220>
 <223> PCR primer
 50 <400> 121
 aagccgccag ttcattctttt t 21
 <210> 122
 <211> 25
 55

	<212> DNA	
5	<213> Artificial sequence	
	<220>	
10	<223> PCR primer	
	<400> 122	
	cttgtggttc aagtcaaattg ttcag	25
	<210> 123	
15	<211> 21	
	<212> DNA	
	<213> Artificial sequence	
20	<220>	
	<223> PCR primer	
25	<400> 123	
	tctgcctgcg ctctcgtcgg t	21
	<210> 124	
30	<211> 18	
	<212> DNA	
	<213> Artificial sequence	
35	<220>	
	<223> PCR primer	
40	<400> 124	
	gggctgggca cctgactt	18
	<210> 125	
	<211> 20	
45	<212> DNA	
	<213> Artificial sequence	
50	<220>	
	<223> PCR primer	
55	<400> 125	
	cccaacaagg gtcccagact	20

	<210> 126	
5	<211> 17	
	<212> DNA	
	<213> Artificial sequence	
10		
	<220>	
	<223> PCR primer	
15	<400> 126	
	cggcgcattg agcggcg	17
	<210> 127	
	<211> 20	
20	<212> DNA	
	<213> Artificial sequence	
25		
	<220>	
	<223> PCR primer	
	<400> 127	
30	cccaaggac ttcgtgaatg	20
	<210> 128	
	<211> 21	
35	<212> DNA	
	<213> Artificial sequence	
40		
	<220>	
	<223> PCR primer	
	<400> 128	
45	ggcgatccct gatgacaagt a	21
	<210> 129	
	<211> 29	
50	<212> DNA	
	<213> Artificial sequence	
55		

<220>
 <223> PCR primer#
 5 <400> 129
 agcaccaact gtgaaccagg tacaatggc 29
 <210> 130
 10 <211> 19
 <212> DNA
 <213> Artificial sequence
 15
 <220>
 <223> PCR primer
 20 <400> 130
 gagggaggct ctgctttgg 19
 <210> 131
 <211> 21
 25 <212> DNA
 <213> Artificial sequence
 30
 <220>
 <223> PCR primer
 35 <400> 131
 tcacaactag cgggtgagga g 21
 <210> 132
 <211> 21
 40 <212> DNA
 <213> Artificial sequence
 45
 <220>
 <223> PCR primer
 50 <400> 132
 tgcagaggaa cggcgtgagc g 21
 <210> 133
 <211> 22
 55

<212> DNA
 <213> Artificial sequence
 5

 <220>
 10 <223> PCR primer
 <400> 133
 tgagggtttcc toccaaatcg ta 22

 <210> 134
 15 <211> 22
 <212> DNA
 <213> Artificial sequence
 20

 <220>
 25 <223> PCR primer
 <400> 134
 cagctcaagg gaagctgtca tc 22

 <210> 135
 30 <211> 24
 <212> DNA
 <213> Artificial sequence
 35

 <220>
 40 <223> PCR primer
 <400> 135
 cccccacatg ttccccaaga tgct 24

 <210> 136
 45 <211> 21
 <212> DNA
 <213> Artificial sequence
 50

 <220>
 55 <223> PCR primer
 <400> 136
 ggaggcgcta aaggtctacg t 21

	<210>	137	
5	<211>	21	
	<212>	DNA	
	<213>	Artificial sequence	
10	<220>		
	<223>	PCR primer	
15	<400>	137	
		tgatgcttcg caggtcagta a	21
	<210>	138	
20	<211>	26	
	<212>	DNA	
	<213>	Artificial sequence	
25	<220>		
	<223>	PCR primer	
30	<400>	138	
		ctcctgcccc tcctaaagct gaagcc	26
	<210>	139	
35	<211>	17	
	<212>	DNA	
	<213>	Artificial sequence	
40	<220>		
	<223>	PCR primer	
45	<400>	139	
		ggacgcgtgg gcttttc	17
	<210>	140	
	<211>	20	
50	<212>	DNA	
	<213>	Artificial sequence	
55			

<220>
 <223> PCR primer
 5 <400> 140
 tgtggctgtg gacacctttc 20
 <210> 141
 10 <211> 25
 <212> DNA
 <213> Artificial sequence
 15
 <220>
 <223> PCR primer
 20 <400> 141
 ccacaagctg aaggcagaca aggcc 25
 <210> 142
 <211> 20
 25 <212> DNA
 <213> Artificial sequence
 30
 <220>
 <223> PCR primer
 35 <400> 142
 gcggattctc atggaacaca 20
 <210> 143
 <211> 20
 40 <212> DNA
 <213> Artificial sequence
 45
 <220>
 <223> PCR primer
 50 <400> 143
 ggtcagccag gagcttcttg 20
 <210> 144
 <211> 23
 55

<212> DNA
 <213> Artificial sequence
 5
 <220>
 <223> PCR primer
 10 <400> 144
 accaccttgc gcaggttgtc cag 23
 <210> 145
 15 <211> 18
 <212> DNA
 <213> Artificial sequence
 20
 <220>
 <223> PCR primer
 25 <400> 145
 cgcatgcacg acctgaac 18
 <210> 146
 <211> 23
 30 <212> DNA
 <213> Artificial sequence
 35
 <220>
 <223> PCR primer
 40 <400> 146
 gtctcgatct tggacagctt ctg 23
 <210> 147
 <211> 22
 45 <212> DNA
 <213> Artificial sequence
 50
 <220>
 <223> PCR primer
 55 <400> 147
 acactgtcca cacggcccga gg 22

	<210>	148	
5	<211>	21	
	<212>	DNA	
	<213>	Artificial sequence	
10	<220>		
	<223>	PCR primer	
15	<400>	148	
		ctgggcagaa tggaaggatc t	21
	<210>	149	
20	<211>	22	
	<212>	DNA	
	<213>	Artificial sequence	
25	<220>		
	<223>	PCR primer	
30	<400>	149	
		gggactctag cagacccaca ct	22
	<210>	150	
	<211>	22	
35	<212>	DNA	
	<213>	Artificial sequence	
40	<220>		
	<223>	PCR primer	
45	<400>	150	
		cacccacctg gattocctgt tc	22
	<210>	151	
	<211>	23	
50	<212>	DNA	
	<213>	Artificial sequence	
55			

	<220>		
	<223> PCR primer		
5	<400> 151		
	ccttcagaca ggcgtagatg atg	23	
	<210> 152		
10	<211> 29		
	<212> DNA		
	<213> Artificial sequence		
15			
	<220>		
	<223> PCR primer		
20	<400> 152		
	gggtattatt tctttattag gtgccactt	29	
	<210> 153		
	<211> 30		
25	<212> DNA		
	<213> Artificial sequence		
30			
	<220>		
	<223> PCR primer		
35	<400> 153		
	ttccctaagg ctttcagtac ccaggatctg	30	
	<210> 154		
	<211> 18		
40	<212> DNA		
	<213> Artificial sequence		
45			
	<220>		
	<223> PCR primer		
50	<400> 154		
	ccagcttggc cctttcct	18	
	<210> 155		
	<211> 23		
55			

	<212> DNA	
5	<213> Artificial sequence	
	<220>	
10	<223> PCR primer	
	<400> 155	23
	gaatgggtcg cttttgttct tag	
15	<210> 156	
	<211> 22	
	<212> DNA	
20	<213> Artificial sequence	
	<220>	
25	<223> PCR primer	
	<400> 156	22
	tcacggacct cagcctgcc ct	
30	<210> 157	
	<211> 21	
	<212> DNA	
35	<213> Artificial sequence	
	<220>	
40	<223> PCR primer	
	<400> 157	21
	tggtgaagggt gtcagccatg t	
45	<210> 158	
	<211> 21	
	<212> DNA	
50	<213> Artificial sequence	
55		

	<220>		
5	<223> PCR primer		
	<400> 158		
	tcagagtgca gcaatggctt t	21	
	<210> 159		
10	<211> 20		
	<212> DNA		
	<213> Artificial sequence		
15			
	<220>		
	<223> PCR primer		
20	<400> 159		
	acctccttcc ccagctcccc	20	
	<210> 160		
	<211> 24		
25	<212> DNA		
	<213> Artificial sequence		
30			
	<220>		
	<223> PCR primer		
35	<400> 160		
	ggcaacatct tacttgtcct ttga	24	
	<210> 161		
	<211> 25		
40	<212> DNA		
	<213> Artificial sequence		
45			
	<220>		
	<223> PCR primer		
50	<400> 161		
	ccaaggaagc acagacaact atttc	25	
	<210> 162		
	<211> 30		
55			

<212> DNA
 <213> Artificial sequence
 5
 <220>
 10 <223> PCR primer
 <400> 162
 tcctccctat ccattggcact aaaccacttc 30
 <210> 163
 15 <211> 19
 <212> DNA
 <213> Artificial sequence
 20
 <220>
 25 <223> PCR primer
 <400> 163
 tgggcaaggc ctcctatct 19
 <210> 164
 30 <211> 21
 <212> DNA
 <213> Artificial sequence
 35
 <220>
 40 <223> PCR primer
 <400> 164
 gttacccttg gcagacgtat g 21
 <210> 165
 45 <211> 31
 <212> DNA
 <213> Artificial sequence
 50
 <220>
 55 <223> PCR primer
 <400> 165
 tgcctctgag tctgaatctc ccaaagagag a 31

5 <210> 166
 <211> 31
 <212> DNA
 <213> Artificial sequence
 10
 <220>
 <223> PCR primer
 15 <400> 166
 gagtagttat gtgattatct cagctcttga c 31
 <210> 167
 20 <211> 21
 <212> DNA
 <213> Artificial sequence
 25
 <220>
 <223> PCR primer
 30 <400> 167
 tcaaattgtg tccccgagtc t 21
 <210> 168
 <211> 34
 35 <212> DNA
 <213> Artificial sequence
 40
 <220>
 <223> PCR primer
 <400> 168
 45 cagaaattcg gaagacagaa ctattgtcat gcct 34
 <210> 169
 <211> 27
 50 <212> DNA
 <213> Artificial sequence
 55

<220>
 5 <223> PCR primer
 <400> 169
 gattagtaac ccatagcagt tgaaggt 27
 <210> 170
 10 <211> 26
 <212> DNA
 <213> Artificial sequence
 15
 <220>
 20 <223> PCR primer
 <400> 170
 atttactgac ggtggtctga acatac 26
 <210> 171
 25 <211> 31
 <212> DNA
 <213> Artificial sequence
 30
 <220>
 35 <223> PCR primer
 <400> 171
 tgacagactc caaatcaca gcacagtcaa c 31
 <210> 172
 40 <211> 25
 <212> DNA
 <213> Artificial sequence
 45
 <220>
 50 <223> PCR primer
 <400> 172
 tgatggtttg gaggaaagtt tatatt 25
 <210> 173
 55 <211> 24

<212> DNA
 <213> Artificial sequence
 5

 <220>
 10
 <223> PCR primer
 <400> 173
 tttggttggg tcttttagagg aatc 24
 <210> 174
 15
 <211> 24
 <212> DNA
 <213> Artificial sequence
 20

 <220>
 25
 <223> PCR primer
 <400> 174
 tgccaaccat gcatcaggtg gcc 24
 <210> 175
 30
 <211> 20
 <212> DNA
 <213> Artificial sequence
 35

 <220>
 40
 <223> PCR primer
 <400> 175
 cagctcacct ggcaacttca 20
 <210> 176
 45
 <211> 20
 <212> DNA
 <213> Artificial sequence
 50

 <220>
 55
 <223> PCR primer
 <400> 176
 cctgattttc ccagcgatgt 20

	<210>	177	
5	<211>	19	
	<212>	DNA	
	<213>	Artificial sequence	
10	<220>		
	<223>	PCR primer	
15	<400>	177	
		cgccgctccc ggttotgct	19
	<210>	178	
20	<211>	20	
	<212>	DNA	
	<213>	Artificial sequence	
25	<220>		
	<223>	PCR primer	
30	<400>	178	
		tggccaagcg taagctgatt	20
	<210>	179	
	<211>	21	
35	<212>	DNA	
	<213>	Artificial sequence	
40	<220>		
	<223>	PCR primer	
45	<400>	179	
		gctgcagtga tcggatcatc t	21
	<210>	180	
	<211>	22	
50	<212>	DNA	
	<213>	Artificial Sequence	
55			

<220>
 <223> MLLT6
 5 <400> 180
 caccatggag cccatcgtgc tg 22
 <210> 181
 10 <211> 19
 <212> DNA
 <213> Artificial Sequence
 15
 <220>
 <223> MLLT6 for
 20 <400> 181
 atccccgagg tgcaatttg 19
 <210> 182
 <211> 21
 25 <212> DNA
 <213> Artificial Sequence
 30
 <220>
 <223> MLLT6 rev
 35 <400> 182
 agcgatcatg aggcacgtac t 21
 <210> 183
 <211> 29
 40 <212> DNA
 <213> Artificial Sequence
 45
 <220>
 <223> ZNF144
 50 <400> 183
 cctgccagag ataggagacc cagacagct 29
 <210> 184
 <211> 19
 55

<212> DNA

<213> Artificial Sequence

5

<220>

<223> ZNF144 for

<400> 184

atccccctga gcattttca

19

<210> 185

<211> 19

<212> DNA

<213> Artificial Sequence

20

<220>

<223> ZNF144 rev

<400> 185

cagcctctgg tcccaccat

19

<210> 186

<211> 28

<212> DNA

<213> Artificial Sequence

35

<220>

<223> PIP5K2B

<400> 186

tgatcatcaa ttccaaacct ctcccgaa

28

<210> 187

<211> 19

<212> DNA

<213> Artificial Sequence

50

<220>

<223> PIP5K2B for

<400> 187

ccccatggtg ttccgaaac

19

55

5 <210> 188
 <211> 19
 <212> DNA
 <213> Artificial Sequence
 10
 <220>
 <223> PIP5K2B rev
 15 <400> 188
 tgccaggagc ctccatacc 19
 <210> 189
 20 <211> 29
 <212> DNA
 <213> Artificial Sequence
 25
 <220>
 <223> TEM7
 30 <400> 189
 cagccttcta aaacacaatg tattcatgt 29
 <210> 190
 <211> 29
 35 <212> DNA
 <213> Artificial Sequence
 40
 <220>
 <223> TEM7 for
 45 <400> 190
 cctgaactta atggtagaat tcaaagatc 29
 <210> 191
 <211> 27
 50 <212> DNA
 <213> Artificial Sequence
 55

<220>
 5 <223> TEM7 rev
 <400> 191
 tattaacact gagaatccat gcagaga 27
 <210> 192
 10 <211> 35
 <212> DNA
 <213> Artificial Sequence
 15
 <220>
 20 <223> ZNFN1A3
 <400> 192
 tatctggtct cagggattgc tcctatgtat tcagc 35
 <210> 193
 25 <211> 20
 <212> DNA
 <213> Artificial Sequence
 30
 <220>
 35 <223> ZNFN1A3 for
 <400> 193
 cacagagccc tgctgaagtg 20
 <210> 194
 40 <211> 23
 <212> DNA
 <213> Artificial Sequence
 45
 <220>
 50 <223> ZNFN1A3 rev
 <400> 194
 gcgaggtcat tggtttttag aaa 23
 <210> 195
 <211> 22
 55

<212> DNA
 <213> Artificial Sequence
 5
 <220>
 10 <223> WIRE
 <400> 195
 ctgtgatccg aaatggtgcc ag 22
 <210> 196
 15 <211> 20
 <212> DNA
 <213> Artificial Sequence
 20
 <220>
 25 <223> WIRE for
 <400> 196
 ccgtctccac atccaaacct 20
 <210> 197
 30 <211> 20
 <212> DNA
 <213> Artificial Sequence
 35
 <220>
 40 <223> WIRE rev
 <400> 197
 acccatgcat tcggtatggt 20
 <210> 198
 45 <211> 21
 <212> DNA
 <213> Artificial Sequence
 50
 <220>
 55 <223> PSMB3
 <400> 198
 agtggcacct gcgccgaaca a 21

	<210>	199	
5	<211>	21	
	<212>	DNA	
	<213>	Artificial Sequence	
10	<220>		
	<223>	PSMB3 for	
15	<400>	199	
		ccccatggtg actgatgact t	21
	<210>	200	
20	<211>	21	
	<212>	DNA	
	<213>	Artificial Sequence	
25	<220>		
	<223>	PSMB3 rev	
30	<400>	200	
		ccagagggac tcacacattc c	21
	<210>	201	
	<211>	29	
35	<212>	DNA	
	<213>	Artificial Sequence	
40	<220>		
	<223>	MGC9753	
45	<400>	201	
		ccagaaactt tccatcccaa aggcagtct	29
	<210>	202	
	<211>	21	
50	<212>	DNA	
	<213>	Artificial Sequence	
55	<220>		

<220>
 <223> MGC9753 for
 5 <400> 202
 ctgccccaca ggaatagaat g 21
 <210> 203
 10 <211> 23
 <212> DNA
 <213> ARTIFICIAL SEQUENCE
 15
 <220>
 <223> MGC9753 rev
 20 <400> 203
 aaaaatccag tctgcttcaa cca 23
 <210> 204
 <211> 20
 25 <212> DNA
 <213> ARTIFICIAL SEQUENCE
 30
 <220>
 <223> ORMDL3
 <400> 204
 35 agctgccccca gctccacgga 20
 <210> 205
 <211> 21
 40 <212> DNA
 <213> ARTIFICIAL SEQUENCE
 45
 <220>
 <223> ORMDL3 for
 <400> 205
 50 tccctgatga gcgtgcttat c 21
 <210> 206
 <211> 28
 55

<212> DNA

5 <213> ARTIFICIAL SEQUENCE

<220>

10 <223> ORMDL3 rev
<400> 206
tctcagtact tattgattcc aaaaatcc 28

<210> 207

15 <211> 25

<212> DNA

<213> ARTIFICIAL SEQUENCE

20 <220>

25 <223> MGC15482
<400> 207
tccagtggaa gcaaccccag tgttc 25

<210> 208

30 <211> 25

<212> DNA

<213> ARTIFICIAL SEQUENCE

35 <220>

40 <223> MGC15482 for
<400> 208
cacttctaga gctaccgtgg agtct 25

<210> 209

45 <211> 22

<212> DNA

<213> ARTIFICIAL SEQUENCE

50 <220>

55 <223> MGC15482 rev
<400> 209
ccctcacttt gtaacccttg ct 22

<210> 210
 5 <211> 20
 <212> DNA
 <213> ARTIFICIAL SEQUENCE
 10
 <220>
 <223> PPP1R1B
 15 <400> 210
 cagcgtggcg caacaaccca 20
 <210> 211
 20 <211> 21
 <212> DNA
 <213> ARTIFICIAL SEQUENCE
 25
 <220>
 <223> PPP1R1B for
 30 <400> 211
 gggattgttt cgccacacat a 21
 <210> 212
 <211> 20
 35 <212> DNA
 <213> ARTIFICIAL SEQUENCE
 40
 <220>
 <223> PPP1R1B rev
 45 <400> 212
 ccgatgttaa ggcccatagc 20
 <210> 213
 <211> 27
 50 <212> DNA
 <213> ARTIFICIAL SEQUENCE
 55

<220>
 <223> MGC14832
 5 <400> 213
 taaaatgtcc ggccaacatg agttccc 27
 <210> 214
 10 <211> 17
 <212> DNA
 <213> ARTIFICIAL SEQUENCE
 15
 <220>
 <223> MGC14832 for
 20 <400> 214
 cgcagtgccct ggcacat 17
 <210> 215
 25 <211> 20
 <212> DNA
 <213> ARTIFICIAL SEQUENCE
 30
 <220>
 <223> MGC14832 rev
 35 <400> 215
 gacacccccct gacctatgga 20
 <210> 216
 40 <211> 25
 <212> DNA
 <213> ARTIFICIAL SEQUENCE
 45
 <220>
 <223> LOC51242
 50 <400> 216
 cagtgacctc tcccgttccc ttgga 25
 <210> 217
 <211> 20
 55

<212> DNA

5 <213> ARTIFICIAL SEQUENCE

<220>

10 <223> LOC51242 for
<400> 217
tgggtccctg tgtcctcttc 20

<210> 218

15 <211> 20

<212> DNA

<213> ARTIFICIAL SEQUENCE

20 <220>

25 <223> LOC51242 for
<400> 218
agggtcagga gggagaaaac 20

<210> 219

30 <211> 26

<212> DNA

<213> ARTIFICIAL SEQUENCE

35 <220>

40 <223> FLJ20291
<400> 219
ccagtgccca cccgttaaag agtcaa 26

<210> 220

45 <211> 24

<212> DNA

<213> ARTIFICIAL SEQUENCE

50 <220>

55 <223> FLJ20291 for
<400> 220
ttgtgggaca ctcagtaact ttgg 24

	<210>	221	
5	<211>	20	
	<212>	DNA	
	<213>	ARTIFICIAL SEQUENCE	
10	<220>		
	<223>	FLJ20291 rev	
15	<400>	221	
		acaagcactc ccaccgagat	20
	<210>	222	
20	<211>	24	
	<212>	DNA	
	<213>	ARTIFICIAL SEQUENCE	
25	<220>		
	<223>	PRO2521	
30	<400>	222	
		agtctgtcct cactgccatc gcc	24
	<210>	223	
	<211>	21	
35	<212>	DNA	
	<213>	ARTIFICIAL SEQUENCE	
40	<220>		
	<223>	PRO2521 for	
45	<400>	223	
		aagcctctgg gttttccctt t	21
	<210>	224	
	<211>	20	
50	<212>	DNA	
	<213>	ARTIFICIAL SEQUENCE	
55			

<220>
 <223> PRO2521 rev
 5 <400> 224
 cccactggtg acaggatggt 20
 <210> 225
 10 <211> 23
 <212> DNA
 <213> ARTIFICIAL SEQUENCE
 15
 <220>
 <223> LINK-GEFII
 20 <400> 225
 catctgacat ctttcccggtg gag 23
 <210> 226
 <211> 21
 25 <212> DNA
 <213> ARTIFICIAL SEQUENCE
 30
 <220>
 <223> LINK-GEFII for
 35 <400> 226
 ctttgacga tgtctcaacc a 21
 <210> 227
 <211> 18
 40 <212> DNA
 <213> ARTIFICIAL SEQUENCE
 45
 <220>
 <223> LINK-GEFII rev
 50 <400> 227
 tttcccggtg agcaggaa 18
 <210> 228
 <211> 26
 55

<212> DNA

<213> ARTIFICIAL SEQUENCE

5

<220>

<223> CTEN

<400> 228

10 ccgccgccta atatgcaaca ttaggg

26

<210> 229

<211> 23

15

<212> DNA

<213> ARTIFICIAL SEQUENCE

20

<220>

<223> CTEN for

<400> 229

25 cgagtattcc aaagctggta tcg

23

<210> 230

<211> 24

30

<212> DNA

<213> ARTIFICIAL SEQUENCE

35

<220>

<223> CTEN rev

<400> 230

40 atcacagaga gatggccctt atct

24

<210> 231

<211> 25

45

<212> DNA

<213> Artificial Sequence

50

<220>

<223> D17S946 forward primer

<400> 231

55 acagtctatc aagcagaaaa atcct

25

<210> 232
 5 <211> 16
 <212> DNA
 <213> Artificial Sequence
 10
 <220>
 <223> D17S946 reverse primer
 15 <400> 232
 tgccgtgcc gagaga 16
 <210> 233
 20 <211> 20
 <212> DNA
 <213> Artificial Sequence
 25
 <220>
 <223> D17S1181 forward primer
 30 <400> 233
 gacaacagag cgagactccc 20
 <210> 234
 35 <211> 20
 <212> DNA
 <213> Artificial Sequence
 40
 <220>
 <223> D17S1181 reverse primer
 45 <400> 234
 gccagcctg tcacttattc 20
 <210> 235
 50 <211> 18
 <212> DNA
 <213> Artificial Sequence
 55

<220>

<223> D17S2026 forward primer
5 <400> 235
tggtcattcg acaacgaa 18

<210> 236

10 <211> 18

<212> DNA

<213> Artificial Sequence

15

<220>

<223> D17S2026 reverse primer
20 <400> 236
cagcattgga tgcaatcc 18

<210> 237

<211> 20

25 <212> DNA

<213> Artificial Sequence

30

<220>

<223> D17S838 forward primer
35 <400> 237
ctocagaatc cagaccatga 20

<210> 238

<211> 20

40 <212> DNA

<213> Artificial Sequence

45

<220>

<223> D17S838 reverse primer
50 <400> 238
aggacagtgt gtagcccttc 20

<210> 239

<211> 20

55

<212> DNA

<213> Artificial Sequence

5

<220>

<223> D17S250 forward primer

<400> 239

ggaagaatca aatagacaat

20

<210> 240

<211> 24

<212> DNA

<213> Artificial Sequence

20

<220>

<223> D17S250 reverse primer

<400> 240

gctggccata tatatatatta aacc

24

<210> 241

<211> 23

<212> DNA

<213> Artificial Sequence

35

<220>

<223> D17S1818 forward primer

<400> 241

cataggtatg ttcagaaatg tga

23

<210> 242

<211> 18

<212> DNA

<213> Artificial Sequence

50

<220>

<223> D17S1818 reverse primer

<400> 242

tgctactgg aaaccaga

18

55

<210> 243
 5 <211> 23
 <212> DNA
 <213> Artificial Sequence
 10
 <220>
 <223> D17S614 forward primer
 15 <400> 243
 aaggggaagg ggctttcaaa gct 23
 <210> 244
 20 <211> 23
 <212> DNA
 <213> Artificial Sequence
 25
 <220>
 <223> D17S614 reverse primer
 30 <220>
 <221> misc_feature
 <222> (1)..(1)
 35 <223> n=a, c, g or t
 <400> 244
 nggaggttgc agtgagccaa gat 23
 40 <210> 245
 <211> 23
 <212> DNA
 45 <213> Artificial Sequence
 <220>
 50 <223> D17S2019 forward primer
 <400> 245
 caaaagctta tgatgctcaa acc 23
 55

<210> 246
 <211> 22
 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> D17S2019 reverse primer
 <400> 246
 ttgtttccct ttgactttct ga 22
 <210> 247
 <211> 25
 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> D17S608 forward primer
 <400> 247
 taggttcacc ttcattttc ttcag 25
 <210> 248
 <211> 24
 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> D17S608 reverse primer
 <220>
 <221> misc_feature
 <222> (17)..(17)
 <223> n=a, c, g or t
 <400> 248
 atctaatct ttataaact tata 24

<211> 20
 5 <212> DNA
 <213> Artificial Sequence

 10 <220>
 <223> D17S1655 forward primer
 <400> 249
 cggaccagag tgttccatgg 20
 15 <210> 250
 <211> 20
 <212> DNA
 20 <213> Artificial Sequence

 25 <220>
 <223> D17S1655 reverse primer
 <400> 250
 gcatacagca ccctctacct 20
 30 <210> 251
 <211> 25
 <212> DNA
 35 <213> Artificial Sequence

 40 <220>
 <223> D17S2147 forward primer
 <400> 251
 aggggagaat aaataaaatc tgtgg 25
 45 <210> 252
 <211> 22
 <212> DNA
 50 <213> Artificial Sequence

 55

<220>
 <223> D17S2147 reverse primer
 5 <400> 252
 caggagtgag acactctoca tg 22
 <210> 253
 10 <211> 22
 <212> DNA
 <213> Artificial Sequence
 15
 <220>
 <223> D17S754 forward primer
 20 <400> 253
 tggattoact gactcagcct gc 22
 <210> 254
 25 <211> 22
 <212> DNA
 <213> Artificial Sequence
 30
 <220>
 <223> D17S754 reverse primer
 35 <400> 254
 gcgtgtctgt ctccatgtgt gc 22
 <210> 255
 40 <211> 18
 <212> DNA
 <213> Artificial Sequence
 45
 <220>
 <223> D17S1814 forward primer
 50 <400> 255
 tccccaatga cggtgatg 18
 <210> 256
 <211> 20
 55

	<212> DNA	
5	<213> Artificial Sequence	
	<220>	
10	<223> D17S1814 reverse primer	
	<400> 256	
	ctggaggttg gcttgtggat	20
	<210> 257	
15	<211> 18	
	<212> DNA	
20	<213> Artificial Sequence	
	<220>	
25	<223> D17S2007 forward primer	
	<400> 257	
	ggtccacga atttgctg	18
	<210> 258	
30	<211> 20	
	<212> DNA	
35	<213> Artificial Sequence	
	<220>	
40	<223> D17S2007 reverse primer	
	<400> 258	
	ccacccagaa aaacaggaga	20
	<210> 259	
45	<211> 20	
	<212> DNA	
50	<213> Artificial Sequence	
	<220>	
55	<223> D17S1246 forward primer	
	<400> 259	
	tcgatetoot gaccttgtga	20

	<210>	260	
5	<211>	20	
	<212>	DNA	
	<213>	Artificial Sequence	
10	<220>		
	<223>	D17S1246 reverse primer	
15	<400>	260	
		ttgtcacccc attgcctttc	20
	<210>	261	
20	<211>	21	
	<212>	DNA	
	<213>	Artificial Sequence	
25	<220>		
	<223>	D17S1979 forward primer	
30	<400>	261	
		ccttgatag attcagctcc c	21
	<210>	262	
35	<211>	21	
	<212>	DNA	
	<213>	Artificial Sequence	
40	<220>		
	<223>	D17S1979 reverse primer	
45	<400>	262	
		cttgccctt ctcaatcctc c	21
	<210>	263	
50	<211>	25	
	<212>	DNA	
	<213>	Artificial Sequence	
55			

<220>
 <223> D17S1984 forward primer
 5 <400> 263
 ttaagcaagg ttttaattaa gctgc 25
 <210> 264
 10 <211> 21
 <212> DNA
 <213> Artificial Sequence
 15
 <220>
 <223> D17S1984 reverse primer
 20 <400> 264
 gattacagtg ctccctotoc c 21
 <210> 265
 25 <211> 22
 <212> DNA
 <213> Artificial Sequence
 30
 <220>
 <223> G11580 forward primer
 35 <400> 265
 ggttttaatt aagctgcatg gc 22
 <210> 266
 40 <211> 21
 <212> DNA
 <213> Artificial Sequence
 45
 <220>
 <223> G11580 reverse primer
 50 <400> 266
 gattacagtg ctccctctcc c 21
 <210> 267
 55 <211> 20

<212> DNA

<213> Artificial Sequence

5

<220>

<223> D17S1867 forward primer

<400> 267

agtttgacac tgaggcttg

20

<210> 268

<211> 20

<212> DNA

<213> Artificial Sequence

20

<220>

<223> D17S1867 reverse primer

<400> 268

tttagacttg gtaactgccg

20

<210> 269

<211> 24

<212> DNA

<213> Artificial Sequence

35

<220>

<223> D17S1788 forward primer

<400> 269

tgcagatgcc taagaacttt tcag

24

<210> 270

<211> 19

<212> DNA

<213> Artificial Sequence

50

<220>

<223> D17S1788 reverse primer

<400> 270

gccatgatct cccaaagcc

19

55

<210> 271
 5 <211> 18
 <212> DNA
 <213> Artificial Sequence
 10
 <220>
 <223> D17S1836 forward primer
 15 <400> 271
 tcgagggttat ggtgagcc 18
 <210> 272
 20 <211> 24
 <212> DNA
 <213> Artificial Sequence
 25
 <220>
 <223> D17S1836 reverse primer
 30 <400> 272
 aaactgtgtg tgtcaaagga tact 24
 <210> 273
 35 <211> 19
 <212> DNA
 <213> Artificial Sequence
 40
 <220>
 <223> D17S1787 forward primer
 45 <400> 273
 gctgatctga agccaatga 19
 <210> 274
 50 <211> 19
 <212> DNA
 <213> Artificial Sequence
 55

<220>
 5 <223> D17S1787 reverse primer
 <400> 274
 tacatgaagg catggtctg 19
 <210> 275
 10 <211> 23
 <212> DNA
 <213> Artificial Sequence
 15
 <220>
 20 <223> D17S1660 forward primer
 <400> 275
 ctaatataat cctgggcaca tgg 23
 <210> 276
 25 <211> 18
 <212> DNA
 <213> Artificial Sequence
 30
 <220>
 35 <223> D17S1660 reverse primer
 <400> 276
 gctgoggacc agacagat 18
 <210> 277
 40 <211> 22
 <212> DNA
 <213> Artificial Sequence
 45
 <220>
 50 <223> D17S2154 forward primer
 <400> 277
 gataaaaaca agcactggct cc 22
 <210> 278
 55 <211> 20

<212> DNA

<213> Artificial Sequence

5

<220>

<223> D17S2154 reverse primer

<400> 278

cccacggctt tcttgatcta

20

<210> 279

<211> 21

<212> DNA

<213> Artificial Sequence

20

<220>

<223> D17S1955 forward primer

<400> 279

tgtaatgtaa gccccatgag g

21

<210> 280

<211> 25

<212> DNA

<213> Artificial Sequence

35

<220>

<223> D17S1955 reverse primer

<400> 280

cactcaactc aacagtctaa aggtg

25

<210> 281

<211> 25

<212> DNA

<213> Artificial Sequence

50

<220>

<223> D17S2098 forward primer

<400> 281

gtgagttcaa gcatagtaat tatcc

25

55

5 <210> 282
 <211> 23
 <212> DNA
 <213> Artificial Sequence
 10
 <220>
 <223> D17S2098 reverse primer
 15 <400> 282
 attcagcctc agttcactgc ttc 23
 <210> 283
 20 <211> 20
 <212> DNA
 <213> Artificial Sequence
 25
 <220>
 <223> D17S518 forward primer
 30 <400> 283
 gatccagtgg agactcagag 20
 <210> 284
 35 <211> 20
 <212> DNA
 <213> Artificial Sequence
 40
 <220>
 <223> D17S518 reverse primer
 45 <400> 284
 tagtctctgg gacacccaga 20
 <210> 285
 <211> 25
 50 <212> DNA
 <213> Artificial Sequence
 55

<220>
 5 <223> D17S518 forward primer
 <400> 285
 attcctgagt gtctaccctg ttgag 25
 <210> 286
 10 <211> 17
 <212> DNA
 <213> Artificial Sequence
 15
 <220>
 20 <223> D17S518 reverse primer
 <400> 286
 actgactgcg ccaactgc 17
 <210> 287
 25 <211> 20
 <212> DNA
 <213> Artificial Sequence
 30
 <220>
 35 <223> D11S4358 forward primer
 <400> 287
 tcgagaagga caaaatcacc 20
 <210> 288
 40 <211> 20
 <212> DNA
 <213> Artificial Sequence
 45
 <220>
 50 <223> D11S4358 reverse primer
 <400> 288
 gaacagggtt agtccattcg 20
 <210> 289
 55 <211> 19

<212> DNA
 <213> Artificial Sequence
 5
 <220>
 10 <223> D17S964 forward primer
 <400> 289
 gttcttttccct cttgtgggg 19
 <210> 290
 15 <211> 19
 <212> DNA
 <213> Artificial Sequence
 20
 <220>
 25 <223> D17S964 reverse primer
 <400> 290
 agtcagctga gattgtgcc 19
 <210> 291
 30 <211> 20
 <212> DNA
 <213> Artificial Sequence
 35
 <220>
 40 <223> D19S1091 forward primer
 <400> 291
 caagccaaga catcccagtt 20
 <210> 292
 45 <211> 20
 <212> DNA
 <213> Artificial Sequence
 50
 <220>
 55 <223> D19S1091 reverse primer
 <400> 292
 cccacacac agctcatatg 20

5 <210> 293
 <211> 22
 <212> DNA
 <213> Artificial Sequence
 10
 <220>
 <223> D17S1179 forward primer
 15 <400> 293
 ttttctctct cattccattg gg 22
 <210> 294
 20 <211> 20
 <212> DNA
 <213> Artificial Sequence
 25
 <220>
 <223> D17S1179 reverse primer
 30 <400> 294
 gcaacagagg gagactccaa 20
 <210> 295
 35 <211> 19
 <212> DNA
 <213> Artificial Sequence
 40
 <220>
 <223> D10S2160 forward primer
 45 <400> 295
 tcccatcccg taagacctc 19
 <210> 296
 <211> 25
 50 <212> DNA
 <213> Artificial Sequence
 55

<220>
 <223> D10S2160 reverse primer
 5 <400> 296
 tatggagtac ctactctatg ccagg 25
 <210> 297
 10 <211> 20
 <212> DNA
 <213> Artificial Sequence
 15
 <220>
 <223> D17S1230 forward primer
 20 <400> 297
 attcaaagct ggatcccttt 20
 <210> 298
 25 <211> 20
 <212> DNA
 <213> Artificial Sequence
 30
 <220>
 <223> D17S1230 reverse primer
 35 <400> 298
 agctgtgaca aatgcctgta 20
 <210> 299
 40 <211> 20
 <212> DNA
 <213> Artificial Sequence
 45
 <220>
 <223> D17S1338 forward primer
 50 <400> 299
 tcacctgaga ttgggagacc 20
 <210> 300
 55 <211> 18

	<212> DNA	
5	<213> Artificial Sequence	
	<220>	
10	<223> D17S1338 reverse primer	
	<400> 300	
	aagatggggc aggaatgg	18
	<210> 301	
15	<211> 19	
	<212> DNA	
	<213> Artificial Sequence	
20	<220>	
	<223> D17S2011 forward primer	
25	<400> 301	
	tcactgtcct ccaagccag	19
	<210> 302	
30	<211> 20	
	<212> DNA	
	<213> Artificial Sequence	
35	<220>	
	<223> D17S2011 reverse primer	
40	<400> 302	
	aaacaccaca ctctcccctg	20
	<210> 303	
45	<211> 20	
	<212> DNA	
	<213> Artificial Sequence	
50	<220>	
	<223> D17S2011 forward primer	
55	<400> 303	
	ttctttgggt tcccgtagcc	20

<210> 304

5 <211> 20

<212> DNA

<213> Artificial Sequence

10

<220>

<223> D17S2011 reverse primer

<400> 304

ggggcagacg acttctcctt

20

<210> 305

20 **<211>** 23

<212> DNA

<213> Artificial Sequence

25

<220>

<223> D17S2038 forward primer

30 **<400> 305**

ggggatacaa cctttaaagt tcc

23

<210> 306

<211> 25

<212> DNA

<213> Artificial Sequence

40

<220>

<223> D17S2038 reverse primer

<400> 306

attcacctaa tgaggattct tcttt

25

<210> 307

<211> 24

<212> DNA

<213> Artificial Sequence

55

<220>

<223> D17S2091 forward primer

<400> 307

gctgaaatag ccattcttgag ctac

24

<210> 308

<211> 23

<212> DNA

<213> Artificial Sequence

<220>

<223> D17S2091 reverse primer

<400> 308

tccgcatcct ttttaagagg cac

23

<210> 309

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> D17S649 forward primer

<400> 309

ctttcactct ttcagctgaa gagg

24

<210> 310

<211> 25

<212> DNA

<213> Artificial Sequence

<220>

<223> D17S649 reverse primer

<400> 310

tgacgtgcta tttcctgttt tgtct

25

<210> 311

<211> 18

<212> DNA

<213> Artificial Sequence

<220>

No.	Doccode	Number of pages
1	1001	6
2	1002	1
3	DESC	4
4	CLMS	1
5	DRAW	1
6	ABST	1
7	PRIODOC	8

<220>

<223> D17S1190 reverse primer

<400> 312

caacacacta ccccaagga

Claims

1. A method for the prediction, diagnosis or prognosis of malignant neoplasia by the detection of at least 2 markers **characterized in that** the markers are genes and fragments thereof or genomic nucleic acid sequences that are located on one chromosomal region which is altered in malignant neoplasia.
2. A method for the prediction, diagnosis or prognosis of malignant neoplasia by the detection of at least 2 markers **characterized in that** the markers are:
 - a) genes that are located on one or more chromosomal region(s) which is/are altered in malignant neoplasia; and
 - b)
 - i) receptor and ligand; or
 - ii) members of the same signal transduction pathway; or
 - iii) members of synergistic signal transduction pathways; or
 - iv) members of antagonistic signal transduction pathways; or
 - v) transcription factor and transcription factor binding site.
3. The method of claim 1 or 2 wherein the malignant neoplasia is breast cancer, ovarian cancer, gastric cancer, colon cancer, esophageal cancer, mesenchymal cancer, bladder cancer or non-small cell lung cancer.
4. The method of claim 1 or 2 wherein at least one chromosomal region is defined as the cytogenetic region: 1p13, 1q32, 3p21-p24, 5p13-p14, 8q23-q²⁴, 1 1q13, 12q13, 17q12-q24 or 20q13.

5. The method of claim 1 or 2 wherein at least chromosomal region is defined as the cytogenetic region 17q1 1.2-21.3 and the malignant neoplasia is breast cancer, ovarian cancer, gastric cancer, colon cancer, esophageal cancer, mesenchymal cancer, bladder cancer or non-small cell lung cancer.

6. The method of claim 1 or 2 wherein at least one chromosomal region is defined as the cytogenetic region 3p21-24 and the malignant neoplasia is breast cancer, ovarian cancer, gastric cancer, colon cancer, esophageal cancer, mesenchymal cancer, bladder cancer or non-small cell lung cancer.

7. The method of claim 1 or 2 wherein at least one chromosomal region is defined as the cytogenetic region 12q13 and the malignant neoplasia is breast cancer, ovarian cancer, gastric cancer, colon cancer, esophageal cancer, mesenchymal cancer, bladder cancer or non-small cell lung cancer.

8. A method for the prediction, diagnosis or prognosis of malignant neoplasia by the detection of at least one marker whereby the marker is a VNTR, SNP, RFLP or STS **characterized in that** the marker is located on one chromosomal region which is altered in malignant neoplasia due to amplification and the marker is detected in a cancerous and a non-cancerous tissue or biological sample of the same individual.

9. The method of claim 8 wherein the marker is selected from the group consisting of the VNTRs:

D17S946, D17S1181, D17S2026, D17S838, D17S250, D17S1818, D17S614, D17S2019, D17S608, D17S1655, D17S2147, D17S754, D17S1814, D17S2007, D17S1246, D17S1979, D17S1984, D17S1984, D17S1867, D17S1788, D17S1836, D17S1787, D17S1660, D17S2154, D17S1955, D17S2098, D17S518, D17S1851, D11S4358, D17S964, D19S1091, D17S1179, D10S2160, D17S1230, D17S1338, D17S2011, D17S1237, D17S2038, D17S2091, D17S649, D17S1190 and M87506.

10. The method of claim 8 wherein the marker is selected from the group consisting of the SNPs:

rs2230698, rs2230700, rs1058808, rs1801200, rs903506, rs2313170, rs1136201, rs2934968, rs2172826, rs1810132, rs1801201, rs2230702, rs2230701, rs1126503, rs3471, rs13695, rs471692, rs558068, rs1064288, rs1061692, rs520630, rs782774, rs565121, rs2586112, rs532299, rs2732786, rs1804539, rs1804538, rs1804537, rs1141364, rs12231, rs1132259, rs1132257, rs113225, rs113225, rs1132254, rs113225, rs1132268 and rs11322

11. A method for the prediction, diagnosis or prognosis of malignant neoplasia by the detection of at least one marker **characterized in that** the marker is selected from:

a) a polynucleotide or polynucleotide analog comprising at least one of the sequences of SEQ ID NO: 2 to 6, 8, 9, 11 to 16, 18, 19, 21 to 26 or 53 to 75 ;

b) a polynucleotide or polynucleotide analog which hybridizes under stringent conditions to a polynucleotide specified in (a) and encodes a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 2 or 3

c) a polynucleotide or polynucleotide analog the sequence of which deviates from the polynucleotide specified in (a) and (c) due to the generation of the genetic code encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 2 or 3

d) a polynucleotide or polynucleotide analog which represents a specific fragment, derivative or allelic variation of a polynucleotide sequence specified in (a) to (d)

e) a purified polypeptide encoded by a polynucleotide or polynucleotide analog sequence specified in (a) to (e)

f) a purified polypeptide comprising at least one of the sequences of SEQ ID NO: 28 to 32, 34, 35, 37 to 42, 44, 45, 47 to 52 or 76 to 98;

are detected.

12. A method for the prediction, diagnosis or prognosis of malignant neoplasia by the detection of at least 2 markers

characterized in that at least 2 markers are selected from:

a) a polynucleotide or polynucleotide analog comprising at least one of the sequences of SEQ ID NO: 1 to 26 or 53 to 75;

b) a polynucleotide or polynucleotide analog which hybridizes under stringent conditions to a polynucleotide specified in (a) and encodes a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 2 or 3

c) a polynucleotide or polynucleotide analog the sequence of which deviates from the polynucleotide specified in (a) and (b) due to the generation of the genetic code encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 2 or 3

d) a polynucleotide or polynucleotide analog which represents a specific fragment, derivative or allelic variation of a polynucleotide sequence specified in (a) to (c)

e) a purified polypeptide encoded by a polynucleotide sequence or polynucleotide analog specified in (a) to (d)

f) a purified polypeptide comprising at least one of the sequences of SEQ ID NO: 27 to 52 or 76 to 98

are detected.

13. The method of any of the claims 1 or 12 wherein the detection method comprises the use of PCR, arrays or beads.

14. A diagnostic kit comprising instructions for conducting the method of any of claims 1 to 13.

15. A composition for the prediction, diagnosis or prognosis of malignant neoplasia comprising:

a) a detection agent for:

i) any polynucleotide or polynucleotide analog comprising at least one of the sequences of SEQ ID NO: 2 to 6, 8, 9, 11 to 16, 18,19,21 to 26 or 53 to 75;

ii) any polynucleotide or polynucleotide analog which hybridizes under stringent conditions to a polynucleotide specified in (a) encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 2 or 3

iii) a polynucleotide or polynucleotide analog the sequence of which deviates from the polynucleotide specified in (a) and (b) due to the generation of the genetic code encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 2 or 3

iv) a polynucleotide or polynucleotide analog which represents a specific fragment, derivative or allelic variation of a polynucleotide sequence specified in (a) to (c)

v) a polypeptide encoded by a polynucleotide or polynucleotide analog sequence specified in (a) to (d);

vi) a polypeptide comprising at least one of the sequences of SEQ ID NO: 28 to 32, 34, 35, 37 to 42, 44, 45, 47 to 52 or 76 to 98.

or

b) at least 2 detection agents for at least 2 markers selected from:

i) any polynucleotide comprising at least one of the sequences of SEQ ID NO: 1 to 26 or 53 to 75;

ii) any polynucleotide which hybridizes under stringent conditions to a polynucleotide specified in (a) encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 2 or 3

iii) a polynucleotide the sequence of which deviates from the polynucleotide specified in (a) and (b) due to the generation of the genetic code encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 2 or 3

iv) a polynucleotide which represents a specific fragment, derivative or allelic variation of a polynucleotide sequence specified in (a) to (c)

v) a polypeptide encoded by a polynucleotide sequence specified in (a) to (d);

vi) a polypeptide comprising at least one of the sequences of SEQ ID NO: 27 to 52 or 76 to 98.

16. An array comprising a plurality of polynucleotides or polynucleotide analogs wherein each of the polynucleotides is selected from:

a) a polynucleotide or polynucleotide analog comprising at least one of the sequences of SEQ ID NO: 1 to 26 or 53 to 75;

b) a polynucleotide or polynucleotide analog which hybridizes under stringent conditions to a polynucleotide specified in (a) encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 2 or 3

c) a polynucleotide or polynucleotide analog the sequence of which deviates from the polynucleotide specified in (a) and (b) due to the generation of the genetic code encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 2 or 3

d) a polynucleotide or polynucleotide analog which represents a specific fragment, derivative or allelic variation of a polynucleotide sequence specified in (a) to (c)

attached to a solid support.

17. A method of screening for agents which regulate the activity of a polypeptide encoded by a polynucleotide or polynucleotide analog selected from the group consisting of:

a) a polynucleotide or polynucleotide analog comprising at least one of the sequences of SEQ ID NO: 2 to 6, 8, 9, 11 to 16, 18, 19, 21 to 26 or 53 to 75;

b) a polynucleotide or polynucleotide analog which hybridizes under stringent conditions to a polynucleotide specified in (a) encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 2 or 3

c) a polynucleotide or polynucleotide analog the sequence of which deviates from the polynucleotide specified in (a) and (b) due to the generation of the genetic code encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 2 or 3

d) a polynucleotide or polynucleotide analog which represents a specific fragment, derivative or allelic variation of a polynucleotide sequence specified in (a) to (c);

comprising the steps of:

i) contacting a test compound with at least one polypeptide encoded by a polynucleotide specified in (a) to (d); and

ii) detecting binding of the test compound to the polypeptide, wherein a test compound which binds to the polypeptide is identified as a potential therapeutic agent for modulating the activity of the polypeptide in order to prevent or treat malignant neoplasia.

18. A method of screening for agents which regulate the activity of a polypeptide encoded by a polynucleotide or polynucleotide analog selected from the group consisting of:

a) a polynucleotide or polynucleotide analog comprising at least one of the sequences of SEQ ID NO: 2 to 6, 8, 9, 11 to 16, 18, 19, 21 to 26 or 53 to 75;

b) a polynucleotide or polynucleotide analog which hybridizes under stringent conditions to a polynucleotide specified in (a) encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 2 or 3

c) a polynucleotide or polynucleotide analog the sequence of which deviates from the polynucleotide specified in (a) and (b) due to the generation of the genetic code encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 2 or 3

d) a polynucleotide or polynucleotide analog which represents a specific fragment, derivative or allelic variation of a polynucleotide sequence specified in (a) to (c)

comprising the steps of:

i) contacting a test compound with at least one polypeptide encoded by a polynucleotide specified in (a) to (d); and

ii) detecting the activity of the polypeptide as specified for the respective sequence in Table 2 or 3, wherein a test compound which increases the activity is identified as a potential preventive or therapeutic agent for increasing the polypeptide activity in malignant neoplasia, and wherein a test compound which decreases the activity of the polypeptide is identified as a potential therapeutic agent for decreasing the polypeptide activity in malignant neoplasia.

19. A method of screening for agents which regulate the activity of a polynucleotide or polynucleotide analog selected from group consisting of;

a) a polynucleotide or polynucleotide analog comprising at least one of the sequences of SEQ ID NO: 2 to 6, 8, 9, 11 to 16, 18, 19, 21 to 26 or 53 to 75;

b) a polynucleotide or polynucleotide analog which hybridizes under stringent conditions to a polynucleotide specified in (a) encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 2 or 3

c) a polynucleotide or polynucleotide analog the sequence of which deviates from the polynucleotide specified in (a) and (b) due to the generation of the genetic code encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 2 or 3

d) a polynucleotide or polynucleotide analog which represents a specific fragment, derivative or allelic variation of a polynucleotide sequence specified in (a) to (c)

comprising the steps of:

i) contacting a test compound with at least one polynucleotide or polynucleotide analog specified in (a) to (d), and

ii) detecting binding of the test compound to the polynucleotide, wherein a test compound which binds to the polynucleotide is identified as a potential preventive or therapeutic agent for regulating the activity of the polynucleotide in malignant neoplasia.

20. Use of

a) a polynucleotide or polynucleotide analog comprising at least one of the sequences of SEQ ID NO: 2 to 6, 8, 9, 11 to 16, 18, 19, 21 to 26 or 53 to 75;

b) a polynucleotide which hybridizes under stringent conditions to a polynucleotide or polynucleotide analog specified in (a) encoding a polypeptide exhibiting the same biological function as specified for the respective

sequence in Table 2 or 3;

c) a polynucleotide or polynucleotide analog the sequence of which deviates from the polynucleotide specified in (a) and (b) due to the generation of the genetic code encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 2 or 3;

d) a polynucleotide or polynucleotide analog which represents a specific fragment, derivative or allelic variation of a polynucleotide sequence specified in (a) to (c);

e) an antisense molecule targeting specifically one of the polynucleotide sequences specified in (a) to (d);

f) a purified polypeptide encoded by a polynucleotide or polynucleotide analog sequence specified in (a) to (d)

g) a purified polypeptide comprising at least one of the sequences of SEQ ID NO: 28 to 32, 34, 35, 37 to 42, 44, 45, 47 to 52 or 76 to 98;

h) an antibody capable of binding to one of the polynucleotide specified in (a) to (d) or a polypeptide specified in (f) and (g);

i) a reagent identified by any of the methods of claim 17 to 19 that modulates the amount or activity of a polynucleotide sequence specified in (a) to (d) or a polypeptide specified in (f) and (g);

in the preparation of a composition for the prevention, prediction, diagnosis, prognosis or a medicament for the treatment of malignant neoplasia.

21. Use of claim 20 wherein the disease is breast cancer.

22. A reagent that regulates the activity of a polypeptide selected from the group consisting of:

a) a polypeptide encoded by any polynucleotide or polynucleotide analog comprising at least one of the sequences of SEQ ID NO: 2 to 6, 8, 9, 11 to 16, 18, 19, 21 to 26 or 53 to 75;

b) a polypeptide encoded by any polynucleotide or polynucleotide analog which hybridizes under stringent conditions to any polynucleotide comprising at least one of the sequences of SEQ ID NO: 2 to 6, 8, 9, 11 to 16, 18, 19, 21 to 26 or 53 to 75 encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 2 or 3

c) a polypeptide encoded by any polynucleotide or polynucleotide analog the sequence of which deviates from the polynucleotide specified in (a) and (b) due to the generation of the genetic code encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 2 or 3

d) a polypeptide encoded by any polynucleotide or polynucleotide analog which represents a specific fragment, derivative or allelic variation of a polynucleotide sequence specified in (a) to (c) encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 2 or 3

e) or a polypeptide comprising at least one of the sequences of SEQ ID NO: 28 to 32, 34, 35, 37 to 42, 44, 45, 47 to 52 or 76 to 98;

wherein said reagent is identified by the method of any of the claims 17 to 19.

23. A reagent that regulates the activity of a polynucleotide or polynucleotide analog selected from the group consisting of:

a) a polynucleotide or polynucleotide analog comprising at least one of the sequences SEQ ID NO: 2 to 6, 8, 9, 11 to 16, 18, 19, 21 to 26 or 53 to 75;

b) a polynucleotide or polynucleotide analog which hybridizes under stringent conditions to a polynucleotide specified in (a) encoding a polypeptide exhibiting the same biological function as specified for the respective

sequence in Table 2 or 3

c) a polynucleotide or polynucleotide analog the sequence of which deviates from the polynucleotide specified in (a) and (b) due to the generation of the genetic code encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 2 or 3

d) a polynucleotide or polynucleotide analog which represents a specific fragment, derivative or allelic variation of a polynucleotide sequence specified in (a) to (c)_encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 2 or 3

wherein said reagent is identified by the method of any of the claims 17 to 19.

24. A pharmaceutical composition, comprising:

a) an expression vector containing at least one polynucleotide or polynucleotide analog selected from the group consisting of:

i) a polynucleotide or polynucleotide analog comprising at least one of the sequences of SEQ ID NO: 2 to 6, 8, 9, 11 to 16, 18, 19, 21 to 26 or 53 to 75;

ii) a polynucleotide or polynucleotide analog which hybridizes under stringent conditions to a polynucleotide specified in (a) encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 2 or 3

iii) a polynucleotide or polynucleotide analog the sequence of which deviates from the polynucleotide specified in (a) and (b) due to the generation of the genetic code encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 2 or 3

iv) a polynucleotide or polynucleotide analog which represents a specific fragment, derivative or allelic variation of a polynucleotide sequence specified in (a) to (c)_encoding a polypeptide exhibiting the same biological function as specified for the respective sequence in Table 2 or 3;

or the reagent of claim 22 or 23 and a pharmaceutically acceptable carrier.

25. A computer-readable medium comprising:

a) at least one digitally encoded value representing a level of expression of at least one polynucleotide sequence of SEQ ID NO: 2 to 6, 8, 9, 11 to 16, 18, 19, 21 to 26 or 53 to 75

b) at least 2 digitally encoded values representing the levels of expression of at least 2 polynucleotide sequences selected from SEQ ID NO: 1 to 26 or 53 to 75

in a cell from the a subject at risk for or having malignant neoplasia.

26. A method for the detection of chromosomal alterations **characterized in that** the relative abundance of individual mRNAs, encoded by genes, located in altered chromosomal regions is detected.

27. A method for the detection of chromosomal alterations **characterized in that** the copy number of one or more chromosomal region(s) is detected by quantitative PCR.

Figure 1




Figure 1




Figure 2




Figure 3




Figure 4

